Document Type: EA-Administrative Record Index Field: Environmental Assessment Project Name: Golden Triangle Solar II Project

Project Number: 2021-17

# GOLDEN TRIANGLE II SOLAR FACILITY and BATTERY ENERGY STORAGE SYSTEM PROJECT

# **Lowndes County, Mississippi**

## FINAL ENVIRONMENTAL ASSESSMENT

Prepared for: TENNESSEE VALLEY AUTHORITY Knoxville, Tennessee

> Submitted By: MS Solar 6, LLC

Prepared by: Burns & McDonnell

May 2022

To request further information, contact:
J. Taylor Johnson
NEPA Compliance
Tennessee Valley Authority
1101 Market Street, BR 2C-C
Chattanooga, Tennessee 37402
E-mail: jtcates@tva.gov

		<u>P</u> :	<u>age No.</u>
1.0	INTR	RODUCTION	1-1
	1.1	Purpose and Need for Action	
	1.2	Scope of this Environmental Assessment	
	1.3	Public Involvement	
	1.4	Necessary Permits or Licenses	
		1.4.1 Clean Water Act and Rivers and Harbors Act	
		1.4.2 Endangered Species Act of 1973	
		1.4.3 National Historic Preservation Act	1-7
		1.4.4 Mississippi Natural Heritage Program	1-7
		1.4.5 Mississippi Department of Environmental Quality	
		1.4.6 County and Municipality Permitting	1-8
2.0	DES	CRIPTION OF PROPOSED ALTERNATIVES	2-1
	2.1	No Action Alternative	
	2.2	Proposed Action Alternative	
		2.2.1 Project Description	
		2.2.2 Solar Facility Construction	
		2.2.3 Operations	
		2.2.4 Decommissioning and Reclamation	2-8
	2.3	Alternatives Eliminated from Further Consideration	2-8
	2.4	Comparison of Alternatives	2-9
	2.5	Best Management Practices and Mitigation Measures	
		2.5.1 Golden Triangle II Solar Facility	2-12
	2.6	The Preferred Alternative	2-14
3.0	AFF	ECTED ENVIRONMENT AND ENVIRONMENTAL CONSEQUENCE	S3-1
	3.1	Land Use	3-1
		3.1.1 Affected Environment	_
		3.1.2 Environmental Consequences	
	3.2	Geology, Soils, and Prime Farmland	3-1
		3.2.1 Affected Environment	
		3.2.2 Environmental Consequences	
	3.3	Water Resources	
		3.3.1 Affected Environment	
		3.3.2 Environmental Consequences	
	3.4	Biological Resources	
		3.4.1 Affected Environment	
		3.4.2 Environmental Consequences – Biological Resources	
	3.5	Visual Resources	
		3.5.1 Affected Environment	_
	0.0	3.5.2 Environmental Consequences	
	3.6	Noise	_
		3.6.1 Affected Environment	
		3.6.2 Environmental Consequences - Noise	3-19

	3.7	Air Qua	lity and Greenhouse Gas Emissions	
		3.7.1	Affected Environment	
		3.7.2	Environmental Consequences	3-22
	3.8	Cultural	Resources	3-24
		3.8.1	Affected Environment	3-25
		3.8.2	Environmental Consequences - Cultural	3-29
	3.9	Utilities.	'	
		3.9.1	Affected Environment	3-32
		3.9.2	Environmental Consequences - Utilities	
	3.10	Waste N	Management	
		3.10.1	Affected Environment	
		3.10.2	Environmental Consequences	
	3.11		and Occupational Health and Safety	
		3.11.1	Affected Environment	
		3.11.2	Environmental Consequences	
	3.12	_	conomics	
	· · · -	3.12.1	Affected Environment	
		3.12.2		
	3.13	-	mental Justice	3-45
		3.13.1	Affected Environment	
		3.13.2	Environmental Consequences	
	3.14	Transpo	ortation	
			Affected Environment	
		3.14.2	Environmental Consequences	3-47
			'	
4.0	ΔΝΤΙ	IDATEC	ENVIRONMENTAL IMPACTS AND CUMULATIVE	
4.0		CTS		<b>⊿</b> _1
	4.1	_	dable Adverse Environmental Impacts	
	4.2		nship of Short-Term Uses and Long-Term Productivity	
	4.3	Irrevers	ible and Irretrievable Commitments of Resources	4-3
	4.4		tive Impacts	_
	7.7	4.4.1	Federal Projects	
			State and Local Projects	
		<b>⊤.</b> ⊤.∠	Otate and Local Projects	
г o	LICT	عد ممدر	ADEDO	<b>5</b> 4
5.0	LIST	JF PREI	PARERS	5-1
6.0	LITER	RATURE	CITED	6-1
APPE	NDIX A	A TVA'S	SITE CLEARING AND GRADING SPECIFICATIONS	
APPE	NDIX E	3 WETLA	AND AND WATERBODY DELINEATION REPORT	
APPE	NDIX (	C ONSIT	E AND OFFSITE DRAINAGE REPORT	
APPE	NDIX [	D PROTI	ECTED SPECIES INFORMATION AND REPORTING	
APPE	NDIX E	E AGEN	CY CONSULTATIONS	
APPE	NDIX F	СОММ	ENTS RECEIVED DURING THE PUBLIC SCOPING	
		PER	IOD (FEBRUARY 18 – MARCH 20, 2022)	

## **LIST OF TABLES**

			Page No.
Table	1-1:	Golden Triangle II Solar Project Permit and Approval List	1-5
Table	2-1:	Comparison of Impacts by Alternative	
Table	3-1:	Land Use and Percent Cover within the Project Site	
Table	3-2:	Soils within the Project Site	
Table	3-3:	Waterbodies Potentially Affected by the Golden Triangle II Solar	
		Project Access Road Crossings	
Table	3-4:	Protected Species with Potential to Occur within the Survey Area.	3-8
Table	3-5:	TVA Natural Heritage Database - Plants Potentially Occurring	
		within the Survey Area	3-5
Table	3-6:	TVA Natural Heritage Database – Aquatic Species Potentially	
		Occurring within 10-miles of Survey Area.	3-6
Table	3-7:	Typical Sound Pressure Levels Associated with Common Sound	
		Sources	3-18
Table		Average Emissions of NAAQS Pollutants in Lowndes County	
		Previously Recorded Archaeological Sites within the Study Area	
		Previous Archaeological Surveys within the Project Study Area	
		Waste Facilities Near the Project Site	3-35
Table	3-12:	Summary of Special Handling Precautions for Large Quantity	
		Hazardous Materials	3-37
Table	3-13:	Summary of Construction Waste Streams and Management	
		Methods	3-38
Table	3-14:	Summary of Operational Waste Streams and Management	
		Methods	3-39
Table	4-1:	Unavoidable Impacts and Proposed Mitigation for the Golden	
		Triangle II Solar Project	
Table	4-2:	Potential Cumulative Impacts Associated with the Golden Triangle	
		Il Solar Project	4-1
Table	5-1:	Golden Triangle II Solar Project Environmental Assessment Team	5-1

# **LIST OF FIGURES**

			Page No.
Figure	1-1:	Golden Triangle II Solar Project Location Map	1-3
		Aerial Map of the Golden Triangle II Solar Project Site	
		Aerial Map of the Preliminary Golden Triangle II Solar Project	
Ü		Layout	2-2
Figure	2-3:	PV Solar System Energy Flow Diagram	
_		Land Cover within the Project Site	
Figure	3-2:	Soils within the Project Site	3-1
Figure	3-3:	Prime Farmland Soils within the Project Site	3-1
Figure	3-4:	Floodplains within the Project Site	3-1
Figure	3-5:	Surface Water and Wetlands within the Project Site	3-1
Figure	3-6:	Chalk Outcrop on Parcel Associated with the Project	3-4
Figure	3-7:	Potentially Suitable Habitat for Federally Protected Species within	
_		the Project Site	3-1
Figure	3-8:	General Site Photo of Golden Triangle II Project Site	3-13
Figure	3-9:	General Site Photo of Golden Triangle II Project Site	3-14
Figure	3-10	:Representative Photo of Single-Axis Tracking PV System in the	
_		Morning, Facing East	3-15
Figure	3-11	:Representative Photo of Single-Axis Tracking PV System at	
_		Midday, Laying Flat	3-16
Figure	3-12	:Observed and Projected Temperature Change in Mississippi	
Figure	3-13	:Solar Glare Ocular Hazard Plot	3-41
Figure	4-1:	Cumulative Impact Map 4-1	

#### LIST OF ABBREVIATIONS

<u>Abbreviation</u> <u>Term/Phrase/Name</u>

AADT Annual Average Daily Traffic

AC alternating current

Alp Airport Layout Plan

APE Area of Potential Effects

BCC Birds of conservation concern

BESS battery energy storage system

BFE base flood elevation

BGEPA Bald and Golden Eagle Protection Act

BMP best management practice

BMS Battery management system

BZA Board of Zoning Appeals

CBMPP Construction best management practices plan

CEQ Council on Environmental Quality

CFR Code of Federal Register

CLOMR Conditional Letter of Map Revision

CO Carbon monoxide

CRA Cultural Resource Analysts

CWA Clean Water Act

dBA A-weighted decibel

dbh diameter at breast height

DC direct current

DNL day-night sound level

EA Environmental Assessment

EO Executive Order

ESCP Erosion and Sedimentation Control Plan

FAA Federal Aviation Administration

F-A-R Forestry-Agriculture-Recreation

<u>Abbreviation</u> <u>Term/Phrase/Name</u>

FEMA Federal Emergency Management Agency

FIRM Flood Insurance Rate Map

FPPA Farmland Protection Policy Act

GHG greenhouse gas

GT2 Golden Triangle II Solar Project
GTR Golden Triangle Regional Airport

HVAC Heating, ventilation, and air conditioning

IRP Integrated Resource Plan

kV kilovolts

L<sub>dn</sub> Day-night sound level

L<sub>eq</sub> 24-hour equivalent sound level

LFP Lithium iron phosphate

Li-ion Lithium ion

Lmax maximum noise level

LTO Lithium titanate oxide

μg/m³ micrograms per cubic meter

MBTA Migratory Bird Treaty Act

MDAH Mississippi Department of Archives and History

MDEQ Mississippi Department of Environmental Quality

MDOT Mississippi Department of Transportation

MDWFP Mississippi Department of Wildlife, Fisheries, and Parks

MFC Mississippi Forestry Commission

MLRA Major Land Resource Area

MPT main power transformer

MS Solar 6 MS Solar 6, LLC

MSSHPO Mississippi State Historic Preservation Office

MVT mid-voltage transformer

MW megawatt

Abbreviation Term/Phrase/Name

NAAQS National Ambient Air Quality Standards

NCM Lithium nickel cobalt manganese oxide

NEPA National Environmental Policy Act

NFPA National Fire Protection Association

NHL National Historic Landmark

NHPA National Historic Preservation Act

NO<sub>2</sub> nitrogen dioxide

NOI notice of intent

NPDES National Pollution Discharge Elimination System

NPS National Park Service

NRCS Natural Resources Conservation Service

NRHP National Register of Historic Places

NSA Noise sensitive areas

OSHA Occupational Safety and Health Administration

PEL Permissible exposure limit

PM<sub>10</sub> Particulate matter having a diameter of less than or equal to 10

microns

PM<sub>2.5</sub> Particulate matter having a diameter of less than or equal to 2.5

microns

PPA power purchase agreement

ppb parts per billion

ppm parts per million

Project The Golden Triangle II Solar Project

PV photovoltaic

ROW right-of-way

RNHD regional natural heritage database

RSO Renewable Standard Offer

SGHAT Solar Glare Hazard Analysis Tool

SHPO State Historic Preservation Officer

Abbreviation Term/Phrase/Name

SO2 sulfur dioxide

Solar Facility A 150-MW AC single-axis tracking photovoltaic solar facility

SPCC Spill Prevention, Control, and Countermeasures

STEL Short-term exposure limit

SWPPP Stormwater Pollution Prevention Plan

TLV Threshold limit value

TVA Tennessee Valley Authority

TVARAM Tennessee Valley Authority Rapid Assessment Method

TWA Time weighted average

USACE U.S. Army Corps of Engineers

USCB U.S. Census Bureau

USDA U.S. Department of Agriculture

USEPA U.S. Environmental Protection Agency

USFWS U.S. Fish and Wildlife Service

USGS U.S. Geological Service

WOTUS Waters of the U.S.

#### **GLOSSARY OF TERMS**

- Area of Potential Effect (APE): The APE includes any area within the Project Site where impacts could occur during construction, operation, or both. Because the Project layout is still preliminary and subject to change, MS Solar 6 has selected an APE that is larger than what would likely be needed. The APE for cultural resources also includes areas within a half mile radius that are within the visual line of sight of the parcels where above ground facilities are proposed.
- **Artesia Substation:** The Artesia Substation is TVA's existing 161-kV substation, located along the western end of Mims Road.
- **Artesia Switching Station:** The Artesia Switching Station is TVA's proposed switching station for the Project. TVA's proposed 0.85-acre Artesia Switching Station would be located directly east of the existing TVA Artesia Substation.
- **Collection Lines:** Collection lines are typically buried (at least three feet under the surface) electrical connections that are installed between different sections of arrays and other facilities within the Project Site.
- **Fencerow:** A fencerow is typically located along the perimeter of a parcel that is comprised of agricultural land, pastureland, or open space. It may actually contain a fence or did at one point in the past. There is typically a row of trees or shrubs that grow along the fencerow. Many of the parcels associated with the Golden Triangle Solar Facility have fencerows along the perimeter of the parcels.
- **Gen-Tie:** Approximately 1,665-foot-long tie-in that would connect the proposed Golden Triangle I Substation to TVA's proposed Artesia Switching Station.
- Golden Triangle I Solar and Battery Energy Storage System (BESS) Facility: The Golden Triangle I Solar and BESS Facility was developed by MS Solar 5, LLC under a separate PPA with TVA. It resulted in 200 MW of AC generating capacity and an additional 50 MW of battery energy storage.
- Golden Triangle II Solar and Battery Energy Storage System (BESS) Facility: The Golden Triangle II Solar and BESS Facility is proposed by MS Solar 6, LLC. It would result in 150 MW of AC generating capacity and an additional 50 MW of battery energy storage. The Golden Triangle II BESS would be collocated with the BESS for Golden Triangle I.
- Golden Triangle I Substation: The Golden Triangle I Substation was developed by MS Solar 5, LLC as part of the Golden Triangle I Solar and BESS Project. The Golden Triangle II Solar and BESS Facility would connect to the Golden Triangle substation by way of collection lines. A separate and new substation will not be developed as part of this Project.
- **Project or Proposed Action (Project or GT2):** The Project or Proposed Action includes the proposed Golden Triangle II Solar and Battery Energy Storage System (Solar Facility), collection lines, access roads, and the PPA between TVA and MS Solar 6, LLC.
- **Project Area:** The "Project Area" includes the Project Site, as defined below, and the land, roadways, businesses, and homes within approximately ½ mile of the Project Site.

- **Project Site:** The "Project Site" includes all six parcels that would be affected during Project construction and/or operation. The total Project Site is approximately 1,524 acres. Only portions of the total Project Site would be impacted during construction and/or operation of the Project.
- **Shared Space/Area:** This is an approximately 150-acre area of land that was initially identified for placement of PV arrays for Golden Triangle II, that also may contain some PV arrays for Golden Triangle I (only in the event that additional capacity is needed to achieve the 200 MW of AC capacity.
- **Survey Area:** The Survey Area includes the proposed Project Site as well as additional areas that were surveyed by environmental and cultural resources specialists. The Survey Area applies to all parcels associated with the Solar Facility. Slightly more than 1,524 acres were surveyed as part of the environmental and cultural surveys. Visual surveys expanded beyond the limits of disturbance by approximately ½ mile for historic resources and architecture.

#### 1.0 INTRODUCTION

The Tennessee Valley Authority (TVA) entered into a Power Purchase Agreement (PPA) with MS Solar 6, LLC (herein referred to as "MS Solar 6") to purchase power generated by the proposed Golden Triangle II Solar and Battery Energy Storage System (BESS) Facility (Solar Facility) Project (GT2 or Project) in Lowndes County, Mississippi, subject to satisfactory completion of all environmental reviews. The Proposed Action of constructing the Golden Triangle II Solar and BESS Project (GT2 or Project) by MS Solar 6 would operate separately, under a separate PPA, from the previously evaluated Golden Triangle I Solar and BESS Facility Project. GT2 is expected to generate up to 150 megawatts (MW) of alternating current (AC) capacity with a 50 MW AC – 200-megawatt hour (MWh) BESS. Under the terms of the conditional PPA between TVA and MS Solar 6, dated December 15, 2020, TVA would purchase the electric output generated by the proposed Solar Facility for an initial term of 20 years, subject to satisfactory completion of all applicable environmental reviews.

The proposed a GT2 is located south and east of the town of Artesia, Mississippi (Figure 1-1). The Project Site includes six individual parcels on approximately 1,524 acres of land, on which the solar array footprint, connection lines, and associated access roads would be constructed. The Solar Facility would consist of multiple parallel rows of photovoltaic (PV) panes on single-axis tracking structures, along with direct current (DC) and AC inverters and transformers. MS Solar 6 would enter into long-term lease agreements or land purchases for all of the Project parcels.

TVA's connection with the Project would occur at the existing TVA Artesia Substation, located approximately three miles north of GT2. Network upgrades to TVA's existing system are not anticipated as a result of the additional capacity to be generated by GT2.

## 1.1 Purpose and Need for Action

The TVA is a corporate agency of the United States that provides electricity for business customers and local power companies serving nearly 10 million people in parts of seven southeastern states called the Tennessee Valley. TVA's mission is to serve the people of the Tennessee Valley region, and it does that through three main areas of work – energy, the environment, and economic development.

TVA produces or obtains electricity from a diverse portfolio of energy sources, including solar, hydroelectric, wind, biomass, fossil fuel, and nuclear. In June 2019, TVA completed an Integrated Resource Plan (IRP) and associated Environmental Impact Statement (EIS) (TVA 2019a and TVA 2019b). The IRP identified the various resources that TVA intends to use to meet the energy needs of the TVA region over the 20-year planning period while achieving TVA's objectives to deliver reliable, low-cost, and cleaner energy and reducing environmental impacts. These energy resources from the 2019 IRP included the addition of between 1,500 and 8,000 MW (AC) of solar capacity by 2028 and up to 14,000 MW by 2038 (TVA 2019a). Customer demand for cleaner energy prompted TVA to release a Request for Proposal (RFP) for renewable energy resources (2020 Renewable RFP). The MS Solar 6 PPA that resulted from this RFP will help TVA meet

immediate needs for additional renewable generating capacity in response to customer demands and fulfill the renewable energy goals established in the 2019 IRP. The Proposed Action would provide cost-effective renewable energy consistent with the IRP and TVA goals.

## 1.2 Scope of this Environmental Assessment

Pursuant to the National Environmental Policy Act of 1969 (NEPA), federal agencies are required to evaluate the potential environmental impacts of their proposed actions. This environmental assessment (EA) was prepared consistent with TVA regulations at 18 CFR 1318 issued in 2020 (85 FR 17434, Mar. 27, 2020), and is consistent with Council on Environmental Quality (CEQ) regulations for implementing NEPA at 40 CFR 1500-1508, which were recently revised (87 FR 23453, April 20, 2022) to amend certain provisions and reinstate long-standing provisions from the 1978 regulations (43 FR 55990, Nov. 29, 1978).

TVA's Proposed Action would result in the construction and operation of the proposed Project by MS Solar 6. The scope of this EA therefore focuses on impacts related to the construction and operation of the proposed Solar Facility.

This EA describes the existing environment at the Project Site (Figure 1-1), analyzes potential environmental impacts associated with the Proposed Action and the No Action Alternatives, and identifies and characterizes potential cumulative impacts from the proposed Project in relation to other recent, ongoing, and reasonably foreseeable future proposed activities within the surrounding area of the Project Site.

Under the PPA, TVA's obligation to purchase power is contingent upon the satisfactory completion of the appropriate environmental review and TVA's determination that the Proposed Action will be "environmentally acceptable." To be deemed "environmentally acceptable", TVA must assess the impact of the Project on the human environment to determine whether (1) any significant impacts would result from the location, operation, and/or maintenance of the proposed Project and/or associated facilities, and (2) the Project would be consistent with the purposes, provisions, and requirements of applicable federal, state, and local environmental laws and regulations.

Based on internal scoping and identification of applicable laws, regulations, executive orders, and policies, TVA identified the following resource areas necessary for analysis within this EA: Land Use (includes Natural Areas and Recreation); Geology, Soils, and Prime Farmland; Water Resources; Biological Resources; Visual Resources; Noise; Air Quality and Greenhouse Gas Emissions; Cultural Resources; Utilities; Waste Management; Public and Occupational Health and Safety; Socioeconomics; Environmental Justice; and Transportation.

This EA consists of six chapters discussing the Project alternatives, potentially impacted resource areas, and analyses of these impacts. Additionally, this document includes five appendices, which generally contain more detail on technical analyses and supporting data. The structure of the EA is outlined below:

- **Section 1.0**: Describes the purpose and need for the Project, the decision to be made, related environmental reviews and consultation requirements, necessary permits or licenses, and the EA overview.
- **Section 2.0**: Describes the Proposed Action and No Action Alternatives, provides a comparison of alternatives, and discusses the Preferred Alternative.
- **Section 3.0**: Discusses the affected environment and the potential direct and indirect impacts on these resource areas. Mitigation measures are also proposed, as appropriate.
- **Section 4.0**: Discusses the cumulative impacts in relation to other ongoing and reasonably foreseeable proposed activities within the surrounding area of the Project Site.
- **Section 5.0**: Provides the List of Preparers for the preparation of this EA.
- **Section 6.0:** Provides the list of Literature Cited in this EA.
- Appendix A: TVA's Site Clearing and Grading Specifications
- Appendix B: GT2 Wetland and Waterbody Delineation Report
- Appendix C: GT2 Solar Project Hydrology Study
- Appendix D: Protected Species Information and Reporting for GT2
- Appendix E: Agency Consultations

Golden Triangle II Solar Project Introduction




Figure 1-1: GT2 Location Map, Lowndes County, MS

#### 1.3 Public Involvement

A copy of the draft EA was sent to local, state, and federal agencies and individuals who indicated an interest in the Project. TVA notified interested federally recognized Native American Tribes, elected officials, and other stakeholders that the draft EA was available for review and comment for a 30-day period. An electronic version of the document was posted on the TVA website where comments could also be submitted electronically. Public notices were locally published to solicit comments from other agencies, the general public, and any interested organizations. Federal agencies that received the notification consisted of the United States Department of Agriculture (USDA) Natural Resource Conservation Service (NRCS), the United States Army Corps of Engineers (USACE), and the U.S. Fish and Wildlife Service (USFWS). State and local agencies that received the notification consisted of the Mississippi Department of Agriculture, Mississippi Department of Economic and Community Development, Mississippi Department of Environmental Quality (MDEQ), and Mississippi Wildlife Department of Wildlife and Fisheries (MDWF). Tribes that received notification of the Project consisted of the Eastern Shawnee Tribe of Oklahoma, Choctaw Nation of Oklahoma, The Chickasaw Nation, Thlopthlocco Tribal Town, Kialegee Tribal Town, Absentee Shawnee Tribe of Indians of Oklahoma, Coushatta Tribe of Louisiana, Muscogee (Creek) Nation, Alabama-Coushatta Tribe of Texas, Mississippi Band of Choctaw Indians, Jena Band of Choctaw Indians, and the Shawnee Tribe.

During the 30-day public review and comment period of the draft EA, one comment was received from the U.S. Environmental Protection Agency (EPA). A copy is included in Appendix F. No additional comments were received from the general public or other interested agencies.

## 1.4 Necessary Permits or Licenses

There are multiple permits, approvals, registrations, and consultations that would be required for the Project. Table 1-1, below, provides an overview of the anticipated permits and approvals. Further details on anticipated permits and approvals are provided in the following sections.

Permit/Approval **Associated Documentation Lead Agency** Federal Permits, Approvals, Registrations, or Consultations **Endangered Species Act** Protected species Habitat U.S. Fish and Wildlife Service Section 7 (ESA) Assessment Report. Informal (USFWS) Section 7 ESA consultation Migratory Bird Treaty Act package. (MBTA) Bald and Golden Eagle Protection Act (BGEPA) None. FPPA applies to Projects Natural Resources Farmland Protection Policy Act receiving federal funding. The Conservation Service (FPPA) Solar Facility would be funded by (NRCS) MS Solar 6.

Table 1-1: Golden Triangle II Solar Project Permit and Approval List

Permit/Approval	Associated Documentation	Lead Agency		
Section 404 Clean Water Act Nationwide Permits 3 and/or 14 for access road crossings	Wetland and Waterbody Delineation Report	U.S. Army Corps of Engineers		
State Permits, Approvals, Regis	strations, or Consultations			
Section106 National Historical Preservation Act Consultation	Phase I Cultural Resources Survey Report	Mississippi Department of Archives and History / State Historic Preservation Officer		
§401 Clean Water Act – Water Quality Certification	Wetland and Waterbody Delineation Report	Mississippi Department of Environmental Quality (MDEQ)		
Construction General Permit No. MSR10 under the National Pollutant Discharge Elimination System (NPDES)	Stormwater Pollution Prevention Plan and Spill Prevention, Control, and Countermeasures Plan.	MDEQ		
Natural Heritage Program Consultation	Protected Species Habitat Assessment Report	MDEQ, Dept. of Wildlife, Fisheries, and Parks		
Driveway/Overhead Electrical Encroachment Permit	Site Plan/Driveway Access Plan	Mississippi Department of Transportation (MDOT)		
Local Permits, Approvals, Registrations, or Consultations				
Floodplain Development Permit	Individual Elevation Certificates for each tracker array in floodplain	Lowndes County Building Inspection Department		
Site Plan Permit	Construction Drawings, MDEQ NPDES Approval	Lowndes County Buildings and Grounds Office		
Building Permit	Electrical/Structural Plans	Lowndes County Buildings and Grounds Office		

#### 1.4.1 Clean Water Act and Rivers and Harbors Act

The U.S. Army Corps of Engineers (USACE) administers Section 404 of the Clean Water Act (CWA), which regulates dredge and fill activities in waters of the U.S. (33 USC 1344), as well as Section 10 of the Rivers and Harbors Act, which regulates the placement of structures in waters of the U.S. (33 USC 403). The Project is located within the Mobile District.

Impacts to jurisdictional waters of the U.S. (as defined in 40 CFR 230.3[s]) would require authorization from USACE under Section 404 of the CWA. Impacts resulting in a permanent loss of wetlands greater than 0.5 acre would require a Section 404 Individual Permit (IP), which may include mitigation and public involvement. The USACE established the Nationwide Permit (NWP) program to streamline the Section 404 permitting process for actions that would have no more than a minimal effect on the environment. If permanent wetland impacts (i.e., permanent fill resulting in the loss of wetland function) range from 0.1 to 0.5 acre, USACE notification and application is required to obtain an NWP. The Project would result in minimal impacts on wetlands, as discussed in greater detail in section 3.3 of this EA. Impacts on USACE-jurisdictional waterbodies would occur in areas where facility access roads and collection lines must cross

streams within the Project Site. Based on conversations with the Mobile District USACE, approval under an NWP 14 is anticipated.

## 1.4.2 Endangered Species Act of 1973

The Endangered Species Act of 1973 (ESA), as amended, was enacted to protect, and recover imperiled species and the ecosystems upon which they depend. The law requires federal agencies, in consultation with the U.S. Fish and Wildlife Service and/or the NOAA Fisheries Service, to ensure that actions they authorize, fund, or carry out are not likely to jeopardize the continued existence of any listed species or result in the destruction or adverse modification of designated critical habitat of such species. The law also prohibits any action that causes a "taking" of any listed species of endangered fish or wildlife. Further details regarding species listed as Threatened and Endangered (T&E) under the ESA are included in section 3.4.1.4.

#### 1.4.3 National Historic Preservation Act

Section 106 of the National Historic Preservation Act (NHPA) requires federal agencies to take into account the effects of their undertakings (including issuance of permits) on historic properties, and afford the Advisory Council on Historic Preservation a reasonable opportunity to comment. State Historic Preservation Officers (SHPOs) and Tribal Historic Preservation Officers (THPOs) serve a critical role in implementing many responsibilities under the NHPA. Central to this framework is the National Register of Historic Places (NRHP), which is the official list of historic properties worthy of preservation. The Mississippi Department of Archives and History (MDAH), which acts as the SHPO for the State of Mississippi, has reviewed and commented on potential Project impacts on resources of cultural significance within the state. A phase I cultural resources survey, utilizing pedestrian and shovel tests along pre-approved transects, was required for this Project. The SHPO does not issue permits; however, approvals by other federal agencies cannot be final without review by and clearance from the SHPO.

## 1.4.4 Mississippi Natural Heritage Program

The Mississippi-protected species list includes animal species for which legal protection is provided under the Mississippi Nongame and Endangered Species Conservation Act of 1974. The act declares that "Species or subspecies of wildlife indigenous to the state should be accorded protection in order to maintain and to the extent possible enhance their numbers." Under state law, the Mississippi Department of Environmental Quality (MDEQ), Department of Wildlife, Fisheries, and Parks has responsibility for reviewing this list and providing recommendations for revisions as needed every two years.

Mississippi law specifically states that rules and regulations related to the protection of state protected species will not affect rights on private property. Prohibitions are limited to the capture, killing, or selling of protected species and the protection of the habitat of these species on public lands.

## 1.4.5 Mississippi Department of Environmental Quality

The MDEQ Stormwater and 401 Water Quality Branch administers the Section 401 Water Quality Certification program in conjunction with the USACE. MDEQ offers general permits for activities that would result in only very minor impacts on wetlands. Proposed minor wetland alteration activities may obtain coverage by submitting a signed and completed application for a general permit, along with any other required information. Work shall not commence until a written Notice of Coverage (NOC) from MDEQ is received.

The MDEQ Stormwater and 401 Water Quality Branch also administers the National Pollution Discharge Elimination System (NPDES) construction stormwater permitting program in Mississippi. In compliance with the Mississippi Water Quality Control Act, MDEQ authorizes point source discharges of stormwater into waters of the state. A notice of intent (NOI) for General NPDES Permit for Stormwater Discharges from Large Construction Activities (permit no. MSR10) would be required for the Project. The NOI form, Stormwater Pollution Prevention Plan (SWPPP), and an Erosion, Sediment and Pollution Control Plan (ES&PCP) would be submitted at least 30 days prior to the commencement of land disturbing activities. Written approval through a NOC must first be received prior to initiating land disturbing activities. These documents will include specific information about the construction site, construction best management practices (BMPs), and stormwater discharge receiving waters.

## 1.4.6 County and Municipality Permitting

No ordinances or requirements specific to solar facilities exist in Lowndes County; however, Lowndes County does require a development review process. Based on initial coordination with the County, a building permit for the structural and electrical scope of the Project will be required. This review will be handled by the County Buildings and Inspection Department. The County will provide a site plan permit once the construction drawings are reviewed and approved by the County Board of Supervisors. A copy of the MDEQ NPDES approval must be provided to the County. Since portions of the Project would occur within a floodplain, MS Solar 6 would require a no rise/no impact and elevation certification. The County has coordinated with Mississippi Emergency Management Agency (MEMA) on what would be required. The County has indicated that it will require an individual elevation certificate for each tracker array located within the Federal Emergency Management Agency (FEMA) 100-year floodplain. The certificate must show that the elevation of the "drip line", or lowest point of the individual array at full tilt, is a minimum of one foot above the 100-year base flood elevation.

Vegetative waste from clearing activities would be burned or chipped onsite. If open burning of debris from tree clearing on the site is planned, the appropriate open burning permits would be obtained from the Mississippi Forestry Commission (MFC). Information on open or surface burning issued by MFC would be followed. Only trees and brush from the Project Site would be burned. Weather conditions would be monitored and considered to ensure safety and minimize degradation to air quality during the open burning of any vegetation cleared from the Project Site.

#### 2.0 DESCRIPTION OF PROPOSED ALTERNATIVES

This section provides a description of alternatives considered and compares the alternatives to the Proposed Action. This EA evaluates two alternatives: The No Action Alternative and the Proposed Action Alternative. A description of the analysis and criteria used in identifying the Preferred Alternative is described.

#### 2.1 No Action Alternative

The No Action Alternative provides a baseline of conditions against which the impacts of the Proposed Action Alternative are measured. Under the No Action Alternative, TVA would not purchase the power generated by the Project under the 20-year PPA with MS Solar 6 (i.e., TVA would not be involved with the Project). If TVA were to select this alternative and MS Solar 6 elected not to proceed with the Project, then MS Solar 6 would not construct or operate the Solar and BESS Facilities. Existing conditions (land use, natural resources, visual resources, physical resources, and socioeconomics) at the Project Site would remain unchanged. TVA would continue to rely on other sources of generation described in the 2019 IRP (TVA 2019a) to ensure an adequate energy supply and to meet its goals for increased renewable energy and low greenhouse gas (GHG)-emitting generation.

## 2.2 Proposed Action Alternative

Under the Proposed Action Alternative, MS Solar 6 would construct and operate an up to 150-MW AC single-axis tracking PV Solar and up to 50-MW – 200 MWh BESS Facility in Lowndes County, Mississippi, and TVA would purchase renewable energy from the facility under the 20-year PPA with MS Solar 6. GT2 would generate up to 150 MW AC output for transmission to TVA's electrical network. MS Solar 6 would enter into long-term leases or purchases on approximately 1,524 acres of land on six individual parcels along the southeastern town limits of Artesia, Mississippi. The power generated from the Solar Facility would be sold to TVA under the terms of the PPA. The Project would connect to the existing TVA electrical network via the existing Golden Triangle gen-tie line to TVA's proposed Artesia Switching Station within the existing Artesia Substation. Figure 2-1 provides an overview of the Project Site as well as the parcel boundaries of properties that would be affected during construction.

As discussed in Section 1.2, this EA assesses (1) the impact of TVA's action to enter into the PPA with MS Solar 6, (2) the associated impacts of the construction and operation of the Solar Facility by MS Solar 6, and (3) the interconnection components by TVA.

## 2.2.1 Project Description

The proposed Solar Facility would occupy portions of the Project Site that are predominantly comprised of cultivated agricultural fields and pastureland (Figure 2-1). The perimeter of the developed facilities would be enclosed with security fencing. Within the limits of the fenced facility would be the arrays of solar panels, inverters, electrical cabling, and other related infrastructure such as the access roads. The remaining portions of the Project Site would remain undeveloped. Additional information regarding existing land use conditions is detailed in Section 3. Figure 2-2 displays the preliminary solar array layout within the Project Site.

Golden Triangle II Solar Project



Figure 2-1: Aerial Map of the GT2 Solar Project Site




Figure 2-2: Aerial Map of the Preliminary GT2 Layout

GT2 would convert sunlight into direct current (DC) electrical energy within the PV panels (modules) as very generally depicted in Figure 2-3. PV power generation is the direct conversion of light into electricity at the atomic level. Some materials exhibit a property known as the photoelectric effect that causes them to absorb energy as photons of light and then release energy as electrons. When the free electrons are captured, an electric current is produced, which can be used as electricity (TVA 2014).




Figure 2-3: PV Solar System Energy Flow Diagram

The Project would be composed of PV modules mounted together in arrays. Groups of panels would be connected electrically in series to form "strings" of panels, with the maximum string size chosen to ensure that the maximum inverter input voltage is not exceeded by the string voltage at the Project's high design temperature. The panels would be in individual arrays consisting of the PV arrays and an inverter station on a concrete pad or steel piles, to convert the DC electricity generated by the solar panels into AC electricity. The portions of the Project Site outside the fenced-in areas would not be developed. The modules would be attached to single-axis trackers that allows all the panels to pivot along an axis to follow the path of the sun from east to west across the sky. The trackers would be attached to steel pile foundations.

Collections of strings or rows of panels would be connected by underground DC cabling to a central inverter that would convert the DC electricity into AC electricity so that it could be transmitted to the electrical grid. Each inverter would have a collocated mid-voltage transformer (MVT) which boosts the AC voltage to account for the standard electrical loss between the central inverters and the Artesia substation. From the MVTs, a network of underground AC power cables would connect to a single main power transformer (MPT) which would be located within the previously reviewed 161-kV Golden Triangle I Substation. Cables would be installed in trenches approximately 3- to 4-feet deep and 2- to 12-inches wide.

Earth-compacted roads would provide access to each inverter block for the purposes of operations, maintenance, and repairs.

## 2.2.2 Solar Facility Construction

Construction activities would take approximately 17 months to complete using a crew that ranges from 200 to 300 workers. Work would generally occur up to seven days a week during daylight hours. Additional hours after dark could be necessary to make up schedule deficiencies or to complete critical construction activities. Night-time construction, if determined necessary, would require lighting in some areas of the Project Site. Any additional night-time lighting would be downward-facing and timer- and/or motion-activated to minimize impacts to wildlife and any surrounding receptors, including nearby households.

Site preparation is generally required prior to construction of the Solar Facility and assembly of the solar arrays. Site preparation typically includes surveying and staking, removal of tall vegetation/small trees, light grading, clearing and grubbing, installation of security fencing around components near one another and not separated by public roads, erosion prevention and sediment control BMPs, and preparation of construction laydown areas. Solar array assembly and construction includes driving steel piles into the ground for the tracker support structures, installation of solar panels, and electrical connections and testing/verification.

Approximately 40 acres of the Project Site would be used as construction assembly areas (also called laydown areas) for worker assembly, safety briefings, vehicle parking, temporary offices, and material storage during construction. Some of these assembly areas, which would be spread out across the Project Site, would be staged within the locations proposed for the PV arrays. The laydown areas would be onsite for the duration of construction. Temporary construction trailers for material storage and office space would be parked onsite. Following completion of construction activities, most trailers, unused materials, and construction debris would be removed from the Project Site. If appropriate, a construction trailer may be utilized as an operations and maintenance building. Construction materials would be transported by truck and/or rail to the Project Site, where materials would be staged, assembled, and moved into place.

MS Solar 6 would use the existing landscape, such as slope, drainages, and roadways where feasible, minimizing grading work where practicable. Grading activities that could not be avoided would be performed using mobile earthmoving equipment, resulting in a moderately consistent slope on land. Native topsoil would be preserved to the greatest extent practicable during grading. Native topsoil would be stockpiled onsite and preserved for redistribution over the disturbed area after grading is complete. After construction, the disturbed areas would be seeded with a native seed mixture of certified weed-free, low-growing, noninvasive grasses, and herbaceous plants. Flowering vegetation also would be used, if available, to attract pollinator species such as honeybees and butterflies. Erosion control BMPs would be regularly inspected and maintained until vegetation in the disturbed areas has been established to the extent it meets permit restoration requirements. Water would be used for fugitive dust control and/or soil compaction

during construction on an as-needed basis. Water used during construction would either be trucked in from a municipal source or withdrawn from an onsite water well.

To manage stormwater during construction, onsite temporary sedimentation basins, sediment traps, or diversion berms would be constructed within the disturbed area of the Project Site. If needed, a diversion berm would be constructed along portions of the Project Site perimeter to contain stormwater onsite. Any necessary sedimentation basins and/or traps would be compliant with MDEQ requirements. If necessary, sedimentation basins and traps would be constructed either by impoundment of natural depressions or by excavating the existing soil. The floor and embankments of the basins would be allowed to naturally reestablish native vegetation after construction (or replanted as necessary) to provide natural stabilization and minimize subsequent erosion. Sediment traps would be placed in strategic drainage areas to prevent sediment from entering onsite jurisdictional streams and wetlands. Offsite sediment migration would be minimized by the placement of silt fencing around each area of ground disturbance within the Project Site. These stormwater BMPs would minimize the potential for sediment to enter onsite jurisdictional streams and wetlands and to minimize sediment migration offsite during construction. Once sufficient revegetation cover is achieved, the Project Site would be considered stabilized and temporary construction BMPs would be discontinued or removed.

Construction activities would be sequenced to minimize the time that bare soil in disturbed areas is exposed. In addition to the silt fencing described above, other appropriate controls, such as temporary cover, would be used as needed to minimize exposure of soil and eroded soil from leaving the work area. Disturbed areas, including road shoulders, construction office and laydown areas, ditches, and other Project-specific locations, would be seeded post-construction. If conditions require, soil may be further stabilized by mulch or sprayable fiber mat. Where required, hay mulch would be applied at three tons per acre and well distributed over the area. As part of NPDES permit authorization (see section 3.2), the site-specific SWPPP would be finalized with the final grading and civil design and would address all construction-related activities prior to construction commencement. Further guidance for clearing and construction activities can be found in Appendix A.

The design of the PV tracker support structures could vary depending on the final PV technology and vendor selected. Based on preliminary geotechnical survey results for the Project Site, the trackers would be attached to driven steel pile foundations. The steel pile foundations are typically galvanized and used where high load-bearing capacities are required. The pile is driven with a hydraulic ram. Soil disturbance is restricted to the pile insertion location to a depth typically less than 20 feet below grade; there is also potential for temporary soil disturbance from the hydraulic ram machinery, which is about the size of a small tractor. The tracker design and pile foundation design would be sealed by a registered Professional Engineer and Structural Engineer, respectively. Screw piles are another option for PV foundations which are drilled into the ground with a truck-mounted auger. Screw piles create a similar soil disturbance footprint as driven piles.

Solar panels would be manufactured offsite and shipped to the Project Site ready for installation. All final electrical collection cables would be underground, and electricians and assistants would

run the electrical cabling throughout the Solar Facility. The trenches to hold the cabling would be approximately 3- to 4-feet deep and 2- to 12-inches wide. The trenches would be backfilled with native soil and appropriately compacted.

The MPT would be supported on a concrete foundation within the Golden Triangle I Substation. An aboveground transmission cable would be constructed to connect the MPT through a circuit breaker.

Access roads would be needed to allow vehicular access within the solar facility. Typically, new permanent or temporary access roads are designed and located to avoid severe slope conditions and to minimize impacts to environmental resources. Information on access road siting is not available at this time; an additional analysis would be conducted if there are changes or upgrades to the scope of this Project. Access roads are typically about 12 to 16 feet wide and are surfaced with dirt, mulch, or gravel. Culverts and other drainage devices, fences, and gates would be installed, as necessary. Culverts installed for temporary access roads in any perennial streams would be removed following construction. However, in ephemeral streams (also known as wet weather conveyances - streams that only flow following a rainfall) the culverts would be left or removed, depending on the wishes of the landowner or any permit conditions that might apply. If desired by the property owner, TVA would restore new temporary access roads to previous conditions following construction. Additional applicable ROW clearing and environmental quality protection specifications are listed in TVA ROW Clearing Specifications, Environmental Quality Protection Specifications for Transmission Line Construction, and Transmission Construction Guidelines Near Streams.

## 2.2.3 Operations

Operation of the Golden Triangle Solar 2 Facility would require up to 1-2 full-time staff to manage the solar facility and conduct regular inspections. Inspections would include identifying any physical damage of panels, wiring, central inverters, transformers, and interconnection equipment, and drawing transformer oil samples. Vegetation on developed portions of the Project Site would be maintained to control growth and prevent overshadowing or shading of the PV panels. Trimming and mowing would likely be performed several times per year, depending on growth rate, to maintain an appropriate ground cover height of no more than approximately 12 to 18 inches. During operation of the Solar Facility, selective use of EPA-approved spot herbicides may also be employed around structures to control invasive weeds.

The proposed Solar Facility would be monitored remotely from the MS Solar 6 Headquarters Energy Control Center in Austin, Texas, 24 hours a day, seven days a week to identify security or operational issues. In the event a problem is discovered during nonworking hours, a repair crew or law enforcement personnel would be contacted if an immediate response were warranted.

Moving parts of the Solar Facility would be restricted to the east-to-west tracking motion of the single-axis solar modules, which amounts to a movement of less than a one degree angle every few minutes. This movement is barely perceptible. In the late afternoon, module rotation would

start to move from west-to-east in a similar slow motion to minimize row-to-row shading. At sunset, the modules would track to a flat or angled stow position. Otherwise, the PV modules would simply collect solar energy and transmit it to the TVA power grid. With the exception of fence repair, vegetation control, and periodic array inspection, repairs, and maintenance, the Solar Facility would have relatively little human activity during operation. No significant physical disturbances would occur during operation. Permanent lighting is anticipated as a potential onsite need during facility operations, independent of the potential operations and maintenance building. Permanent lighting would be downward-facing and timer- and/or motion-activated to minimize impacts to surrounding areas.

Rainfall in the region should be adequate to remove dust and other debris from the PV panels while maintaining acceptable energy production; therefore, manual panel washing is not anticipated unless a site-specific issue is identified. If later identified, module washing would occur no more than twice a year and would comply with appropriate BMPs.

## 2.2.4 Decommissioning and Reclamation

MS Solar 6 would operate the Project and sell power to TVA under the terms of the PPA for the first 20 years of its life. At the end of the term of the PPA, MS Solar 6 would assess whether to cease operations at the Solar Facility, replace equipment and attempt to enter into a new power purchase agreement, or make some other arrangement to sell the power. If operations were ceased, the facility would be decommissioned and dismantled, and the Project Site would be restored to its pre-construction conditions. In general, the majority of decommissioned equipment and materials would be recycled. Materials that could not be recycled would be disposed of at an approved facility in accordance with federal, state, and local laws and regulations. As the lease agreements with the landowners are for at least 35 years, site control would be maintained for longer than the 20-year PPA period, and MS Solar 6 may attempt to renegotiate further PPA terms with TVA. At the end of the 20-year contract period, TVA may also choose to purchase and operate the facility. If additional PPA terms are arranged or if TVA chooses to operate the facility, these activities would be evaluated through separate NEPA processes.

## 2.3 Alternatives Eliminated from Further Consideration

Numerous criteria were considered throughout the process of identifying a suitable site within TVA's service area that would meet the purpose and need of this Project as well as expanding TVA's renewable energy portfolio (TVA 2019a). The following is a list of key factors MS Solar 6 took under consideration during the site selection process.

- Site must be large-scale enough to accommodate enough PV panels to generate 150-MW AC output for transmission to the electrical network.
- Large contiguous parcels of land with at least 1,500 acres available for solar panel installation and additional acreage for related infrastructure.
- Availability of nearby electric infrastructure for interconnection to TVA's system with sufficient available transmission capacity.

- Generally flat landscape with minimal slope; preferably previously disturbed contiguous land with minimal existing infrastructure obstacles.
- Minimal presence of forested areas and wetlands.
- Parcels with appropriate local zoning regulations and located away from densely populated areas.
- Land that would allow developers to avoid and/or minimize impacts on known sensitive biological, visual, and cultural resources.

The process of screening potential locations and ultimately eliminating sites that did not possess the necessary attributes led to the selection of the Project Site.

## 2.4 Comparison of Alternatives

This EA evaluates the potential environmental effects that could result from implementation of the No Action Alternative or the Proposed Action Alternative at the proposed Project Site in Lowndes County, Mississippi. The analysis of impacts described in this EA is based on current conditions as well as potential future conditions on the parcels associated with the Project and the surrounding area. A comparison of potential impacts from each alternative is summarized in Table 2-1.

Table 2-1: Comparison of Impacts by Alternative

Resource Area	Impacts from the No Action Alternative	Potential Impacts from the Proposed Action Alternative	
Land Use	No direct or indirect impacts anticipated.	7	
	No direct or indirect changes anticipated.	Geology: Minor direct impacts on potential shallow subsurface geological resources.	
Geology, Soils, and Prime Farmlands		Soils: Minor direct impacts on soils from potential minimal increases in erosion and sedimentation during construction. Once stabilized and facility is operational, impacts on soils would be offset by the beneficial effects to soil health with the use of native and noninvasive vegetation.	
		Farmlands: Direct impacts on farmland from the conversion of agricultural land to solar for the duration of the Project.	
Water Resources	No direct or indirect changes to current conditions anticipated.	Groundwater: Negligible direct impacts on the supply from use of a new water well during operation of Solar Facility. Minor beneficial effects are anticipated from the reduction in fertilizer and pesticide application.	

Resource Area   Impacts from the No Action Alternative		Potential Impacts from the Proposed Action Alternative	
		Surface water: Minor beneficial impacts on surface water due to the reduction in fertilizer and pesticide application once agricultural operations are not occurring. Minor short-term impacts from road crossings, erosion, and sedimentation during construction (until site is stabilized).	
		Wetlands: No permanent impacts on wetlands. No proposed placement of permanent structures within wetlands.	
		Floodplains: There are approximately 135 acres of FEMA designated floodplain within the Project Site; however, MS Solar 6 would not install arrays within the entire floodplain. Required permits and approvals would be obtained for placement of PV panels within floodplains prior to initiating construction. Access roads would be constructed such that upstream flood elevations would not be increased by more than 1.0 foot.	
	No direct or indirect impacts anticipated.	Vegetation: Three main vegetation communities would be affected: actively cultivated bean and corn fields, hay pastures, and forested uplands.	
Biological Resources		Wildlife: Moderate adverse impacts on wildlife during construction. Some forested areas would be converted to open land for solar array placement. Wildlife that can use early successional habitat is expected to return to the area once operational. The Project is not anticipated to have long-term significant impacts on migratory bird species of concern.	
		Protected Species: No known occurrences of protected species were found within the Survey Area. The Project is not anticipated to significantly affect federal or statelisted species.	
	No direct or indirect impacts anticipated.	During construction, minor temporary impacts on visual resources would occur due to the alteration of the existing agricultural viewshed and increased construction activity.	
Visual Resources		During operation, moderate direct impacts in the immediate Project vicinity due to the presence and quantity of PV panels. Impacts on residents and visitors to the town of Artesia would be minimized through the presence of existing natural screening buffers including forest areas and topography. If existing buffers are not sufficient in shielding residents in Artesia from the Solar Facility, MS Solar 6 would install a privacy fence or shrubbery along the perimeter of the Project Site on a case-by-case basis.	
Noise	No direct or indirect impacts anticipated.	Minor temporary noise impacts would be experienced during construction. Negligible adverse impacts from noise associated with operation.	

Resource Area	Impacts from the No Action Alternative	Potential Impacts from the Proposed Action Alternative
Air Quality and	No direct or indirect impacts anticipated. Continue using fossil fuels.	Air Quality: Minor direct impacts on air quality could occur during site preparation involving heavy, earth moving construction equipment (temporary emissions). No adverse impacts on air quality from operations.
Greenhouse Gas Emissions		GHG: Temporary and minor increases in GHG emissions would be expected during construction from operation of equipment. Offsetting beneficial effects would also occur, due to the nearly emissions-free power generated by the solar facility, offsetting power that would otherwise be generated by the combustion of fossil fuels.
Cultural Resources	No direct or indirect impacts anticipated.	Archaeological Resources: Due to TVA's Avoidance Agreements for known NRHP-eligible and NRHP-listed sites, no impacts on archaeological resources would occur.
		Architectural Resources: Recommendation of no adverse effect on architectural resources.
Utilities	No direct or indirect impacts anticipated.	No direct or indirect adverse impacts are anticipated to utilities. The region would experience long-term benefits to electrical services.
Waste Management	No direct or indirect impacts anticipated.	With the implementation of appropriate BMPs, no impacts on waste management would be anticipated.
Public and Occupational Health & Safety	No direct or indirect impacts anticipated.	Minor, temporary impacts during construction. No public health or safety hazards would be anticipated during operation.
Transportation impacts anticipated. and from impact of construction construction		Due to moderate increases from workers commuting to and from the Project Site during construction, a moderate impact on transportation would be anticipated during construction. Negligible direct impacts and no indirect impacts on transportation would occur during operation.
	No direct or indirect impacts anticipated.	Short-term beneficial economic impacts would result from construction, including the purchase of materials, equipment, and services and a temporary increase in employment, income, and population.
Socioeconomics		Positive, long-term, direct impacts on economics and population from Project operation. The local tax base would experience an increase from construction of the Solar Facility which would benefit Lowndes County and the Golden Triangle region of eastern Mississippi.
		There would not be disproportionately high or adverse direct or indirect impacts to minority or low-income populations.

## 2.5 Best Management Practices and Mitigation Measures

MS Solar 6 would implement minimization and mitigation measures for resources potentially affected by the Project. These measures would be developed in conjunction with industry supported BMPs, requirements of regulatory permits, and adherence to the following plans:

- Stormwater Pollution Prevention Plan (SPPP)
- Spill Prevention, Containment, and Countermeasures (SPCC) Plan, and
- Cultural and Architectural Resources Unanticipated Discovery Plan.

Additional details are provided in sub-sections 2.5.1 and 2.5.2.

## 2.5.1 Golden Triangle II Solar Facility

MS Solar 6 would implement the following BMPs and mitigation measures associated with potentially affected resources, as follows:

BMPs include actions to:

- o Install anti-reflective, PV panel surfaces to minimize glare and reflection.
- Install silt fencing along the perimeter of areas that would be cleared, consistent with local and state stormwater regulations.
- Maintain stormwater BMPs in each area until stabilization (adequate vegetation regrowth) has been achieved.
- Avoid direct impacts on perennial and intermittent streams by maintaining a riparian buffer at most perennial and intermittent streams and jurisdictional wetlands.
- Plant or seed with noninvasive vegetation and include native and naturalized plant species to encourage beneficial habitat, reduce erosion, and limit the spread of invasive species.
- Utilize vegetation that benefits pollinator species to the extent practicable.
- Utilize timer- and/or motion-activated downward facing security lighting to limit attracting wildlife, such as migratory birds and bats.
- Use dust mitigation activities such as watering dry exposed soils, covering openbody trucks, and establishing a speed limit to minimize fugitive dust.
- Install temporary construction fencing around natural resources that should be avoided.
- Cover or stabilize exposed soil stockpiles.

## Mitigation Measures include:

- Should traffic flow become a problem, consider implementation of staggered worker shifts during construction and a flag person along the roadside during deliveries that may coincide with heavy commute times to manage the flow of traffic near the Project Site.
- There are three residences within visual proximity to GT2. Where existing natural buffers are not sufficient in shielding residents from the Solar Facility, MS Solar 6 would install vegetative screening along the perimeter of the Project Site (additional information in section 3.5.2).
- Temporary laydown areas, construction trailers, and parking areas would be placed outside the floodplain during construction.
- Access roads within 100-year floodplains would be constructed such that upstream flood elevations would not be increased by more than 1.0 foot.
- The portions of the fencing within the floodplain should be designed to withstand flooding with minimum damage.
- Any excavated material or debris generated when GT2 is decommissioned and dismantled would be disposed of in an area outside 100-year floodplains.
- The lowest section of each solar module, at full tilt, is at least one foot above the 100-year base flood elevation.
- At the end of the Project's useful life, non-recyclable and/or non-reusable debris would be disposed of at an appropriate location outside 100-year floodplains.
- Any permanent operations and maintenance building(s) will be located outside 100year floodplains and elevated to at least one foot above the 100-year floodplain at that location.
- For residences that are within 500 feet of an inverter, a pre-construction sound study including an ambient survey would be conducted to quantify the existing ambient environment. After the project reaches commercial operation, MS Solar 6 would measure the sound levels at residential property lines and identify any equipment that generates a Ldn sound level that exceeds 55 dBA at the property line. If there are locations where noise levels exceed that threshold, MS Solar 6 would install sound buffers (walls, fences with screening, or vegetation) to minimize the noise levels from operating equipment.
- The Project exclusion areas would be avoided and not allow any development, disturbance, or other construction activities associated with the development of the project or future activities associated with the operation and maintenance of the solar facility. Exclusion areas include those identified plant species, jurisdictional wetlands, 100-year floodplains, and cultural resources. Exclusion area cultural resources would

- be avoided with a 100-foot (ft) buffer if they were identified as eligible or unassessed sites.
- Interpretive signs would be installed. One interpretive sign would mark the location of the Thomas Wilburn Harness Horse Racing Track (22Lo1017) discussing the significance of the track to the community. The second interpretive sign will discuss the Oakland Plantation Historic District, the significance of the history of plantations and tenant farms in the region.

#### 2.6 The Preferred Alternative

TVA's preferred alternative for fulfilling its purpose and need is the Proposed Action Alternative. This alternative would generate renewable energy for TVA and its customers with only minor direct and indirect environmental impacts due to the implementation of BMPs and minimization and mitigation efforts, as described in Section 2.5.1 and Section 2.5.2. Implementation of the Project would help meet TVA's renewable energy goals and would help TVA meet customer-driven energy demands on the TVA system.

#### 3.0 AFFECTED ENVIRONMENT AND ENVIRONMENTAL CONSEQUENCES

This chapter describes the importance, nature, and extent of environmental resources in their current setting at the Project Site. This chapter also provides a measure for the assessment of potential effects of the alternatives described in Section 2.0. The scope of environmental consequences assessed in this EA for the Proposed Action focuses on impacts related to the construction and operation of the proposed Solar Facility.

#### 3.1 Land Use

This section provides an overview of the existing and surrounding land use at the Project Site. Potential impacts on land use associated with the alternatives are described below.

#### 3.1.1 Affected Environment

The EPA defines "land use" as the human use of land for activities such as agricultural, residential, and recreational uses (USEPA, 2020a). The GT2 Project Site is east of U.S. Hwy Alt 45 and south of Gilmer Wilburn Road in Artesia, Mississippi. Imagery data collected from the National NRCS Land Use Land Cover Database identify the Project Site as primarily cultivated crops and pastures.

The Project Site is generally flat with minor changes in elevation and ranges from approximately 225 to 250 feet above mean sea level. Elevation is generally higher in the northern portion of the Project Site south of Gilmer Wilburn Road and gets lower toward the southern portion of the Project Site near North Branch Magowah Creek. MS Solar 6 intends to maximize the use of agricultural and pasturelands to the greatest extent practicable, thus minimizing tree clearing and impacts on wetlands. Table 3-1 provides a detailed breakdown of land use and land cover within the Project Site.

Table 3-1: Land Use and Percent Cover within the Project Site

Land Use	Approximate Area (acres)	% Cover
Open Water	1.5	0.1
Developed, Open Space	5.7	0.4
Deciduous Forest	262.4	17.2
Evergreen Forest	49.2	3.2
Mixed Forest	16.9	1.1
Shrub/Scrub	15.5	1.0
Hay/Pasture	593.5	39.0
Cultivated Crops, Agriculture	494.8	27.9
Emergent Herbaceous Wetlands	1.9	0.1
Woody Wetlands	152.1	10.0
Total	1523.6	100.0

The closest town to the Project Site is Artesia, approximately one mile to the northwest of the Project boundary. Artesia has a population of approximately 425 people (City-Data, 2021). The Project Site is located about halfway between the municipalities of Columbus and Starkville, Mississippi. Columbus and Starkville have populations of 23,573 and 25,653 people, respectively (USCB, 2021c). There is one residence within the Project Site, and both are south of Gilmer Wilburn Road. The Golden Triangle Regional Airport and a variety of other industrial facilities are located northeast of the Project vicinity. The areas immediately surrounding the Project Site are similar in land use and are primarily agriculture/pasture or undeveloped.

According to historical aerial imagery and topographic quadrangle maps, the current land use of the Project Site has remained primarily undeveloped or used for agriculture/pasture with no significant land use changes in recent history. Figure 3-1 provides the land use classifications provided in the NRCS Land Use Land Cover Dataset.




Figure 3-1: Land Cover within the Project Site

# 3.1.2 Environmental Consequences

#### 3.1.2.1 No Action Alternative

Under the No Action Alternative, no Project-related impacts to land use would occur. Existing land use would be expected to remain a mix of agricultural and undeveloped land for the foreseeable future.

## 3.1.2.2 Proposed Action

Implementation of the Proposed Action would result in construction and operation of the Solar Facility; therefore, the land use within the Project Site would change from primarily agriculture and pastureland to industrial renewable energy production. The undeveloped forested portions of the Project Site would remain undeveloped. The undeveloped areas that are currently agricultural would remain undeveloped with no farming or other activities occurring. The Project Site is in a rural area with limited zoning restrictions and would be compatible to land uses in the surrounding areas. Golden Triangle Regional Airport is approximately three miles to the northeast of the Project Site and an approximately 1,200-acre industrial megasite facility is three miles northeast of the Project Site along with a variety of other small industrial facilities northeast of the Project Site. Installation of the Solar Facility would increase industrial development southward where agriculture is currently the dominant land use. If the Solar Facility were to be decommissioned, the land could be returned to agriculture or used for a variety of other development strategies as allowed by local zoning legislation. Minor direct impacts are anticipated from the conversion of actively cultivated agricultural land and pasture/livestock grazing land, and forested areas to solar generation.

Construction and operation of the Solar Facility is proposed in an area primarily used for agriculture and pasture. There are no outdoor recreation areas in the vicinity of the Project Site and development of the Project would not impact public recreational activities or facilities associated with recreational activities. Therefore, adverse direct or indirect impacts on land use are not anticipated.

# 3.2 Geology, Soils, and Prime Farmland

#### 3.2.1 Affected Environment

This section describes the existing geological resources at the Project Site and the potential impacts on these geological resources that would be associated with the No Action and Proposed Action Alternatives. Components of geological resources that are analyzed include geology, geological hazards, paleontology, soils, and prime farmland.

## 3.2.1.1 Geology

The Project Site is in the Alabama and Mississippi Blackland Prairie Major Land Resource Area (MLRA) in Northeast Mississippi. This Project Site is in the East Gulf Coastal Plain section of the

Coastal Plain Physiographic Province (NPS 2018). The Coastal Plain Province is generally underlain by poorly consolidated clastic rocks from the Mesozoic and Cenozoic (Jurassic to Quaternary) age.

# 3.2.1.2 Geological Hazards

Geological hazards can include landslides, volcanoes, earthquakes/seismic activity, and subsidence/sinkholes. The Project Site and surrounding areas are relatively stable without significant slopes within several miles, mitigating potential risk for landslides. There are no volcanoes within several hundred miles of the Project Site (U.S. Geological Service [USGS] 2019).

Sinkholes can be common when subsurface rock composition is evaporite rock (salt, gypsum, and anhydrite) and carbonates (limestone and dolomite) which can naturally be dissolved by groundwater circulating through them. When rock dissolves, spaces and caverns develop underground. These types of formations are referred to as karst topography. Land over sinkholes may stay intact until there is not enough support for the land above the spaces. Then, a collapse of the land surface can occur. Land collapses can vary greatly in size and shape (USGS, 2016). Human activities can also expedite cavity formation in more susceptible materials and trigger a collapse or collapse an existing sub-surface cavity site. A geotechnical survey will be performed within the Project Site to determine that no carbonate bedrock units underlie the Project Site.

Surface faulting, ground motion and deformation, liquification, and subsidence as a result of seismic activity were assessed at the Project Site. Susceptibility of structures or humans to experience seismic activity are often shown via the Modified Mercalli Scale. Values on the Mercalli scale are translated into a peak ground acceleration (PGA) value to measure the maximum force experienced. The PGA is the maximum acceleration experienced by a building or object at ground level during an earthquake on uniform, firm-rock site conditions. The PGA is measured in terms as a percentage "g," the acceleration due to gravity. The USGS Earthquake Hazards Program publishes seismic hazard map data layers that display the PGA with ten percent (one in 500-year event) probability of exceedance in 50 years. The potential ground motion for the Project Site is 0.1 g, for a PGA with a two percent probability of exceedance within 50 years (USGS 2014). A 0.1 g earthquake will have a weak to moderate perceived shaking with a low moderate potential for structural damage. The Project Site has a low to moderate risk for earthquakes that will cause structural damage (USGS 2019a).

## 3.2.1.3 Paleontology

There are no Precambrian rocks in the Project Site because the land that is now the State of Mississippi did not exist during this period. During the Paleozoic Era, Mississippi was primarily under the sea and provided habitat to organisms such as mollusks, crinoids, brachiopods, and trilobites. As water levels receded, broad coastal plains remained, and by the end of the Paleozoic era the entire state of Mississippi would have been above sea level and exposed to erosional pressures. Rocks deposited during flooding of the late Mesozoic Era can potentially contain fossils of invertebrates, vertebrates, and driftwood. Fluctuating sea levels from glacial influence on

climate during the Cenozoic Era provided layers of windblown loess eroded from the Mississippi River to cover a large portion of the state to the northwest. Fossils of mollusks, other invertebrates, and large terrestrial mammals have been recovered from the loess deposits (Paleontology Portal, 2020).

#### 3.2.1.4 Soils

There are 12 soil map units within the Project Site, as identified in the USDA NRCS geospatial data for Lowndes County. Each type is listed along with its percent coverage in the table below. The soil map units are illustrated in Figure 3-2. Brooksville silty-clay, Okolona silty-clay, and Vaiden silty-clay soils are considered prime farmland and Catalpa silty-clay and Leeper silty clay soils are considered prime farmland if protected from flooding or not frequently flooded during the growing season. Sumter-silty-clay-loam soils are considered farmland of statewide importance. Prime farmland and farmland of statewide importance are discussed further below.

Table 3-2: Soils within the Project Site

Soil Map Unit	Farmland	Hydric	Area (acres)	Percent
Brooksville silty clay, 0 to 1 percent slopes	All areas are prime farmland	Not Hydric	158.5	10.4%
Brooksville silty clay, 1 to 3 percent slopes	All areas are prime farmland	Not Hydric	299.4	19.6%
Okolona silty clay, 0 to 1 percent slopes	All areas are prime farmland	Not Hydric	28.4	1.9%
Okolona silty clay, 1 to 3 percent slopes	All areas are prime farmland	Not Hydric	11.2	0.7%
Vaiden silty clay, 0 to 2 percent slopes	All areas are prime farmland	Hydric	214.0	14.0%
Vaiden silty clay, 2 to 5 percent slopes, eroded	All areas are prime farmland	Not Hydric	399.9	26.2%
Sumter silty clay loam, 2 to 5 percent slopes, eroded	Farmland of statewide importance	Not Hydric	15.9	1.0%
Sumter silty clay loam, 5 to 12 percent slopes, eroded	Farmland of statewide importance	Hydric	62.6	4.1%
Demopolis-Binnsville complex, 2 to 8 percent slopes, eroded	Not prime farmland	Hydric	52.8	3.5%
Sumter-Demopolis-Chalk outcrop complex, 5 to 20 percent slopes, severely eroded	Not prime farmland	Hydric	6.0	0.4%
Water	Not prime farmland	NA	2.8	0.2%
Leeper silty clay, 0 to 2 percent slopes, occasionally flooded	Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season	Hydric	33.3	2.2%

Soil Map Unit	Farmland	Hydric	Area (acres)	Percent
Catalpa silty clay	Prime farmland if protected from flooding or not frequently flooded during the growing season	Hydric	238.9	15.7%
		Total	1,523.6	100%

Source: NRCS 2021

The Project Site is located within MLRA-135A (Alabama and Mississippi Blackland Prairie). Therefore, upland soils exhibit a low chroma matrix, which is characteristic of the native parent material and is not necessarily caused by extensive soil saturation. The Leeper soil series is typically found in floodplains of MLRA-135A and consists of very deep, somewhat poorly drained soils. These soils have dark grayish brown "A" and "B" horizons and are derived from clayey alluvium parent material. The Vaiden soil series is typically found in uplands and old stream terraces of MLRA-135A and consist of very deep somewhat poorly drained, very slowly permeable soils that formed in clayey sediments over chalk or calcareous clays. These soils have yellowish brown "A" and "B" horizons and are principally used for cropland, pasture, hay production and woodland. The Catalpa soil series is also typically found in floodplains or low terraces of streams that drain areas of MLRA-135A. The Catalpa series soils are somewhat poorly to moderately well drained and derived from clayey alluvial sediments. These soils have color ranging from a very dark grayish brown "A" horizon to a dark grayish brown and olive brown "B" horizon. The Okolona series soils are deep, well drained very slowly permeable soils found in upland areas of MLRA-135A. These are generally level to gently sloping soils derived from calcareous clayey parent material underlain by marly clay and chalk. These soils have color ranging from a very dark grayish brown "A" horizon to olive "B" horizon. The Brooksville soil series are deep, somewhat poorly drained soils found in uplands of MLRA-135. Brooksville soils have an "A" horizon that is very dark grayish brown to dark olive gray and a dark grayish "B" horizon (USDA, 2018).



Figure 3-2: Soils within the Project Site

#### 3.2.1.5 Prime Farmland

Prime Farmland is a designation assigned by the USDA NRCS that identifies soils and land that has the best combination of physical and chemical properties for producing food, feed, forage, fiber, and oilseed crops. The land could be cropland, pastureland, rangeland, forest land, or other land, but not urban built-up land or water. The Farmland Protection Policy Act ([FPPA]; 7 United States Code [U.S.C.] 4201 et seq.) requires federal agencies to consider the effects of their actions on prime or unique farmlands. The purpose of the FPPA is "to minimize the extent to which federal programs contribute to the unnecessary and irreversible conversion of farmland to nonagricultural uses." (USDA, 2021). Table 3-2 describes the soil types and farmland classification within the Project Site. The table is representative of soils that would be affected by the preliminary array layout (shown in Figure 2-2). Locations of prime farmland soils on the Project Site are shown in Figure 3-3. Data analysis from the NRCS indicates prime farmland soils and farmland of statewide importance soils make up approximately 95% of the Project Site.

Hydric rating is an indicator of the percentage of a map unit that meets the criteria for hydric soils (USDA ND). Hydric soils are formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part. Each map unit is designated as follows:

- Hydric: All of the components that make up the map unit are rated as being hydric.
- Predominantly Hydric: 66 99 percent of the components that comprise the map unit are rated as being hydric.
- Partially Hydric: 33 66 percent of the components that comprise the map unit are rated as being hydric.
- Predominantly Nonhydric: Up to 33 percent of the components that comprise the map unit are rated as being hydric.
- Nonhydric: None of the components that comprise the map unit are rated as being hydric.




Figure 3-3: Prime Farmland Soils within the Project Site

## 3.2.2 Environmental Consequences

This section describes the potential impacts to Geology, Soils, and Prime Farmland from implementation of the Proposed Action or the No Action Alternative.

## 3.2.2.1 No Action Alternative

Under the No Action Alternative, no Project-related impacts on geology, geological hazards, paleontological, soil resources, or prime farmlands would result. Existing land use would be expected to remain as a mix of agriculture and undeveloped land. If current land use remains unchanged, impacts to soils from continued agricultural use could cause a depletion of nutrients, causing a minor change to the Project Site.

## 3.2.2.2 Proposed Action

Under the Proposed Action, temporary, minor direct impacts to geology and soil resources would occur from construction and operation of the Project. Minor grading and clearing for the Solar Facility would cause temporary, minor, localized increases in erosion and sedimentation, resulting in minor impacts to geology and soils.

# Geology

Implementation of the Proposed Action could result in minor impacts to geology. In addition to minor grading, installation of pilings to support solar arrays would be mechanically driven to depths of up to six feet. Trenching up to approximately three feet for underground wiring connections between solar panels would also be required. Due to limited areas and potential shallow subsurface disturbances, minor direct impacts to subsurface geological resources are anticipated.

## Geological Hazards

The Project Site is in an area with carbonate bedrock geology and karst landforms associated with a risk for sinkholes. Geological hazards associated with potential sinkholes will be investigated within the Project Site to ensure there are no known sinkholes anywhere within the Project Site, and the future formation of sink holes are unlikely. There is also minor to moderate potential for small to moderate intensity seismic activity. The Solar Facility would be designed to comply with applicable standards. In the unlikely event that seismic activity and/or sinkholes would occur at the Project Site, only minor impacts to the Solar Facility and associated infrastructure would occur. Impacts to resources outside of the Project Site from geological hazards associated with construction of the proposed Solar Facility are unlikely.

## <u>Paleontology</u>

If paleontological resources are identified during initial construction or operational activities, a qualified paleontologist will be consulted to recover and analyze the resources for potential impacts. MS Solar 6 would develop and implement a recovery plan and mitigation strategy.

# Soils

During construction, all soils within the Project Site could potentially be disturbed from site preparation and construction activities. In areas where vegetation or tree removal is proposed, soil would be stockpiled and replaced to the greatest extent practicable. Due to the limited vegetation and tree clearing activities likely to occur, only a nominal amount of off-site soil may be required or hauling away of on-site soil may be necessary. If other borrow material such as sand, gravel, rip rap, or other aggregate is necessary during site preparation, these resources may be used from on-site sources within the Project Site or nearby previously permitted off-site sources.

Implementation of BMPs outlined in the Erosion and Sedimentation Control Plan (ESCP) during construction, in addition to a re-vegetation strategy of planting or seeding non-invasive vegetation, including native and naturalized plant species post-construction would minimize the potential for increased soil erosion. Activities associated with construction of the Proposed Action would not result in adverse impacts to soils on the Project Site.

Minor disturbance to soils would occur during operation of the Proposed Action. Activities ranging from routine and non-routine maintenance of the arrays, array inspections, facility maintenance, fence repairs, and vegetation control would be an on-going potential disturbance to soils within the Project Site. Vegetation control would be conducted using mechanized equipment such as tractors, mowers, and trimmers. MS Solar 6 proposes to re-vegetate the Project Site with low-growing native vegetation that would reduce the routine vegetation maintenance while also limiting interference with the solar arrays. Broad application of herbicides is not anticipated. However, if selective herbicides are necessary for small applications around problematic areas, they would be applied by a licensed contractor or qualified staff. Maintenance activities discussed in this section would not result in adverse impacts to soils on the Project Site during operation.

## Prime Farmland

Implementation of the Proposed Action would result in a large portion of the Project Site being developed into GT2, changing the land use to renewable energy from the existing agriculture and pastureland. Since the entire Project Site is essentially prime farmland or farmland of statewide importance, implementation of the Proposed Action would result in direct impacts to prime farmland from installation of the solar arrays and construction of permanent structures necessary for operation of the Solar Facility. Based on information provided by the local NRCS office, there is approximately 330,000 acres of farmable land in Lowndes County, of which 250,000 acres is FPPA-defined farmland. Any area within the Project Site not developed for the Solar Facility would likely remain undeveloped or continue to remain agricultural land. Depending on the size of undeveloped agricultural areas within the Project Site, MS Solar 6 may allow certain landowners to resume agricultural activities if they are interested. Based on the availability of farmable land in Lowndes County, the Project would not result in long-term adverse impacts on agricultural operations within Lowndes County or the town of Artesia.

During site preparation and grading activities, topsoil would be stockpiled and re-applied to the respective surface areas once grading is complete. Soils within the Project Site do not have

characteristics that would require specific construction requirements or techniques. If the Solar Facility is decommissioned and closure occurs, facility components would be removed, and farming could subsequently be resumed with limited long-term loss of agriculture production.

Implementation of the Proposed Action would result in temporary adverse effects to prime farmland during operation of the Solar Facility. Stockpiling topsoil for reuse and installing appropriate erosion control devices would preserve topsoil at the Project Site. Adhering to BMPs during construction and operation of the Solar Facility and revegetating the site with native plant cover would limit erosion. Implementation of these BMPs would result in minor impacts to prime farmland. If the Solar Facility is decommissioned and Project equipment is removed, the Project Site could be returned to agricultural, and pastureland uses with a negligible loss to soil productivity. Beneficial impacts to soil health could result from a re-vegetation strategy using native and non-invasive species while terminating the need for broad application of herbicides, pesticides, and fertilizers. Based on the above mitigation measures as well as the availability of farmable land in Lowndes County, impacts on prime farmland soils would be minor and mostly reversible.

## 3.3 Water Resources

This section provides an overview of existing water resources in the Project Site, and the potential impacts on these water resources that would be associated with the No Action and Proposed Action Alternatives. Water resources discussed in this section include groundwater and surface water, including wetlands and floodplains.

#### 3.3.1 Affected Environment

#### 3.3.1.1 Groundwater

Groundwater supplies more than 38 percent of water needs for public water supply, agriculture, industry, mining, thermoelectric power, and domestic and commercial use in the Arkansas, Louisiana, and Mississippi area, which encompasses Lowndes County (Segment 5 of the Ground Water Atlas of the United States). In Mississippi, groundwater provides approximately 68 percent of the freshwater used in the state, most of which is used for agriculture (USGS, 1998). In Lowndes County, approximately 8.37 million gallons per day of groundwater is withdrawn from public supply self-supplied groundwater, and an estimated 192 gallons per person per day are used (USGS 2018).

Major aquifers near the Project Site are the Southeastern Coastal Plains (Black Warrior River) aquifer system and the Mississippi embayment aquifer system. These aquifers consist primarily of an unconsolidated to poorly consolidated Coastal Plain strata of gravel, sand, clay, and some limestone of the Cretaceous and Holocene age. Specifically, the Mississippi embayment aquifer system consists of permeable sedimentary rock from the late Cretaceous to middle Eocene period and is the largest aquifer system in the Coastal Plain. In the mid-1980's, the Mississippi embayment aquifer system provided nearly six percent of the total groundwater withdrawn in Mississippi, Louisiana, and Arkansas. Large quantities of groundwater withdrawn via wells from

the Coastal Plain aquifer systems during the last 100 years have not only lowered water levels, but they have also caused encroachment of salt water, decreased thickness of several aquifers, and even altered regional groundwater flow (USGS 1998).

Prior to the development of the Coastal Plain, regional groundwater flow was primarily driven topographically from elevated interstream recharge areas on the east and west sides of the Mississippi River to discharge areas in the valleys. Groundwater in the region is generally safe for most uses and quality in the aquifers correlate with groundwater flow, depth, and principal chemical constituents. Chemical constituents in groundwater below the Project Site are calcium bicarbonate, sodium bicarbonate, and sodium chloride where areas of the aquifer are deeply buried (USGS, 1998).

## 3.3.1.2 Surface Water

Surface waters are defined as water features that are on the Earth's surface, typically consisting of streams, lakes, ponds, and wetlands. Surface water features are further segregated into perennial, intermittent, and ephemeral. Perennial waters are permanent surface water features that have water present throughout the year. They typically exist as streams, rivers, lakes, springs, and swamps. During periods of little or no rain, the water level is maintained by groundwater contributions. Intermittent classification is generally restricted to streams that have a well-defined channel but only contain water part of the year, typically during winter and spring seasons when the stream bed is below the water table. Intermittent streams often do not support the diversity of biological and hydrological characteristics that perennial streams do. Ephemeral waters are features that only hold water for part of the year or flow as a result of stormwater events. Ephemeral streams (also known as wet-weather conveyances) are features that only flow in direct response to precipitation events. Flow would only occur during and shortly after large precipitation events. These features typically lack the biological, hydrological, and physical characteristics of intermittent and prennial streams. Examples of ephemeral streams/drainages include topographic swales and/ or dry drainages with poor bed and bank development.

Wetlands are those areas inundated by surface water or groundwater such that vegetation adapted to saturated soil conditions is prevalent. Examples include swamps, marshes, bogs, and wet meadows. Wetland habitat provides valuable public benefits including flood/erosion control, water quality improvement, wildlife habitat, and recreation opportunities.

Surface waters that meet certain physical and hydrologic criteria (defined bed and bank, ordinary high water mark, or specific hydrologic, soil, and vegetation composition) as defined in the Clean Water Act (CWA) are considered Waters of the U.S. (WOTUS) (or jurisdictional waters) and are under the regulatory jurisdiction of USACE. The CWA is the primary federal law that regulates discharges of pollutants and/or fill materials into WOTUS as outlined in Sections 402, 404 and 401. A jurisdictional determination by the USACE typically governs the activities affecting WOTUS.

The Project is within the EPA Blackland Prairie Ecoregion (Level 4) and is within the Middle Tombigbee River [Hydrologic Unit Code (HUC) 03160106] and Tibbee Creek (8-digit HUC)

03160104) sub-basins (USGS 2021b). The northern half of the Project Site generally drains south and west into North Branch Magowah Creek which flows in a westerly direction eventually into Magowah Creek which drains to the Tombigbee River. North Branch Magowah Creek, (designated use is "Aquatic Life Support") is not listed on Mississippi's 2020 303(d) List (MDEQ 2020).

#### **3.3.1.3 Wetlands**

Field surveys were conducted on March 7 through March 13, 2021, to determine the presence of potentially jurisdictional wetlands and waterbodies within a prescribed Survey Area which encompasses the more refined Project Site (Appendix B). The delineation was conducted in accordance with the 1987 Corps of Engineers Wetlands Delineation Manual (1987 Manual) and the 2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Atlantic and Gulf Coast Plain Region – Version 2.0 (Regional Supplement).

Wetlands are those areas inundated by surface water or groundwater such that vegetation adapted to saturated soil conditions is prevalent. Examples include swamps, marshes, bogs, and wet meadows. Wetland habitat provides valuable public benefits including flood/erosion control, water quality improvement, wildlife habitat, and recreation opportunities. Appendix B contains the *Golden Triangle II Solar Project Wetland and Waterbody Delineation Report*. Environmental Field Surveys were performed over a large survey area that included almost 1,525 acres. Based on the results of the surveys, MS Solar 6 refined the site design to minimize or avoid impacts on environmental resources, most specifically wetlands, waterbodies, and habitat that could be suitable for rare, threatened, or endangered plants and animals.

Under Wetlands Protection Executive Order (EO) 11990, federal agencies shall avoid, to the extent possible, the long- and short-term adverse impacts associated with the destruction or modification of wetlands and avoid direct or indirect support of new construction in wetlands wherever there is a practicable alternative. Under Section 404 of the CWA, unavoidable impacts to wetlands should be compensated through a process known as compensatory mitigation. Wetlands on the Project Site were classified by hydrologic regime and vegetation cover type in accordance with the Cowardin Classification System (Cowardin et. al. 1979). Three wetland types were identified within the Project Area: palustrine emergent (PEM; 11.89 acres), palustrine forested (PFO; 2.8 acres), and palustrine scrub-shrub (PSS; 2.49 acres) for a total of 17.18 acres of potentially jurisdictional wetlands. The common overstory and understory vegetation of PFO wetlands consists of water oak (Quercus nigra) sugarberry (Celtis laevigata), Osage orange (Maclura pomifera), green ash (Fraxinus pennsylvanica), black willow (Salix nigra), American sycamore (Platanus occidentalis), red maple (Acer rubrum), Chinese privet (Ligustrum sinense), American elm (Ulmus americana) and willow oak (Quercus phellos). Common vegetation comprising the understory includes poison ivy (Toxicodendron radicans), butterweed, roundleaf greenbrier (Smilax rotundifolia), Virginia creeper (Parthenocissus quinquefolia), yellow nutsedge (Cyperus esculentus), and soft rush (Juncus effusus). Typical, dominant scrub shrub species encountered during surveys included green ash, black willow (Salix nigra), butterweed, bulbous bittercress (Cardamine bulbosa), and soft rush. Species commonly found in the PEM within the

Project Area included shallow sedge (*Carex lurida*), black willow, alligator weed (*Alternanthera philoxeroides*), rough 3-6ocklebur (*Xanthium strumarium*), butterweed, annual ragweed (*Ambrosia artemisiifolia*), great ragweed (*Ambrosia trifida*), woolgrass (*Scirpus cyperinus*), curly dock (*Rumex crispus*), blackberries (*Rubus* spp.), spike rush (*Eleocharis palustris*), yellow nutsedge, and soft rush.

## 3.3.1.4 Floodplains

A floodplain is the relatively level land area along a stream or river that is subject to periodic flooding. The area subject to a one-percent chance of flooding in any given year is normally called the 100-year floodplain. The area subject to a 0.2-percent chance of flooding in any given year is normally called the 500-year floodplain. It is necessary to evaluate development in floodplains to ensure that the Project is consistent with the requirements of EO 11988, Floodplain Management.

As shown in Figure 3-4, portions of the Project Site would be located within the 100-year floodplains of North Branch Magowah Creek and several unnamed tributaries The following floodplain/floodway requirements have been previously established by Lowndes County for the Project Site:

Floodplains: Lowndes County requires that an elevation certificate be submitted for each individual solar panel tracking string. The elevation certificate must demonstrate that the lowest section of the module, at full tilt, is at least one foot above the 100-year base flood elevation.

Floodways: Lowndes County requires that a no-rise condition be reflected through hydraulic analysis in order to avoid the requirement of a Conditional Letter of Map Revision (CLOMR). If a no-rise condition cannot be met, a CLOMR must be obtained as the best mitigation approach to prevent major impacts on flood heights.

There are no FEMA Regulatory Floodways associated with GT2.




Figure 3-4: Floodplains within the Project Site

# 3.3.2 Environmental Consequences

#### 3.3.2.1 No Action Alternative

Under the No Action Alternative, no Project-related impacts on water resources would occur. Existing land use would remain as farmland and surface waters would remain as they are at the present time. Increases in erosion and sediment runoff could occur over time if best-practices in agriculture were not maintained to prevent erosion and runoff. In addition, if broad applications of chemical fertilizers or pesticides are continually used, this could result in nutrient-rich runoff that degrades the quality of surface waters within the site and throughout the broader drainage basin.

## 3.3.2.2 Proposed Action

Under the Proposed Action, minor short-term impacts from construction would be expected on streams, wetlands, and floodplains from sedimentation of exposed soils. Standard BMPs would be installed, inspected, and maintained until satisfactory stabilization is achieved. Beneficial, indirect impacts to surface water and groundwater would result from a reduction in broad applications of pesticides, herbicides, and fertilizers used in support of the current agricultural land use activities. Additionally, water quality may be improved through filtering by native plant cover as opposed to crop cover, which could reduce erosion and sedimentation from stormwater events.

# Groundwater

During construction, hazardous materials would be on-site that could potentially contaminate groundwater resources, including petroleum products for fuel and lubrication of construction equipment, hydraulic fluids, and a variety of other chemicals commonly used for general construction projects. Implementation of BMPs would provide measures to minimize potential for leaks or spills from construction equipment and outline procedures and protocols to quickly address potential spills that may occur. Construction activities would be in accordance with BMPs outlined in TVA's A Guide for Environmental Protection and Best Management Practices for Tennessee Valley Authority Construction and Maintenance Activities (TVA, 2017b) to avoid contamination to groundwater. Fertilizers and herbicides are not proposed to be used during construction or operation of the Proposed Action which would be considered a direct long-term benefit to groundwater. However, if minor application of fertilizer is needed for initial re-vegetation, applications would be in accordance with the manufacturer's recommendation and would be short-term. Currently, most of the land proposed for use is actively cultivated cropland with frequent application of fertilizers, herbicides, and pesticides. Change in land use from agriculture to solar would be a long-term beneficial impact to groundwater.

Water needed for construction would be provided from existing or proposed temporary groundwater wells or water delivery trucks. The construction contractor would use water for the purposes of fugitive dust mitigation (during dry conditions), concrete mixtures, and other temporary construction needs. If practicable, groundwater wells and holding tanks would reduce and avoid impacts to groundwater. Construction activities requiring water would primarily be for

dust control and compaction during grading activities for access roads, pads, and foundations for structures.

Water usage within the Project Site during operations would be in accordance with BMPs outlined in TVA's *A Guide for Environmental Protection and Best Management Practices for Tennessee Valley Authority Construction and Maintenance Activities*. Local rainfall is generally consistent enough to avoid the need for dust control on PV arrays. Therefore, regular panel washing is not anticipated. Water needs during the operation and maintenance phase of the Proposed Action, would be addressed by new or existing groundwater wells. Groundwater withdrawal volumes are expected to be less than the existing volume needed for agricultural irrigation, thus resulting in a net positive impact on groundwater resources.

Broad application of fertilizers and herbicides are not proposed during construction or operation of the Project, which would be considered a direct long-term benefit to groundwater. However, if minor applications of fertilizer or herbicides are needed for initial re-vegetation or maintenance, applications would be in accordance with manufacturer's recommendations and would be short-term. Currently, most of the land proposed for use is actively cultivated cropland with frequent application of fertilizers, herbicides, and pesticides. Change in land use from agriculture to solar would therefore be a long-term beneficial impact to groundwater.

If the Solar Facility were to be decommissioned or closed, a Decommissioning and Closure Plan would be developed. The Decommissioning and Closure Plan would detail procedures to control erosion and sedimentation and maintain compliance with NPDES requirements and permits. Water usage for potential decommissioning and closure is not likely to exceed that used for operation and maintenance. Therefore, impacts to groundwater resulting from a decommissioning and closure of the Solar Facility are not anticipated.

There were no sinkholes identified within the Project Site, therefore the potential for direct surface to groundwater contamination from stormwater or chemical and solid waste runoff is not anticipated. Herbicide and pesticide applications are not expected to be used during construction or operation of the Project. However, if pesticides or herbicides would be required at any point during construction or maintenance activities, applications would be consistent with Mississippi Department of Agriculture and Commerce permit requirements. Proper application of herbicides and pesticides, if needed, may result in minor impacts to surface water but would be significantly less than applications from current land use.

Overall, impacts to local aquifers and groundwater are not anticipated due to the limited volume of groundwater required for initial construction, operation, and maintenance, or decommissioning and closure of the Solar Facility. BMPs and a Decommissioning and Closure Plan would reduce the potential for hazardous materials to reach groundwater resources during any stage of the Project.

Additionally, minor, indirect beneficial impacts to groundwater could occur from the discontinued use of broad applications of herbicides, pesticides, and fertilizers, due to change in land use from agriculture to solar.

# Surface Water

Due to capacity requirements and land constraints, complete avoidance of jurisdictional water features was not practicable with the Proposed Action. Due to the need to cross certain streams within GT2, temporary and permanent access road crossings would be necessary. The installation of temporary and permanent bridges and/or culverts could affect up to 300 linear feet of intermittent and perennial streams. During all stages of the design process, efforts have been made to avoid and minimize impacts to jurisdictional wetlands and waterbodies to the greatest extent practicable. New stream crossings would, to the extent practicable, occur at a 90-degree angle to the stream. Best Management Practices for stream crossings would be implemented, such as using a bridge span that avoids direct impacts to the stream beds and top of bank, placing geotextile fabric along bridges to minimize dirt and debris from entering the stream channel, and minimizing the width of each crossing to the smallest width allowable (within safety requirements).

The Proposed Action would result in minor, direct, permanent impacts to jurisdictional streams at locations where permanent culverts are installed for construction of access roads. While complete avoidance of stream features was not possible, with implementation of the above mentioned BMPs, the USACE and MDEQ Section 404 and 401 permit requirements, and the Project's MDEQ-approved SWPPP to control erosion and sediment runoff, impacts to streams would be short-term and minimal.

Table 3-3 provides a breakdown of potential impacts on streams due to facility access road crossings. Currently, MS Solar 6 anticipates the need for eight stream crossings, totaling approximately 160 linear feet of impacts. MS Solar 6 intends to limit stream crossings to 300 total linear feet (or less) in order to comply with the USACE and MDEQ impact limits under NWP 3 and/or NWP 14. Existing access roads and bridges would be prioritized with construction of new waterbody crossings and culvert installation only being used where necessary.

Table 3-3: Waterbodies Potentially Affected by GT2
Access Road Crossings

Waterbody ID	Waterbody Type	Top of Bank Width (feet)	Length of Stream (feet) in Project Site	Impact Type	Length of Impact (feet)	Area of Impact Sq. ft.
S-29	Intermittent	5	3,430	bridge/culvert	20	100
S-05	Perennial	5	4,243	bridge/culvert	20	100
S-16	Perennial	20	11,766	bridge/culvert	20	400
S-16	Perennial	20	11,700	bridge/culvert	20	400
S-33	Perennial	6	4,837	bridge/culvert	20	120
S-33	Perennial	6	4,037	bridge/culvert	20	120
S-23	Perennial	25	2,790	bridge/culvert	20	500
S-21	Intermittent	2	378	bridge/culvert	20	40
				Total Impact	160	1780

## Wetlands

Impacts on wetlands would be avoided to the extent practicable. As the current layout shows, the area of impact has been designed to avoid any impacts on wetlands entirely. The placement of permanent aboveground facility components, such as PV panels, inverters, and access roads were limited to upland areas. Minor impacts from stormwater discharges could occur to wetlands within the Project Site. However, buffers of at least 15 feet have been established around wetlands; and BMPs such as silt fencing would be installed, inspected, and maintained along the perimeter of the construction area where wetlands are present. In some cases, buried tie-in lines may be required to connect one area of panels to another. To avoid disturbances to wetland features at the Project Site, MS Solar 6 would bore under wetlands as necessary to install collection cables. If a wetland crossing were identified by the construction contractor, the appropriate permits and clearances would be obtained through the USACE Mobile Regulatory District and the Mississippi DEQ. Figure 3-5 shows the preliminary array layout and delineated features within the Project Site. As shown, MS Solar 6 would avoid jurisdictional wetlands and waterbodies to the greatest extent practicable.




Figure 3-5: Surface Water and Wetlands within the Project Site

## Floodplains

As a federal agency, TVA adheres to the requirements of EO 11988, Floodplain Management. The objective of EO 11988 is "... to avoid to the extent practicable the long- and short-term adverse impacts associated with the occupancy and modification of floodplains and to avoid direct and indirect support of floodplain development wherever there is a practicable alternative..." The EO is not intended to prohibit floodplain development in all cases, but rather to create a consistent government policy against such development under most circumstances (United States 1978). The EO requires that agencies avoid the 100-year floodplain unless there is no practicable alternative.

Based on Lowndes County, Mississippi, FIRM panels 28087C0225K and 28087C0250K, both effective 2/18/2011, and aerial maps, access roads, buried collection lines, security fencing, security lighting, and portions of the overhead wire would be located within 100-year floodplains. Consistent with EO 11988, these structures and activities are considered to be repetitive actions in the 100-year floodplain, which would result in minor impacts (TVA 1981).. To minimize adverse impacts, any roads within 100-year floodplains would be constructed such that upstream flood elevations would not increase by more than 1.0 foot.

Portions of the solar arrays would also be located in 100-year floodplains. As depicted in Figure 2-2, the preliminary layout of solar arrays has been designed to minimize the number of PV panels installed within floodplains. However, a small section of arrays would be installed within the floodplain. The location of arrays within the floodplain was determined based on the results of MS Solar 6's Hydrology Study (Appendix C). Solar arrays are not considered to be repetitive actions in the 100-year floodplain (TVA 1981). Due to existing site constraints, including cultural resources, surface waters, and property reserved for potential future development, about up to acres of solar panels cannot be located outside of the 100-year floodplain. Further, the number of proposed panels are required to meet the PPA between GT2 and TVA. Therefore, TVA has determined that there is no practicable alternative to locating about up to 50 acres of solar panels in the 100-year floodplain. To minimize adverse impacts, the solar panels would be placed at a height such that the lowest section of the solar module, at full tilt, would be located at least one foot above the BFE (base flood elevation; based on the Hydrology Study for the Project Site).

During construction, the temporary laydown areas, construction trailers, soil stockpiles, and parking areas would be located outside the floodplain, which would be consistent with EO 11988. MS Solar 6 would employ stabilization measures to prevent erosion of stockpiled soil.

MS Solar 6 would need new access roads throughout the Solar Facility in order to access arrays and inverters during construction and operation. MS Solar 6 would utilize existing farm roads to the greatest extent practicable. The exact location of any new compacted earth access roads is not yet final. However, it is anticipated that at least one new access road would be needed within the 100-year floodplain. Lowndes County participates in the National Flood Insurance Program, and any development must be consistent with its floodplain regulations. MS Solar 6 has discussed with Lowndes County the potential Project activities within the 100-year floodplain. Prior to

construction, a review by the Lowndes County Floodplain Administrator would occur; and the appropriate permissions would be obtained for work and facilities within the floodplain. Any access roads would be designed and permitted as compacted earth facility roads. Also, access roads within 100-year floodplains would be constructed such that upstream flood elevations would not be increased by more than 1.0 foot.

By adhering to the following mitigation measures, the proposed GT2 solar facility and TVA PPA would have no significant impact on floodplains and their natural and beneficial values.

- Access roads within 100-year floodplains would be constructed such that upstream flood elevations would not be increased by more than 1.0 foot.
- The portions of the fencing within the floodplain should be designed to withstand flooding with minimum damage.
- Any excavated material or debris generated when GT2 is decommissioned and dismantled would be disposed of in an area outside 100-year floodplains.
- The lowest section of each solar module, at full tilt, will be at least one foot above the 100year base flood elevation.
- At the end of the Project's useful life, non-recyclable and/or non-reusable debris would be disposed of at an appropriate location outside 100-year floodplains.
- Any permanent operations and maintenance building(s) will be located outside 100-year floodplains and elevated to at least one foot above the 100-year floodplain at that location.

# 3.4 Biological Resources

This section provides an overview of the existing biological resources within the Project Site and the potential impacts to those resources from implementation of the No Action and Proposed Action Alternatives. Biological resources analyzed in this section include natural areas, vegetation, wildlife, and rare, threatened, and endangered species.

The Project is within the EPA Blackland Prairie Ecoregion (Level IV) and is within the Middle Tombigbee River [Hydrologic Unit Code (HUC) 03160106] and Tibbee Creek (HUC 03160104) watersheds. Blackland prairie ecoregions usually consist of gently rolling hills with little bluestem (*Schizachyrium scoparium*), big bluestem (*Andropogon geradii*), Indianagrass (*Sorghastrum nutans*), eastern gamma grass (*Tripsacum dactyloides*), switchgrass and sidesoats grama (*Bouteloua curtipendula*) with sparse pockets of pecan (*Carya illinoinensis*), cedar elm (*Ulmus crassifolia*), hackberry (*Celtis occidentalis*), and a variety of oak trees (*Quercus sp.*). Average temperatures are usually between 66- and 70-degrees Fahrenheit and rainfall averages 30 to 40 inches annually (Texas Parks and Wildlife, 2021). Agriculture products such as corn, soybeans, and hay are produced in large quantities within the Project Site and surrounding area.

Prior to conducting field work, biologists reviewed the USFWS Information for Planning and Consultation (IPaC) for the Project Area (Appendix D), regarding special status species that may occur within the Project Site (Consultation Code: 04EM1000-2021-SLI-0745, Event Code: 04EM1000-2021-E-01680) and assessed whether the proposed Project had potential to affect ESA species (i.e., ESA listed, proposed and candidate species), bald eagles (*Haliaeetus leucocephalus*), golden eagles (*Aquila chrysaetos*), migratory birds (including raptor species), and associated habitat within the Survey Area. Additionally, consultation was conducted with Mississippi Department of Wildlife, Fisheries, and Parks' (MDWFP) National Heritage Program which provided a list of state protected species and state ranked species of special concern with potential to occur within two miles of the Survey Area. U.S. Fish and Wildlife's Range Wide Indiana Bat Survey Guidelines (USFWS, 2019) was also reviewed. Lastly, TVA's regional Natural Heritage Database was reviewed which provides known occurrences of rare species within three and five miles of the Project Site for terrestrial and botanical species, respectively.

Field surveys were conducted from March 7 through March 13, 2021. During the field surveys, data was collected on vegetative cover/land use and protected species habitats. Regulations for biological resources potentially relevant for the Proposed Action include:

- ESA (16 U.S.C. §§ 1531-1544).
- Migratory Bird Treaty Act (MBTA) of 1918 (16 U.S.C. §§ 703-712) (for actions of nonfederal entities).
- Executive Order for Migratory Birds (EO 13186 of January 10, 2001) (for actions of federal agencies).
- Bald and Golden Eagle Protection Act (BGEPA) (16 U.S.C. §§ 668-668c); and
- Mississippi Nongame and Endangered Species Conservation Act (Miss Code Ann. §§ 49-5-101 to 49-5-119).

Additionally, TVA reviewed data from the regional Natural Heritage Database for federal and state protected species occurring within or near the Project Site or generally listed for Lowndes County, MS.

## 3.4.1 Affected Environment

Biological resources within the Project Site include natural areas, vegetation, wildlife, and the potential for rare, threatened, or endangered species.

#### 3.4.1.1 Natural Areas

Natural areas include managed areas such as Wildlife Management Areas, National Wildlife Refuges and Habitat Protection Areas, ecologically significant sites, and river segments listed in the Nationwide Rivers Inventory. There is only one natural area within three miles of the Project. The closest natural area to the Project Site is the Black Prairie Wildlife Management Area, which is approximately 2.1 miles to the southeast (MDWFP 2022). The Black Prairie Wildlife Management Area, is managed and maintained by MDWFP, and is used primarily for deer hunting and wildlife viewing.

## 3.4.1.2 Vegetation

The Project Site is located within the Black Belt Prairie (also known as the "Blackland Prairie" and the "Cotton Belt"), which is a subdivision of the East Gulf Coastal Plain physiogeographic province. The Black Belt Prairie is best known for its dark, low chroma soils and high crop yield; however, prior to the influx of farming practices, there were three dominant plant communities that naturally occurred in this region: open prairie, chalk outcrop, and forests (MacGown, Brown, and Hill 2006). Naturally occurring, mature hardwood forests are found within and beyond the Project Site.

Figure 3-6: Chalk Outcrop on Parcel Associated with the Project Site



Chalk outcrops occur in areas within the Black Belt Prairie where erosion has exposed the underlying formation. Two outcrops were identified within a parcel of land that would be leased to MS Solar 6 as part of the Project Site. These scattered outcrops, which have undergone severe erosion, add up to approximately 12 acres in total. Chalk outcroppings are known to support several endemic and rare species of plants and insects. The photo to the left in Figure 3-6 was

taken within the Project Site in March 2021. Most of the Project Site on the south side of Artesia Road are within part of a prairie that is now in use as pastureland for cattle grazing (Phillips 2012). Very little undisturbed prairieland still exists today due primarily to human disturbances such as agricultural practices, conversion to pastureland for livestock, and other developments (Phillips 2012).

There are presently three major vegetation communities currently within the Project Site: active agriculture (row crops and pastureland), bottomland hardwood forests, and upland forests.

More than half of the Project Site is comprised of pastureland and approximately 9 percent is agricultural fields. Vegetation in these communities is maintained in an early successional state due to herbicide application, crop growth/harvesting, and cattle grazing. Corn is planted in late spring and cover the row crop fields. Vegetation observed in pastures consists of primarily tall fescue grass (Festuca arundinacea), Johnson grass (Sorghum halepense), annual bluegrass (Poa annua), couch grass (Elymus repens), bermuda (Cynodon dactylon), annual ragweed (Ambrosia artemisiifolia), cheatgrass (Bromus tectorum), perennial ryegrass (Lolium perenne), green foxtail (Setaria viridis), butterweed (Packera glabella), bulbous bittercress, soft rush, Cherokee sedge (Carex cherokeensis), fox sedge (Carex vulpinoidea), red sorrel (Rumex acetosella), curly dock, prairie fleabane (Erigeron strigosus), jimsonweed (Datura stramonium), Carolina horsenettle (Solanum carolinense), bull thistle (Cirsum vulgare), wild violet (Viola sororia), field garlic (Allium vineale), and sensitive partridge pea (Chamaecrista nictitans).

Bottomland hardwood forests are also present within the Project Site. These communities are composed of a tree canopy associated with a mature second-growth forest. Dominant vegetation observed consisted of water hickory (*Carya aquatica*), willow oak, cherrybark oak (*Quercus pagoda*), swamp chestnut oak (*Quercus michauxii*), silky dogwood (*Cornus amomum*), osage orange, green ash, eastern red cedar (*Juniperus virginiana*), water locust, shagbark hickory (*Carya ovata*), box elder (*Acer negundo*), red maple, American sycamore (*Platanus occidentalis*), sugarberry, Cherokee sedge, wild petunia (*Ruellia humilis*), tall fescue, poison ivy, greenbrier, Virginia spiderwort (*Tradescantia virginiana*), Virginia creeper, hairy buttercup (*Ranunculus sardous*), and resurrection fem (*Pleopeltis polypodioides*),

The upland forests within the Project Site are also mostly composed of a canopy age associated with a mature second growth forest. Dominant vegetation observed consisted white oak (*Quercus alba*), southern red oak (*Quercus falcata*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), mockernut hickory (*Carya tomentosa*), shagbark hickory, pignut hickory (*Carya glabra*), loblolly pine (*Pinus taeda*), eastern red cedar, American elm (*Ulmus americana*), honey locust (*Gleditsia triacanthos*), osage orange, Chinese privet, Devil's walkingstick (*Aralia spinosa*), Christmas fern (*Polystichum acrostichoides*), Japanese honeysuckle (*Lonicera japonica*), multiple greenbrier species (*Smilax* spp.), muscadine (*Vitis rotundifolia*.), Virginia creeper, blackberry (*Rubus* spp.), wooly panic grass (*Dichanthelium acuminatum*), and wild violet. The upland forest in the southwest corner of the Project Site is primarily composed of eastern red cedar.

#### 3.4.1.3 Wildlife

Surveys for protected species and habitat assessments were conducted from March 7 through March 13, 2021. The entire Project Site was surveyed simultaneously during the wetland delineation survey. In addition to habitat assessment, detailed vegetative communities were described. Areas within the Project Site that are not currently used for agriculture may provide suitable habitat for wildlife common to the region both seasonally and year-round.

Mammals commonly found throughout Mississippi include coyotes (*Canis latrans*), white-tailed deer (*Odocoileus virginianus*), otters (*Lontra canadensis*), beavers (*Castor canadensis*), black bears (*Ursus americanus*), skunks (*Mephitis mephitis*), opossums (*Didelphis virginiana*), racoons (*Procyon lotor*), squirrels (*Sciurus* spp.), and armadillos (*Dasypus novemcinctus*) (MDWFP 2021). During field surveys for protected species and habitat assessments, evidence of deer, raccoon, squirrel, opossum, and armadillo were physically observed and generally confined to the forested areas. Observations of deer and deer tracks were made throughout the site.

Reptiles and amphibians commonly found in the region include a variety of turtles, lizards, frogs, and snakes. Fence lizards (*Sceloperus* spp.), five-lined skinks (*Plesitiodon fasciatus*), and anoles (*Anolis* spp.) are commonly observed lizards. Chorus frog (*Pseudacris* sp.), cricket frog (*Acris gryllus*), green tree frog (*Hyla cinerea*), bullfrog (*Lithobates catesbeianus*), American toad (*Anaxyrus americanus*), eastern spadefoot toad (*Scaphiopus holbrookii*), marbled salamander (*Ambystoma opacum*), spotted salamander (*Ambystoma maculatum*), and dusky salamander (*Desmognathus fuscus*) are often observed in the region. Non-venomous snakes include garter snake (*Thamnophis sirtalis*), king snake (Lampropeltis spp.), rat snake (*Pantherophis* spp.), and

water snake (*Nerodia* spp.), while venomous species include coral snake (*Micrurus fulvius*), cottonmouth (*Agkistrodon piscivorus*), copperhead (*Agkistrodon contortrix*), pygmy rattlesnake (*Sistrurus miliarius*), and diamondback rattlesnake (*Crotalus adamanteus*) (MDWFP, 2021). Several observations of garter snakes, red-eared sliders (*Trachemys scripta*), *upland chorus frogs* (*Pseudacris feriarum*), a cottonmouth, an eastern king snake (*Lampropeltis getula*), and a banded water snake (*Nerodia fasciata*) were observed during field surveys. No other specific observations were noted for reptiles or amphibians.

Birds commonly found in the region include wild turkey (*Meleagris gallopavo*), northern mockingbird (*Mimus polyglottos*), red-winged blackbird (*Agelaius phoeniceus*), northern cardinal (*Cardinalis cardinalis*), American crow (*Corvus brachyrhynchos*), mourning dove (*Zenaida macroura*), red-tailed hawk (*Buteo jamaicensis*), American robin (*Turdus migratorius*), starling (*Sturnus vulgaris*), brown thrasher (*Toxostoma rufum*), and turkey vulture (*Cathartes aura*) (Sibley, 2017). Flyovers from red-tailed hawk, American crow, and black vultures were observed during field surveys. Observations of American robin, American crow, northern mockingbird, field sparrow (*Spizella pusilla*), cedar waxwing (*Bombycilla cedrorum*), downy woodpecker (*Dryobates pubescens*), killdeer (*Charadrius vociferus*), great horned owl (*Bubo virginianus*) and mallard duck (*Anas platyrhynchos*) were also made throughout the Project Site. Wading birds to include great blue heron (*Ardea herodius*) and great egrets (*Ardea alba*) were observed near the norther portion of the project site near the aquaculture ponds across from Gilmer Wilburn Rd.

No caves or mines were identified on the Project Site during field surveys. Additionally, in speaking with landowners during site visits, no known caves or mines exist in proximity to the Project Site.

## 3.4.1.4 Threatened, Endangered, and Rare Species

The USFWS IPaC result letter (Appendix D) was obtained from the IPaC website in regard to special status species that may occur within the Survey Area (Consultation Code: 04EM1000-2021-SLI-0745, Event Code: 04EM1000-2021-E-01680). This IPaC result letter identified federally protected species or Designated Critical Habitats that occur or have the potential to occur in the Survey Area ((i.e., Endangered Species Act of 1973 (ESA) listed, proposed and candidate species), bald eagles, golden eagles, and migratory birds (including raptor species)). Additionally, consultation was conducted with MDWFP National Heritage Program which provided a list of state protected species and state ranked species of special concern with potential to occur within two miles of the Survey Area.

Three federally listed species were identified as potentially occurring within the Survey Area: the northern long-eared bat (*Myotis septentrionalis*), wood stork (*Mycteria americana*), and Price's potato-bean (*Apios priceana*). Critical habitat for federally protected species has not been designated within Lowndes County, Mississippi. The Mississippi Natural Heritage Program is managed under the MDWFP, Museum of Natural Science. The state of Mississippi does not designate state-statuses for plants, but MDWFP does track plant species there are rare in Mississippi. However, there are aquatic species (mussels, fish, crayfish), amphibians, snakes, and birds that hold special state-endangered status. Several state-protected species were listed as

potentially occurring in the Survey Area. The federal and state protected species listed as having potential to occur in the Survey Area are provided in Table 3-4 and discussed in further detail below.

Table 3-4: Protected Species with Potential to Occur within the Survey Area

Common	Scientific	Federal	State	Preferred Habitat Description	Habitat
Name	Name	Status	Status Mamma	-	Present
Northern long- eared bat	Myotis septentrionalis	LT	-	Summer roosts occur in tree cavities and under exfoliating bark, but this species has also been found in buildings and behind shutters. During the winter, northern long-eared bats hibernate in tight crevices in caves and mines. Foraging is done primarily on forested hillsides and ridges	Yes
Eastern Spotted Skunk	Spilogale putorius	1	S1	Open prairies, brushy lands, and cultivated areas (MDC 2021)	Yes
			Birds		
Wood Stork	Mycteria americana	LT	LE	Freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches (USFWS 2020b)	Yes
Painted Bunting	Passerina ciris	ı	S3B	Dense weedy habitats as well as the dense understory of semi- open forests (Cornell 2019a)	Yes
Scissor-tail Flycatcher	Tyrannus forficatus	1	S1B; S1N	Open habitats in the southern Great Plains and south Texas, often around sparsely scattered trees or utility lines (Cornell 2019b)	Yes
			Plants		
Price's Potato Bean	Apios priceana	LT	-	Lightly disturbed areas such as forest openings, wood edges and where bluffs descend to streams	Yes
Big-head Evax	Evax <i>Prolifera</i>	1	S1	Meadows, grasslands, sagebrush steppe, and plains (MNHP 2021)	Yes
Slender Sedge	Carex gracilescens	-	S1	Mesic deciduous forest; often in sandy, rocky, and sometimes calcareous soils (WDNR 2020)	Yes
Stiff Greenthread	Thelesperma filifolium	-	S1	Disturbed sites in dry sandy or gravely soil (UTA 2018)	Yes
Nebraska Sedge	Carex jamesii	-	S1S2	Rich mesic woodlands, wooded slopes, wooded groves (where the ground vegetation is infrequently mowed underneath large trees), and edges of woodland paths (Hilty 2020a)	Yes
Big Shellbark Hickory	Carya Iaciniosa	-	S2	Rich alluvium of floodplains and rich mesic soil along riverbanks and marshes (NYNHP 2021)	Yes

Common Name	Scientific Name	Federal Status	State Status	Preferred Habitat Description	Habitat Present
Bur Oak	Quercus macrocarpa	-	S2	Moist woodlands and bottomland forests to prairies and sandhills (USFS 2021)	Yes
Ohio Buckeye	Aesculus glabra	-	S2	Moist to mesic deciduous woodlands, wooded valleys along rivers, and rocky wooded slopes in sheltered areas (Hilty 2020b)	Yes
Prairie Parsley	Polytaenia nuttallii	-	S2	Upland prairies, hill prairies, limestone glades, chert glades, thinly wooded bluffs, and savannas (Hilty 2018).	Yes
Prairie Pleatleaf	Nemastylis geminiflora	-	S2	Glades, prairies, and rocky slopes, on calcareous substrates – Occasionally on glades in the shade of eastern red cedars or Ashe's junipers (MDC 2020)	Yes
Small-toothed Sedge	Carex microdonta	-	S2	Dry to moist, calcareous substrates in open rocky or wet prairies, swales, seeps, and ditches - Relic patches of Blackland prairies, limestone glades, and chalk openings (FNA 2020)	Yes
Mead's Sedge	Carex meadii	-	S3S4	Moist to dry black soil prairies, dolomite prairies, hill prairies, savannas with sparse ground vegetation, sedge meadows, limestone glades, and areas along railroads (Hilty 2019a)	Yes
Nutmeg Hickory	Carya myristiciformis	-	S3S4	Banks of rivers and swamps in rich moist soils, occasionally on higher ground and often on limestone (PFAF 2021)	Yes

Key: Statuses are LE= Listed Endangered, LT= Listed Threatened State Rank

## Federally Listed Species

Three species listed as threatened under the ESA were identified as having potential to occur within the Project Site (Figure 3-7). None of the species described herein were observed during environmental field surveys, and no critical habitats were identified on the IPaC report (Appendix D).

S1 - Critically imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

S2 - Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

S3 - Rare or uncommon in Mississippi (on the order of 21 to 100 occurrences).

Sources: USFWS, 2021 and Mississippi Natural Heritage Program, 2021.

The wood stork (*Mycteria americana*) is a wading bird that is federally listed as threatened. It is found primarily in freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches. No nesting records are known from this area of Mississippi (USFWS 2020b). Foraging habitat for the species was observed within all the open waters observed throughout the Survey Area.

The northern long-eared bat (*Myotis septentrionalis*) (NLEB) is federally listed as threatened under the ESA. It usually roosts in tree cavities and under exfoliating bark, but this species has also been found in buildings and behind shutters. During the winter, NLEBs hibernate in tight crevices in caves and mines. Foraging is done primarily on forested hillsides and ridges (USFWS 2015). No records of this species are known from Lowndes County, Mississippi. Approximately 358 acres of suitable summer roosting habitat for the NLEB was observed within forested areas at the Project Site, and suitable foraging habitat was observed within the perennial stream corridors, fence rows, and forests throughout the Project Site. An acoustic survey was approved by USFWS and performed throughout the suitable habitat following the 2020 U.S. Fish and Wildlife *Range-wide Indiana Bat Survey Guidelines*. Acoustic monitors were run Aug 3-5 at the Level of Effort dictated by the Guidelines (26 detector nights). NLEB was determined to be likely absent based on automated identification results.

Price's potato-bean (*Apios priceana*) is a plant that is federally listed as threatened under the ESA. It prefers lightly disturbed areas such as forest openings, wood edges and where bluffs descend to streams. It also grows along highway ROWs and powerline corridors (USFWS 2019). Potential habitat for Price's potato bean is present in central portion of the Project site within some ravines and hillslopes and the nearby forested bluffs along Gilmer Creek and its tributaries, adjacent to the central western portion the Project Site. Although potentially suitable habitat is present within the central portion of the Project Site, there were no observations of Price's potato bean during targeted botanical surveys.




Figure 3-7: Potentially Suitable Habitat for Federally Protected Species within the Project Site

#### State Listed Mammals

The eastern spotted skunk is a small weasel-like skunk with dense fur baring stripes and spots in a distinctive pattern. This skunk can be found across the central and southeastern United States and northeastern Mexico (Bullock 2008). In Mississippi, the eastern spotted skunk's status is S1: Critically imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation. Typical habitat preferences include brushy, rocky, woody habitats with extensive vegetation; and they are rarely identified in open areas or around wetlands. No observations of eastern spotted skunks occurred during field surveys; however, they are more likely to be active at night. If the eastern spotted skunk did occur within the Project area, it would likely prefer the large, dense forest just south of North Branch Mogawah Creek, which is located at the very southern boundary of the Project Site.

## Bald Eagles and Migratory Birds

In Mississippi, the bald eagle is protected under the Bald and Golden Eagle Protection Act (BGEPA) and the Migratory Bird Protection Act (MBTA). No bald eagles or nests were confirmed to be present during the environmental field surveys within the Survey Area or along public roadways near the Project. However, one potential bald eagle nest was reported to field crews by a landowner. The potential nest is within private property outside of the Survey Area. The nest was observed from a distance in an adjacent parcel outside of the Survey Area. Photographs were attempted but were unable to provide value due to distance and obstruction from existing mature trees. The field crews were unable to confirm whether the nest was that of a bald eagle or some other bird of prey. The status of the nest is unknown, and no individuals were observed. No blasting or other extremely loud noises are anticipated during construction. At its closest point, Project-related activities would occur within approximately 1,200 feet from the potential nest and would be in compliance with National Bald and Golden Eagle Management Guidelines.

There are certain birds that are protected under the MBTA as Birds of Conservation Concern (BCC). In the April 2021, USFWS IPaC Report for the Survey Area, no USFWS-designated BCCs were identified. However, numerous migratory birds were identified through the TVA's Natural Heritage Database. Migratory birds of conservation concern identified in TVA's natural heritage data include American kestrel, bald eagle, cerulean warbler, dunlin, Kentucky warbler, lesser yellowlegs, marbled godwit, prothonotary warbler, red-headed woodpecker, rusty blackbird, short-billed dowitcher, swallow-tailed kite, willet, and wood thrush. Additionally, MDWFP indicated that scissor-tail flycatcher and the painted bunting could also occur within two miles of the Survey Area. Both species are state species of concern.

The American Kestrel is a small member of the falcon family and prefers habitats consisting of open farmland, wood edges, and cities. It breeds from April through August most likely in areas outside the Project Site (Cornell 2019c). It is most commonly found in Lowndes County, Mississippi, during December and January. No individuals or nests were observed during field surveys; however, suitable nesting and foraging habitat does occur within the Survey Area.

Ospreys (*Pandion haliaetus*) are large birds of prey that typically nest in elevated nest sites near rivers, lakes, reservoirs, lagoons, swamps, and marshes with an adequate supply of accessible fish within about twelve miles of the nest (Cornell 2019d). According to TVA's natural heritage data, there is a record from 2011 of an osprey's nest approximately 2.7 miles from the Survey Area. No nests were observed during field surveys; however, osprey could exist near the Survey Area due to the proximity of the aquaculture ponds.

Lesser yellowlegs (*Tringa flavipes*) is a small "piper" bird that breeds in meadows and open woodlands in Canada. In North America, lesser yellowlegs typically occurs in marshes, shallow wetlands, shorelines, and flooded fields during migration. Most often, they prefer vegetated wetlands rather than bare habitats (Cornell 2019e). Marginal foraging habitat is present at PEM wetlands throughout the Project Site. PEM wetlands within the Survey area were low quality, isolated, and are regularly disturbed during crop and hay harvesting.

Marbled godwit (*Limosa fedoa*) is a small shorebird with a two-toned, long, slightly upcurved bill that breeds in the shortgrass prairies near wetlands in Canada and the northern central U.S. While migrating south, marbled godwits typically inhabit native grass prairies with green needle grass, western wheatgrass, blue gram, needle-and-thread, and blue stem. Wintering grounds typically consist of coastal mudflats, estuaries, and sandy beaches (Cornell 2019f). The migration corridor for the marbled godwit is further west of the Survey Area, closer to Texas and Arkansas. Due to lack of suitable habitat and the location of the Survey Area outside of the migration corridor, marbled godwit is unlikely to occur within the Survey Area.

Red-headed woodpecker (*Melanerpes erythrocephalus*) is a medium-sized woodpecker with a bright red head, white buff, and black back. Red-headed woodpeckers breed in river bottoms, beaver swamps, burned areas, recent clearing, deciduous woodlands with oak or beech and groves of dead or dying trees. Dead or partially dead trees are important for nest cavities in areas where they breed. In Mississippi, red-headed woodpeckers typically inhabit pine and pine-oak forests (Cornell 2019g). Most of the undisturbed forested areas within the Survey Area would provide suitable habitat for red-headed woodpeckers.

As its name entails, the rusty blackbird (*Zanata Canadiense*) has distinctively rusty feather edges and pale-yellow eyes. Rusty blackbirds typically breed in wet forests, swamps, bogs, and beaver ponds. They also winter in swamps, woodlands, and pond edges. This species is considered a common bird in steep decline due to habitat loss (Cornell 2019h). Potentially suitable habitat for rusty blackbird is present near the pond edges and riparian woodlands to the open ponds.

Short billed dowitcher (*Limnodromus griseus*) is a full-bodied orange, brown, and golden shorebird with a long bill used for rhythmic probing of mudflats and flooded field for food. Short-billed dowitchers breed in the taiga shield ecotone, nest in wetlands usually near edges of bogs, small lakes, or wet meadows. During winter months, short-billed dowitchers are founds in saltwater and brackish environments such as estuaries and lagoon with tidal influences (Cornell 2019i). Marginal foraging habitat for migrant short-billed dowitchers is present in some of the shallow ponds and potentially flooded farm fields when excessive precipitation occurs.

Willet (*Tringa semipalmata*) is a straight billed, mottled brown, shorebird with distinct wing markings and a piercing call. Willets typically inhabit open beaches, marshes, mudflats, and rocky coastal zones. During the breeding season eastern willets are typically found in saltmarshes, barrier islands, and barrier beaches (Cornell 2019j). Marginal foraging habitat for migrant willets is present in some of the shallow ponds and potentially flooded farm fields from excessive precipitation.

MDWFP's Natural Heritage Data also identified two migratory birds with potential to occur within two miles of the Survey Area:

Scissor tail flycatcher (*Tyrannus forficatus*) was identified by the MDWFP as potentially occurring within two miles of the Survey Area. Scissor tail flycatchers prefer open habitats in the southern Great Plains and south Texas, often around sparsely scattered trees, or utility lines (Cornell 2019b). Open pasture within the Survey Area provides marginal foraging habitat for the scissortailed flycatcher.

Painted Bunting (*Passerina ciris*) was also identified by the MDWFP as potentially occurring within two miles of the Survey Area. Painted bunting prefers dense weedy habitats as well as the dense understory of semi open forests (Cornell 2019j). Although they are typically more often found further south and west, most of the forested areas within the Survey Area provide foraging habitat.

If tree clearing activities associated with construction of the Project overlap with the primary nesting season (March 15 – September 15), short-term inadvertent impacts could occur on bird species that nest in or near the construction areas. To the extent practicable, Project construction will be scheduled to minimize potential effects to bird species by minimizing clearing activities within forested areas during nesting season. Other mitigation measures MS Solar 6, LLC plans to implement include:

- Designing Project facilities to avoid sensitive resources where possible.
- Maximizing locations where the Project utilizes agricultural areas.
- Limiting the construction and operation workspaces to the minimum necessary.
- Conducting mitigation for effects to sensitive resources (e.g., wetlands) through agency permit conditions.
- Avoiding forested areas, to the greatest extent practicable.
- Minimizing routine mowing/maintenance during the bird nesting season once the Project is complete (generally March 15 through September 15 in the general Project vicinity) to the extent feasible.

On Dec. 22, 2017, the U.S. Department of the Interior (DOI) revised its guidance on incidental take of migratory birds in Memorandum M-370501, which specifies that incidental take prohibitions apply only to actions that have as their purpose the taking or killing of migratory birds. Under the Biden administration, the Department of the Interior withdrew its 2017 legal opinion

which preceded and formed the basis of the guidance. The new administration may return policies initiated under the Obama administration, which considered a proposal to develop an incidental take program. Because the purpose of MS Solar 6 is the lawful construction of a clean energy facility, and not the intentional take of migratory birds, MS Solar 6 does not anticipate further coordination with USFWS regarding migratory birds; however, legal developments to MBTA should be monitored.

Terrestrial species documented in TVA's natural heritage data included northern long-eared bat, wood stork, osprey and bald eagle which are discussed above. Mitchell's satyr (*Neonnympha mitchellii*) was also identified has having the potential to occur within the Survey Area; however, that species has not been documented within three miles of the project area or from Lowndes County, Mississippi.

# Mississippi State-Listed Species

The Mississippi Natural Heritage Program is managed under the MDWFP, Museum of Natural Science. Consultation was initiated with the MDWFP on February 25, 2021, and database results were issued on March 18, 2021, which indicate occurrences of state listed species or species of special concern that are known to occur within two miles of the Survey Area. Results of that correspondence are in Appendix C. According to the results, one state-listed species, the federally threatened wood stork, and 15 state ranked species of special concern have potential to occur within the Survey Area. Furthermore, MDWFP concluded that if best management practices are properly implemented, monitored, and maintained, the proposed project likely poses no threat to listed species or their habitats. Each of the species and a description of its preferred habitat is provided below. The state protected species identified are listed in Table 3-4 and discussed in further detail below.

# Rare Plants

As of 2018, there are 495 plants of special concern in Mississippi. The designation of plants of special concern in Mississippi was developed for flagging sensitive species that may be adversely affected by proposed projects, determining protection priorities of natural areas that contain special plants, and determining priorities for inventory and protection for special plants to include proposing special plants for federal protection (Mississippi Natural Heritage Program, 2018).

Twenty-eight rare plant species were included in the results of the TVA Heritage Database search as potentially occurring within five miles of the Survey Area. Because MS Solar 6 is avoiding most of the wetland and stream riparian areas and chalk outcroppings and associated buffers from the construction and operation footprint, suitable habitat only exists for 13 of these plant species within the Project Site. The table below provides additional details for these 13 species.

Table 3-5: TVA Natural Heritage Database - Plants Potentially Occurring within the Survey Area

Common Name	Scientific Name	Rank	Preferred Habitat Description
Ridgestem false foxglove	Agalinis oligophylla	S2	Typically found from east Texas to southwest Louisiana on coastal plains primarily in prairies – Prefers moist to dry sand or clay soils and also found in pine hardwood forest and pine savannahs (USGS 2020)
Wild hyacinth	Camassia scilloides	S2	Black soil prairies, moist savannas, moist open woodlands along stream banks, rocky wooded slopes, and limestone glades - Typically prefer moist conditions to full sun to light shade in rich loamy soil (Hilty 2020c)
James's Sedge	Carex jamesii	S1S2	Mesic woodlands, wooded slopes, wooded groves, and edges of woodland paths - May also occur in upland woods and swampy woodlands and occasionally found in degraded woodland habitats (Hilty 2020a)
Small-toothed sedge	Carex microdonta	S3	Dry to moist, calcareous substrates in open rocky or wet prairies, swales, seeps, and ditches - Relic patches of Blackland prairies, limestone glades, and chalk openings (FNA 2020)
Kingnut Hickory	Carya laciniosa	S2	Bottomland woodlands, upland woodlands, swamps, savannas, and limestone glades. Usually not far from rivers (Hilty 2020d)
Scarlet Indian- paintbrush	Castilleja coccinea	S1	Sandy soils in prairies and open woods (USFS 2020)
Pumpkin ash	Fraximus profunda	S3	Moist wet conditions often in swamps or floodplains, wet bottomlands, river valleys, and low areas. Often in bald cypress swamps, cottonwood, and tupelo swamps (MBG 2020a)
Canada moonseed	Menispermum canadense	S3	Deciduous woodlands, woodland borders, thickets, semi-shaded riverbanks, cleared powerline ROWs, overgrown fencerows, and hedges (Hilty 2020e)
Prairie pleatleaf	Nemastylis geminiflora	S2	Glades, prairies, and rocky slopes, on calcareous substrates – Occasionally on glades in the shade of eastern red cedars or Ashe's junipers (MDC 2020)
Buroak	Quercus macrocarpa	S2	Bottomland soils in woodland and stream valleys (MBG 2020b)
Heath aster	Symphyotrichum ericoides	S2	Mesic to dry black soil prairies, gravel prairies, dolomite prairies, hill prairies, savannas, openings in dry rocky forests, limestone glades, roadsides and railroad sides, and pastures (Hilty, 2020f)
Southern Meadow Rue	Thalictrum debile	S1S2	Floodplain forests over calcareous substrates – prefers rich, rocky limestone woods often near streams (Chafin 2007).
Great Plains Ladies'-tresses	Spiranthes magnicamporum	S2	Open prairie and higher pH soils in both low and, moist areas and higher, gravelly sites (Minnesotawildflowers 2021)

Common Name	ientific Name	Rank	Preferred Habitat Description
----------------	---------------	------	-------------------------------

S1 = Critically Imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation.

S2 = Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

S1S2 = Rank range between S1 and S2.

S3 = Vulnerable in Mississippi due to a restricted range (on the order of 21 to 100 occurrences), relatively few populations or occurrences, recent and widespread declines, or other factors making it vulnerable to extirpation.

Source: TVA's Heritage Database "Botany within 5 Miles"

During field surveys, one species of state listed tree was observed in the forested area along the fence line of the southeast border of the Survey Area: bur oak (*Quercus macrocarpa*) (state ranked S2). Bur oak observations were limited to two individuals in lower elevations along a stream in the same general area. The two individuals were mature trees and had a diameter at breast height (dbh) of three to four feet. Representative photos are provided in the *Protected Species Habitat Assessment Report* (Appendix D).

Two areas of open chalk outcrops and associated buffers were also mapped in the southwest portion of the Survey Areas as areas that could potentially support sensitive plant species and will be recommended as avoidance areas. No rare plants were observed at or near the chalk outcrops during surveys.

## Rare Aquatic Species

With the exception of the white heelsplitter (S3), all aquatic species identified in the TVA Natural Heritage Database (listed below in Table 3-6) as potentially occurring within 10-miles of the Survey Area require aquatic habitats primarily comprised of medium to large rivers with moderate to fast moving currents.

Table 3-6: TVA Natural Heritage Database – Aquatic Species Potentially Occurring within 10-miles of Survey Area.

Common Name	Scientific Name	Rank/Status	Preferred Habitat Description
Blue Sucker	Cycleptus elongatus	S3	Large rivers with deep channels and swift currents that have substrates consisting of sand, gravel or rubble bottoms. (MDNR 2021a)
Crystal Darter	Crystallaria asprella	S1/LE	Medium to large rivers, usually with moderate to swift currents and clean sandy or gravel bottoms (MDNR 2021b)
Cylinder Elimia	Elimia cylindracea	S2	Riffle habitats often in areas with rocky legdes (NatureServe 2021)
Flat Pigtoe	Pleurobema marshalli	SX/LE	Medium to large rivers with moderate to fast current and sandy gravel to gravel cobble substrates and clean water (NatureServe 2021b)

Common Name	Scientific Name	Rank/Status	Preferred Habitat Description
Frecklebelly Madtom	Noturus munitus	S2/LE	Fast moving streams associated with larger rivers that provide firm gravel substrates with pebbles and rock (USFWS 2020).
Freckled Darter	Percina lenticula	S2	Medium to large rivers with deep and swift currents (NatureServe 2021c)
Heavy Pigtoe	Pleurobema taitianum	SX/LE	Riffles and shoals of small to large rivers with sandy gravel to gravel-cobble substrates and moderate to fast currents (NatureServe 2021d).
Southern Clubshell	Pleurobema decisum	S1	Highly oxygenated small to medium rivers with loose sand and gravel substrates (NatureServe 2021e).
Southern Combshell	Epioblasma penita	S1/LE	Medium sized rivers with high gradient and riffles with sandy gravel to gravel cobble substrates in moderate to swift currents (NatureServe 2021f).
Stirrupshell	Quadrula stapes	SX/LE	Medium to large rivers with moderate to fast current and sandy gravel to gravel cobble substrates and clean water (NatureServe 2021g)
Striped Bass	Morone saxatilis	SH	Coastal estuaries; migrates upstream to medium to large rivers during spawning typically within 6 km of the shore (NatureServe 2021h).
White Heelsplitter	Lasmigona complanata	S3	Rivers and stream of various sizes with substrates of gravel, sand, or mud (NatureServe 2021i).

S1 = Critically Imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation.

The six perennial streams identified within the Survey Area had limited flow and fragmented sections of water throughout the reaches. Perennial streams within the Survey Area were characterized by the presence of flow at the time of the site visit as well as secondary characteristics such as observance of fish and rooted aquatic fauna, indicating that water flows year-round. Perennial streams were approximately 6 to 25 feet in width at the OHWM with bank heights ranging from 3 to 25 feet. At the time of the delineation, the depth of water observed was

S2 = Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

S3 = Vulnerable in Mississippi due to a restricted range (on the order of 21 to 100 occurrences), relatively few populations or occurrences, recent and widespread declines, or other factors making it vulnerable to extirpation.

SH = Possibly Extirpated – Known from only historical records in Mississippi, but still some home of rediscovery. There is evidence that the species or ecosystem may no longer be present in the jurisdiction, but not enough to state this with certainty.

SX = Presumed Extirpated – Species or ecosystem is believed to be extirpated from Mississippi. Not located despite intensive searches of historical sites and other appropriate habitat, and virtually no likelihood that it will be rediscovered.

0.5 to 2.5 feet. Perennial streams flowed through upland fields, agricultural fields, and wooded riparian areas.

Observations of gambusia were documented during field surveys. However, no fish or macroinvertebrate sampling was conducted. No other observations of aquatic species were documented. Representative photographs of each stream flow regime type are provided in Appendix D.

Perennial streams within the Survey Area do not meet habitat requirements for these species. White heelsplitters can be found in a variety of freshwater settings and perennial stream within the Survey Area may provide marginal habitat for this species. Additionally, proposed solar array layouts would be designed to avoid perennial stream and their buffers. Therefore, impacts to aquatic species within the Survey Area are not expected.

# 3.4.2 Environmental Consequences

#### 3.4.2.1 No Action Alternative

Under the No Action Alternative, there would be no Project impacts to biological resources at the Project Site. It is assumed that existing agricultural operations would continue, and the undeveloped land would remain as is. There would be no Project-related impacts to existing vegetation at the Project Site. The forested areas within the Project Site could eventually be cleared for other development or landowner preference. Additional minor impacts to vegetation and wildlife could occur through continued applications of pesticides, herbicides, and fertilizers. No Project-related impacts would occur.

### 3.4.2.2 Proposed Action

## **Natural Areas**

The Proposed Action is not anticipated to have any impacts on the biological resources associated with natural areas, given the nature of the activities and distance from the Project Site to the natural areas.

### **Vegetation**

Under the Proposed Action the Solar Facility would be constructed and put into service, which would lead to direct adverse impacts to vegetation. Up to 270 acres of forested upland areas could be removed during initial construction clearing and grading activities to increase available land used for arrays and to reduce shading to the PV arrays.

# Rare & Protected Plant Species

Most of the forested areas that provide habitat for protected species or species that are biologically sensitive are present within avoidance buffers established for riparian areas along jurisdictional water features. Other areas have been excluded from the Project Site during planning and design stages because of topographic and constructability concerns. Avoidance areas will be established onsite with signage or temporary construction fencing as well as

identified on constraints/avoidance plans. Further, these areas will be included on engineering/constraints drawings provided to the construction contractor.

No observations of federal noxious weeds were documented within the Project Site. Upon completion of construction, the site would be planted or seeded with non-invasive vegetation, including native and naturalized plant species to encourage beneficial habitat, reduce erosion, and limit the spread of invasive species as part of the revegetation strategy, in accordance with EO 13112. Seed selection for the re-vegetation strategy would be developed to plant lower growing species in effort to reduce the amount of maintenance required below the PV arrays. If the construction of the Project increases growth of invasive species, these effects would be reduced once the re-vegetation process is initiated, and native and non-invasive species are established. MS Solar 6 would spot treat areas where invasive species/weeds begin growing until the native and non-invasive species are established throughout the site. MS Solar 6 would not implement an invasive species control plan or cogon grass plan within areas that were preestablished as avoidance areas during construction and operation as the intent of the avoidance area is to limit access to these areas by workers and equipment. Implementation of the Proposed Action would not significantly contribute to the introduction or spread of invasive species.

Of the approximately 270 acres of forested uplands within the Project Site, the majority of forested habitat could be developed for the Solar Facility and may be cleared to reduce shading for the PV arrays. Riparian areas would remain intact at and around jurisdictional streams and wetlands and avoided during construction to the greatest extent practicable. Tree removal in general would be minimized to the greatest extent practicable.

A majority of the Project Site is either disturbed, maintained, or actively cultivated cropland. There is potential to remove a minor amount of forested area within the Project Site (<18 percent) during clearing and grading activities. Additionally, the surrounding areas consist of similar vegetation communities and have also been mostly converted to cropland. Adverse impacts associated with vegetation removal resulting from implementation of the Proposed Action would be minor but long-term.

# Wildlife

Under the Proposed Action, the Solar Facility would be constructed, and negligible direct and indirect impacts could occur to wildlife. Most of the area proposed to be cleared is land currently used for agriculture and/or livestock grazing. Clearing of 270 acres out of approximately 480 acres total of forested habitats (woody wetlands and upland forests) could have moderate direct and indirect effects on wildlife utilizing those habitats. Impacts resulting from construction and operation of the Solar Facility would be temporary due to displacement of native species during construction. Direct effects on less mobile species or life stages of wildlife occurring at the site (i.e., nestlings, eggs, larvae, and burrowed animals) could occur during initial land clearing and grading activities. As mentioned above, forested riparian areas would be avoided to the greatest extent possible. Wildlife that are present are expected to disperse to suitable habitat nearby when construction activities commence. Wildlife species able to use early successional habitats would be able to return to the Project Site once construction activities are complete.

The habitat present is similar to the landscape and land use in surrounding areas. It is expected that wildlife in the Project Site would be able to relocate successfully to nearby areas of similar habitat and food resources during construction. Because the Project Site would be revegetated with native and/or naturalized non-invasive herbaceous plants maintained without the extensive use of harmful herbicides and pesticides, the site would provide a small amount of potentially suitable nesting and long-term foraging habitat for some species of songbirds and small mammals. Therefore, short-term direct and indirect impacts to wildlife during construction of the Proposed Action would be moderate. Minor long-term benefits would be realized for some species such as insects, small mammals, and reptiles and amphibians as frequently disturbed crop fields are established with native and non-invasive vegetation after construction.

# **Mississippi State-Listed Species**

Suitable habitat for three state-listed animals was identified within the Project Site: the eastern spotted skunk, painted bunting, and scissor-tail flycatcher. If utilizing areas for nesting within the Project Site, minor, short-term, direct impacts to painted bunting and scissor-tail flycatcher may occur if vegetation removal is implemented during their breeding season. If the eastern spotted skunk is present within the Project Site, it would likely relocate to nearby areas with similar habitat during construction. However, minor, short-term, direct impacts to younger individuals or individuals with slower mobility may occur during construction. Additionally, temporary impacts from construction activities may discourage these species from entering the area; however, they are expected to return to areas of suitable habitat within the Project Site once the project is operational and native grasses have been established. In areas where forested habitat is permanently converted to open space where PV arrays are located, forest-dwelling species would likely relocate to woody areas at the southern portion of the Project Site.

The Project was designed to minimize impacts on natural vegetation communities to the extent practicable. Specifically, most of the chalk prairie complex has been designated as a "no build area" for several reasons, including its potential as habitat for protected species. The chalk outcroppings at the Project Site would be completely avoided. Additionally, where construction impact areas overlapped with areas identified as potentially suitable habitat for rare and protected plants, the solar array layout would be designed to avoid impacts. Two individual bur oak trees were the only listed plant species observed during surveys which were within areas that would be excluded from development. Therefore, many rare plant or insect species that may occur within those habitats would remain unimpacted by the Project. TVA's Natural Heritage Program identified 28 plant species, 12 aquatic species, and 5 animal species within a 10-mile radius, 5-mile radius, and within the Project Site, respectively. However, MS Solar 6 has minimized development activities within areas identified as suitable habitat for rare species. Additionally, a 100-foot-wide buffer has been established around these areas to further reduce potential adverse impacts.

### **Federally Listed Species**

The wood stork is federally protected under the ESA. Wood storks typically develop roosting sites in medium to tall trees that occur in stands either in swamps or on islands surrounded by relatively

broad expanses of open water; primarily bald cypress (*Taxodium distichum*) swamps. Nesting usually occurs in bald cypress trees in swamps. Wetlands within the Survey Area were small, isolated, and low quality with little to no surface water (Appendix D). Therefore, no suitable roosting habitat for wood storks was observed within the Survey Area; however, marginal foraging habitat for the species was observed within most of the small and confined ponds throughout the Survey Area. As the current layout shows, the area of impact has been designed to avoid any impacts on wetlands. There are also large aquaculture/fish farms both north and south of the Project that may attract foraging wood storks. The Project would not affect fish farms or large open waters outside the immediate Project limits. For these reasons, impacts to wood storks or wood stork habitat are not anticipated from implementation of the Proposed Action.

Only four populations of Price's potato-bean are known to exist in Mississippi today because of its specific habitat requirements (USFWS 2016). Areas within the Project Site that could potentially support Price's potato bean will be avoided. During the survey, no individuals were observed. Additionally, the small chalk prairie complex has been designated as an "avoidance area" for several reasons, including its potential as habitat for protected species. For these reasons, the Proposed Action would have no effect to Price's potato-bean.

Suitable foraging habitat for the NLEB was identified over ponds, wetlands, and streams located in the Project Site [Appendix D (Appendix D – Figure A-1)]. Approximately 328 acres of suitable summer roost habitat is also present in the forested areas. However, the results of the acoustic bat surveys indicated that there are no NLEBs within the Project Area.

Informal consultation with the USFWS, under Section 7 of the Endangered Species Act, was conducted in March 2022. The USFWS acknowledged TVA's determination that there would be "no effect" on the wood stork and Price's potato bean. They also concurred with TVA's "may affect, but not likely to adversely affect" determination for the NLEB. Copies of consultation letters are included in Appendix E.

## **Bald Eagles and Migratory Birds**

Sparse habitat along the forest edges and vegetated stream corridors may provide habitat for migratory birds within the Project Site. Wetlands, streams, and forested areas would be avoided to the greatest extent practicable under the Proposed Action. Up to 270 acres of tree removal could be conducted during construction. Implementation of the Proposed Action may reduce negligible amounts of nesting habitat, reduce minor amounts of foraging habitat, and may displace birds to surrounding areas with similar habitat and land use during construction. Most riparian areas will be avoided, however, minor direct impacts to rusty blackbirds may occur if pond edges and riparian woodlands are disturbed. Mortality to rusty blackbirds from vegetation removal could happen if tree clearing is conducted during its breeding season, however, it would most likely cause birds to flush and have to relocate to similar habitat nearby. Implementation of the Proposed Action may also reduce minor foraging habitat for species such as the American kestrel and other birds of prey that use farm fields as foraging habitat. However, it is assumed any birds using the farms fields within the Project Site for foraging would use adjacent lands if there were a significant reduction in prey items. Red headed woodpeckers may be present in forested areas proposed for

clearing. Tree clearing could occur in May, June, July, August, and September which is the bird's breeding season. The most likely impact would be limited to red-headed woodpeckers relocating to nearby areas of similar habitat during active construction. In smaller numbers, inadvertent instances of bird mortality could take place if red headed woodpecker nests occur within trees that would be cleared during its breeding season. Impacts would be direct and adverse, but minor in these instances. Long-term, the Project Site would provide nesting and foraging habitat for some species of songbirds able to use early successional habitats once operational.

Due to limited nesting and foraging habitat for migratory birds currently at the Project Site, no significant direct or indirect adverse impacts on migratory birds or bald eagles are expected from Project activities.

### 3.5 Visual Resources

Visual resources are described as physical features or visual characteristics of a place that define the visual and aesthetic character of an area. The following sections describe the aesthetic and visual characteristics of the Project Site and surrounding areas.

#### 3.5.1 Affected Environment

Visual resources are the visible features of an area and can include both natural viewsheds and viewsheds comprised of manmade attributes. Visual resources can influence how an observer experiences a certain location and how they may distinguish the location from other locations. Visual resources are important to the people living in and traveling through an area. Additionally, visual resources can make up a significant component of historically and/or culturally significant settings. Additional information pertaining to the visual area of potential effects (APE) and historic or culturally significant resources is provided in section 3.8.1.

During preliminary field surveys, Project historians conducted a viewshed analysis to document the actual visibility of the Project within an established Visual APE that accounts for terrain, vegetation, and other setting intrusions that would impede views of the Project. The Project Site constitutes the established APE for physical impacts while the half-mile buffer around the Physical APE was developed to account for visual and other non-physical effects from Project implementation. The Visual APE is comprised of approximately 1,524 acres (Project Site) plus an additional half-mile buffer around the outside of the Project Site.

All but the forested areas of the Project Site are comprised of active agricultural land and pastureland for livestock grazing. The active agriculture area is used to produce a rotational mix of corn and soybeans. There are several mature stands of hardwood forests around the Project Site, as visible in Figure 2-1. The Golden Triangle I Solar and BESS Project is located just north of GT2. The Golden Triangle Regional Airport (GTR) is located approximately three miles northeast of the Project Site. GTR is a commercial airport but also is used by private airlines due to its central location. Small jets land and take off from this airport regularly (an average of approximately 1-2 airplanes per hour).

The photos below provide representative views of the Project Site. Generally, the Project Site is rural and agricultural with isolated single-family homes adjacent to the Project Site. There are three single home residences located within less than half-mile from the Project site. The topography is characterized by generally flat terrain to gently rolling hills interspersed with intermittent and perennial streams. Scenic attractiveness is separated into three classes: Class A – Distinctive; Class B – Typical, and Class C – Indistinctive. The scenic attractiveness of the Project Site is rated as typical, which means the visual aesthetics are considered common of a rural-agricultural and sparsely residential area. The Scenic integrity is assessed as moderate due to the surrounding railroad operations and commercial airport.



Figure 3-8: General Site Photo of GT2 Project Site



Figure 3-9: General Site Photo of GT2 Project Site

The Project is located within the Mississippi Hills National Heritage Area (MHNHA), which was designated by Congress and the President in April 2009, through the Omnibus Public Land Management Act of 2009. The MHNHA fully covers 19 counties and portions of 11 counties in the northeastern part of Mississippi. The MHNHA is representative of a distinct cultural landscape that has been shaped predominantly by the intersection of the Appalachian and Delta cultures. National Heritage Areas are places where cultural, historic, and natural resources unite as a significant national landscape. NHAs are a partnership unit of the National Park Service (NPS), but unlike the national parks managed by the NPS, NHAs are large, lived-in landscapes. The primary theme of the MHNHA is African-American heritage, Native American history, Civil War, and music and literature (Mississippi Hills National Heritage Area, 2021). The MHNHA includes the Elvis Presley Birthplace and Museum, the Natchez Trace Parkway, the homes of William Faulkner and Tennessee Williams, the Corinth Civil War Interpretive Center, and Brices Cross Roads National Battlefield. No associated MHNHA markers or Civil War battle sites were identified within the Physical APE or Study Area (Shaver, Harris, House, and Kepka 2021).

# 3.5.2 Environmental Consequences

#### 3.5.2.1 No Action Alternative

Under the No Action Alternative, no Project-related impacts to visual resources would occur. Existing views would be expected to remain unchanged from the present conditions, which are either natural and undeveloped or open fields under agricultural use.

### 3.5.2.2 Proposed Action

The Project would convert mostly agricultural land and pastureland into industrial/electrical infrastructure consisting mostly of low-profile PV arrays. Figure 2-2 provides the preliminary layout for PV arrays.

During site visits, which occurred between June to September 2021, the Project Area was assessed for the potential for visual impacts on sensitive areas such as residences, churches, schools, and from roadways. There are two minimally traveled gravel public roadways that either abut or are in close proximity to portions of the Project Site:

- Gilmer Wilburn Road: runs east to west along northern project boundary
- Hardy Billups Road: runs north to south approximately 2,000 feet east of GT2.

From the above listed vantage points, the manufactured and structural appearance of the Solar Facility would be most obvious in the morning, when the dark-colored solar panels would be upright, averaging approximately eight feet from the ground, at full tilt, facing east. At midday, the panels would be less obvious as they are laying nearly flat. At midday, when the panels are flat, they would average about five feet off the ground. In the evening, when the panels would be at full tilt and facing west, they would be most obvious along Gilmer Wilburn Road.



Figure 3-10: Representative Photo of Single-Axis Tracking PV System in the Moming, Facing East



Figure 3-11: Representative Photo of Single-Axis Tracking PV System at Midday, Laying Flat

Although portions of the Project would be visible across open fields or otherwise clear areas, the residential properties and roadways in the vicinity of the Project Site generally have mature trees along or near property boundaries and fence lines that would partially or almost fully obscure views of the Solar Facility from many vantage points. The relatively stable elevations and the maintenance of existing vegetation along the perimeter of the Project Site would largely shield views from most vantage points to the Solar Facility.

There are several single-family homes located within close proximity to the Project Site. One home is located approximately 2,200 feet west of GT2 on the south side of Gilmer Wilburn Road. A second home is on a parcel located in between two GT2 parcels. At its closest point, the home is 525 feet from the limits of GT2; however, the home is surrounded by mature trees that will likely obstruct the resident's view of the solar arrays. A third home is located along the north side of Gilmer Wilburn Road, just north of GT2 and east of the aquaculture facility. It is only about 275 feet from the Project Site; however, there is a dense stand of mature trees between the home and GT2. It is unlikely that this resident will be able to view the solar arrays from its property.

Single family homes adjacent to the Project site will likely be able to view the solar arrays when residents are coming or going, but there are vegetated fencerows with mature trees along the parcel boundaries which would likely provide natural screening from any arrays that could be positioned in this area when residents are at home. With the natural vegetation screening and vegetated fencerows, no long-term adverse direct impacts on single-family home areas would occur from the Project.

Construction of the proposed Project would temporarily alter the visual character of the Project Site. During construction, heavy machinery would be present, changing the visual characteristics from vantage points surrounding the Project Site. Within the Project Site, many trees and other tall vegetation would be preserved to the extent practicable, which would provide natural buffers from public roadways. In areas where grading would be necessary, minor changes to the contour, color, and texture of the scenery would be visible. The Project Site would appear as a mixture of

neutral colors such as dark browns and grays due to earthmoving, road construction, and concrete activities. Water would be used to suppress dust. Visual impacts from construction would be minimal at night since most construction is anticipated to occur during the day. Erosion control silt fencing and sediment traps would be removed once construction is complete, and bare areas would be promptly vegetated.

Over the approximately 17-month-long construction phase, indirect impacts to visual resources in the area may occur due to increased traffic and movement of heavy machinery to the Project Site along local roads. Overall, there would be minor temporary direct and indirect impacts to visual resources during the construction phase of the Proposed Action.

Visual impacts during the operation phase of the Project would be minor in the immediate vicinity, due to substantial tree buffers around property boundaries and rolling hills within the Project Site. Visual impacts would be minimal to negligible on a larger scale, due to variation of the visual attributes of the surrounding area as distance from the Project Site increases.

### 3.6 Noise

The magnitude and frequency of noise may vary considerably over the course of the day, throughout the week, and across seasons, in part due to changing weather conditions and the effects of seasonal vegetation cover. Two measures that relate the time-varying quality of environmental noise to its known effect to people are the 24-hour equivalent sound level (Leq) and day-night sound level (Ldn). The Leq is the level of steady sound with the same total (equivalent) energy as the time-varying sound of interest, averaged over a 24-hour period. The Ldn is the Leq plus 10 dBA added to account for people's greater sensitivity to nighttime sound levels (typically considered between the hours of 10:00 p.m. and 7:00 a.m.). The A-weighted scale is used to assess noise impacts because human hearing is less sensitive to low and high frequencies than mid-range frequencies. The human ear's threshold of perception for noise change is considered 3 A-weighted decibels (dBA); 6 dBA is clearly noticeable to the human ear, and 10 dBA is perceived as a doubling of noise (or halving, if the noise is decreasing).

In 1974, the EPA published *Information on Levels of Environmental Noise Requisite to Protect Public Health and Welfare with an Adequate Margin on Safety*, which evaluated the effects of environmental noise on public health and welfare (USEPA 1974). In this document, the EPA indicated that a L<sub>dn</sub> of 55 dBA is the noise threshold that would prevent outdoor activity interference or annoyance from continuous noise. We have adopted this criterion and use it to evaluate potential noise impacts from TVA projects at noise sensitive areas (NSAs) such as residences, schools, or hospitals. A L<sub>dn</sub> of 55 dBA is equivalent to a continuous sound level of 48.6 dBA for facilities that generate constant sound levels. A list of typical sound levels for common sound sources is presented in the table below.

Table 3-7: Typical Sound Pressure Levels Associated with Common Sound Sources

Sound Pressure Level	Subjective	Environ	ment		
(dBA)	Subjective Evaluation	Outdoor	Indoor		
140	Deafening	Jet aircraft at 75 feet			
130	Threshold of pain	Jet aircraft during takeoff at a distance of 300 feet			
120	Threshold of feeling	Elevated train	Hard rock band		
110		Jet flyover at 1,000 feet	Inside propeller plane		
100	Very loud	Power mower, motorcycle at 25 feet, auto horn at 10 feet, crowd sound at football game	-		
90		Propeller plane flyover at 1,000 feet, noisy urban street	Full symphony or band, food blender, noisy factory		
80	Moderately loud	Diesel truck (40 miles per hour) at 50 feet	Inside auto at high speed, garbage disposal, dishwasher		
70	Loud	B-757 cabin during flight	Close conversation, vacuum cleaner		
60	Moderate	Air-conditioner condenser at 15 feet, near highway traffic	General office		
50	Quiet		Private office		
40		Farm field with light breeze, birdcalls	Soft stereo music in residence		
30	Very quiet	Quiet residential neighborhood	Inside average residence (without TV and stereo)		
20		Rustling leaves	Quiet theater, whisper		
10	Just audible		Human breathing		
0	Threshold of hearing				
Sources: Egan, 1988; Ramsey and Sleeper, 1994					

# 3.6.1 Affected Environment

The proposed Solar Facility is located in a rural area adjacent to local roadways, a rail line to the west, and the Golden Triangle Regional Airport is about three miles to the northeast. The major noise sources in this area are vehicles traveling along the gravel roadways, trains running on the railroad, private and commercial jets, farm equipment, wind, and farm animals. Noise levels in rural areas typically range from 45 to 55 dBA. A day-night average sound level of 55 dBA is commonly used as a threshold for noise levels which could result in adverse impacts. Prolonged exposure to levels above 65 dBA is considered unsuitable for residential areas.

There are several single-family homes located within close proximity to the Project Site. One home is located approximately 2,200 feet west of GT2 on the south side of Gilmer Wilburn Road. A second home is on a parcel located in between two GT2 parcels. At its closest point, the home

is 525 feet from the limits of GT2. A third home is located along the north side of Gilmer Wilburn Road, just north of GT2 and east of the aquaculture facility. It is about 275 feet from the Project Site boundaries.

# 3.6.2 Environmental Consequences - Noise

#### 3.6.2.1 No Action Alternative

Under the No Action Alternative, no Project-related noise impacts would occur. Current noise impacts related to air traffic, train operations, and agricultural land use, which is likely minimal, would persist.

## 3.6.2.2 Proposed Action

Under the Proposed Action Alternative, noise levels would be temporarily elevated in the areas immediately surrounding the Project Site when construction is occurring. Construction activities such as tree removal, site grading, and installation of the PV panel support posts would generate noise on a temporary basis. Maximum noise levels produced by the construction equipment are in the range of 80 to 85 dBA at a distance of 50 feet. The nearest occupied residence is approximately 275 feet from the Project Site. Nearby residents would likely experience elevated noise levels caused by the operation of construction equipment during daytime hours (typically 8am – 7pm) as well as an increase in traffic during peak morning and evening commutes. However, the elevated noise levels would be short in duration and would not occur at night. Furthermore, the elevated noise levels at the closest residence would not exceed the 65 dBA for prolonged periods of time.

As the Project design evolves, MS Solar 6 would locate inverters at least 500 feet away from residences. If a 500-foot setback could not be accomplished for the nearest residence, once the Solar Facility is operational, mufflers or sound reducing devices may be installed on inverters that are located within 500 feet of the single-family home if an increase in noise from the Solar Facility occurs. Periodic maintenance activities, such as mowing, would temporarily increase noise levels; however, these levels would be consistent with the existing agricultural operations that have historically occurred in the area. Overall, the noise impacts resulting from both construction and operation of the Project would be negligible.

# 3.7 Air Quality and Greenhouse Gas Emissions

This section describes existing air quality and GHG emissions in the Project region and the potential impacts on air quality and GHG emissions that would be associated with the No Action and Proposed Action Alternatives.

#### 3.7.1 Affected Environment

Ambient air quality is determined by the type and concentration of pollutants emitted into the atmosphere, the size and topography of the air shed in question, and the prevailing meteorological conditions in that air shed. With the issuance of the Clean Air Act of 1970 and its amendments, Congress mandated the protection and enhancement of our nation's air quality. The EPA

established the National Ambient Air Quality Standards (NAAQS) for the following primary criteria pollutants to protect public health and welfare: sulfur dioxide (SO<sub>2</sub>), ozone (O<sub>3</sub>), nitrogen dioxide (NO<sup>2</sup>), particulate matter less than or equal to 10 micrometers (PM<sub>10</sub>), particulate matter less than or equal to 2.5 micrometers (PM<sub>2.5</sub>), carbon monoxide (CO), and lead (Pb).

Primary NAAQS were developed to protect public health and safety, and secondary NAAQS were developed to protect public welfare (e.g., visibility, crops, forests, soils, and materials) from known or potential adverse effects of air pollutants. Areas in compliance with the NAAQS are designated "attainment" areas. Areas not in compliance with the NAAQS are designated as "nonattainment" areas. New sources proposed in or near nonattainment areas may be subject to more stringent air permitting requirements. Nonattainment areas are usually defined by county. National standards may not be exceeded more than once per year (except where noted). Areas that cannot be classified based on available information for a specific pollutant are designated as "unclassifiable" and are treated as attainment areas unless proven otherwise. If an area that was formerly designated as nonattainment for a particular pollutant later qualifies as attainment, that area is then categorized as "maintenance" for that pollutant for the next 20 years as long as the area continues to meet the NAAQS for that pollutant. If an area remains in attainment for the 20-year maintenance period, the status will qualify it to be formally designated back to normal attainment.

# 3.7.1.1 Regional Air Quality

The area where the Project is located is in rural Lowndes County, which has little development in the vicinity apart from that related to rural-residential and agricultural uses. Denser development is approximately 10 miles or more to the west in Starkville and about 12 miles to the east in Columbus. Lowndes County is considered in attainment for all pollutants (USEPA 2021a).

Table 3-8 presents the most recent (2017) EPA emission inventory data (USEPA 2021d) for the most prevalent NAAQS pollutants for Lowndes County. These data represent anthropogenic emissions from all stationary source and mobile source activities. Included in Lowndes County is the City of Columbus, the Columbus Airforce Base, and the Golden Triangle Regional Airport, which offers private and commercial flights. All values fall beneath the USEPA thresholds for NAAQS pollutants.

Table 3-8: Average Emissions of NAAQS Pollutants in Lowndes County

Pollutant	Emissions for Lowndes County (tons per year)
Carbon Monoxide	5,168
Nitrogen Oxides (NO <sub>x</sub> )	2,992
PM <sub>10</sub>	7,393
PM <sub>2.5</sub>	1,300
Sulfur Dioxide (SO <sub>2</sub> )	452
Volatile Organic Compounds (VOCs)	11,491

Source: EPA 2021d

### 3.7.1.2 Regional Climate

The Project Site is located almost halfway between Starkville and Columbus, MS. This region of the country is known as the humid subtropical climate region and is characterized by temperate winters; long, hot summers; and rainfall that is evenly distributed throughout the year. Normal annual precipitation at the Project Site is 55 inches per year. Typical weather conditions in Artesia range from 91° to 72° in the summer months and 54° to 36° in the winter months with mild Spring and Fall temperatures.

The Antecedent Precipitation Tool (APT), developed by the USACE, was used to assess the climate conditions in the months leading up to and during the March 2021 environmental survey. The APT provides a standardized method for evaluating precipitation conditions relative to a climate normal, the presence of drought conditions, and the approximate dates of wet and dry seasons for a given location.

At the time of the March 2021 survey, conditions on site appeared to be dry, with little to no rainfall immediately before field mobilization. The APT results confirm that conditions on site were normal, and the overall drought index (PDSI) was listed as mild wetness. Rainfall data reported from the APT is available as an Appendix to the Wetland and Waterbody Delineation Report (Appendix B).

#### 3.7.1.3 Greenhouse Gas Emissions

GHGs typically consist of natural and man-made compounds that are released into the earth's atmosphere. GHGs also absorb a portion of Earth's infrared radiation and can re-emit some of the radiation back to the earth's surface. When radiation is emitted back to the earth's surface, temperatures are typically warmer than they would naturally be. With that process, GHGs act as insulation and contribute to the maintenance of global temperatures. Increasing levels of GHGs in the atmosphere result in an increase in temperature on earth, commonly known as global warming. Changes in climate associated with global warming produce adverse economic and social consequences globally through changes in weather (e.g., more intense natural disasters, greater risk for forest fires, flooding) (USGCRP 2020). However, as shown in Figure 3-12, for the State of Mississippi, there is currently no noticeable long-term upward trend in temperature.




Figure 3-12: Observed and Projected Temperature Change in Mississippi

Source: NOAA 2019

The primary GHG emitted by human activities in the U.S. is carbon dioxide (other than water vapor), representing approximately 82 percent of total GHG emissions in the US (USEPA 2021e). Dominant sources of carbon dioxide and of overall GHG emissions is fossil fuel combustion. Emissions of the GHG methane, which have declined in the U.S. from levels in 1990, are primarily a result of digestion of domestic livestock, decomposition materials in landfills, coal mining, and natural gas leaks. Agricultural soil management is the major source of the GHG nitrous oxide emissions in the United States, representing approximately 74 percent of its emissions from human activities (USEPA 2021e).

## 3.7.2 Environmental Consequences

#### 3.7.2.1 No Action Alternative

Under the No Action Alternative, no air emissions or GHGs would be generated by equipment or vehicles from construction or operation of the Solar Facility. Existing land use would be expected to remain as a mix of farmland and undeveloped land, and the only ongoing emissions would be due to vehicles or equipment used to operate the farms or maintain the agricultural clearings, and from domestic livestock operations.

### 3.7.2.2 Proposed Action Alternative

Under the Proposed Action, minor direct impacts on air quality would be anticipated as a result of construction and operation of the Project. Temporary impacts to GHG emissions expected during construction would be negligible.

# Regional Air Quality

The majority of potential air quality impacts associated with the Proposed Action would occur during construction. Construction activities would result in emissions from the operation of construction equipment, contracted employees' personal vehicles, and fugitive dust suspension from clearing, grading, and other activities across exposed dry soil. Tree debris from clearing would be removed by either burning or chipping and grinding. As burning may occur, this could generate temporary localized air quality impacts due to smoke particles and gases. Any such burning of vegetative debris would be done in accordance with local ordinances or burn permits and is not expected to have any health consequences for this rural area.

The use of construction equipment would cause a minor temporary increase in GHG emissions during construction. Combustion of gasoline and diesel fuels by internal combustion engines (haul trucks and off-road vehicles) would generate local emissions of PM, NO<sub>x</sub>, CO, VOCs, and SO<sub>2</sub>. The total amount of these emissions would be small and would result in negligible air quality impacts overall.

Approximately 95 percent (by weight) of fugitive emissions from vehicular traffic over paved and unpaved roads would be composed mainly of particles that would be deposited near the roadways on routes taken to reach the Project Site. As necessary, fugitive dust emissions from construction areas and paved and unpaved roads would be mitigated using wet suppression. Wet suppression would reduce fugitive dust emissions from roadways and unpaved areas by as much as 95 percent. Direct impacts to air quality associated with construction activities would be temporary and minor.

## Regional Climate

No noticeable direct or indirect impacts to the regional climate would occur as a result of the proposed Project. Local or regional climate effects can occur, for example, with major changes in land use that affect the hydrological cycle, or that create large impervious surfaces, thus changing the radiative heat balance over a large area. The Project would change the surface characteristics somewhat, but it would have little effect on soil permeability and hydrologic characteristics of the developed area. Vegetation would still grow under and around the solar panels, tending to maintain a landscape with significant evapotranspiration of precipitation, as opposed to creating significant runoff of precipitation which happens with urban development. Therefore, average temperatures of the developed area are not expected to change significantly due to the proposed development.

## Greenhouse Gases

The use of construction equipment would cause a minor temporary increase in GHG emissions during construction activities. Combustion of gasoline and diesel fuels by internal combustion

engines (trucks and off-road vehicles) at the site would generate emissions of CO<sub>2</sub> and very small amounts of other GHGs such as methane and nitrous oxide. Additional GHG emissions would occur due to transporting materials and workers to the Project Site, and GHGs would be emitted in the US or globally for production and transportation of the materials used for construction. The production of construction materials is expected to represent the largest portion of the Project-related GHG emissions. The total GHG emissions resulting from construction should eventually be offset by Project operation over the long term, assuming that the electricity generated by the Project would offset the need for some fossil-fuel-based electricity generation and the GHG emissions associated with the combustion of fossil fuels.

Tree and other tall vegetation removal during construction of the Project would represent a minor loss of sequestered carbon, as well as potential future carbon sequestration. Trees and other tall vegetation currently remove CO<sub>2</sub> from the air and sequester it as biomass. The loss of this carbon sink would constitute a minor adverse direct and indirect impact as sequestration would have continued for the life of the vegetation and long into the future, assuming that other future changes on the Project Site would not result in deforestation. The loss of the carbon sink from tree removal would be at least partially offset by the increased sequestration of CO<sub>2</sub> by the permanent grass-dominated vegetation that would be maintained on the Project Site.

The operation of the Project is not anticipated to have any long-termadverse impacts to air quality or GHG emissions. Operation of the Solar Facility would not produce emissions. Minor emissions would occur during maintenance activities, including facility inspections and periodic mowing. Conversely, overall emissions of air pollutants from the TVA power system would decrease during operation as the nearly emissions-free power generated by the Solar Facility would offset the need for power that would otherwise be generated, at least in part, by the combustion of fossil fuels. The reduction in GHG emissions resulting from the operation of the Solar Facility would have little noticeable effect at regional or larger scales. It would, however, be a component of the larger ongoing system-wide reduction in GHG emissions by the TVA power system through reducing the need for some fossil-fuel based electricity generation. The adverse impacts of GHG emissions and the beneficial impacts of TVA's efforts to reduce GHG emissions are described in more detail in the TVA IRP (2019).

### 3.8 Cultural Resources

Cultural resources include, but are not limited to, prehistoric and historic archaeological sites, historic structures, and historic sites at which important events occurred. Cultural resources are finite, non-renewable, and often fragile. They are frequently threatened by industrial, commercial, and residential development, as well as construction of roads and other infrastructure.

The NHPA provides for a national program to support both public and private efforts to identify, evaluate, and protect the nation's important cultural resources. Once identified, these resources are evaluated for inclusion in the NRHP maintained by the NPS. Tangible cultural resources may qualify for inclusion in the NRHP if they are 50 years of age or older and if found to possess one or more of four different criteria, in accordance with 36 CFR § 60.4:

- Criterion A: association with events that have made a significant contribution to the broad
  patterns of our history. Such events may include a specific occurrence or pattern of
  occurrences, cultural traditions, or historic trends important at a local, regional, or national
  level. To be considered in association with a cultural resource, events must be important
  within the particular context being assessed.
- Criterion B: association with the lives of persons significant in our past. People considered
  may be important locally, regionally, or nationally, and the cultural resources considered
  are limited to properties illustrating a person's achievements rather than commemorating
  them.
- Criterion C: embodiment of the distinctive characteristics of a type, period, or method of
  construction; representative of the work of a master; possessing high artistic values; or
  representative of a significant and distinguishable entity whose components may lack
  individual distinction. Cultural resources considered generally include architectural
  resources such as buildings, objects, districts, and designed landscapes.
- Criterion D: cultural resources that have yielded, or may be likely to yield, information
  important in prehistory or history. Considered cultural resources typically include
  archaeological sites but may also include buildings, structures, and objects if they are the
  principal source of important information not contained elsewhere.

Under Section 106 of the NHPA, each federal agency must consider public views and concerns about historic preservation issues when making final project decisions (GSA, 2020). The NHPA addresses the preservation of "historic properties," which are defined under the Act as any prehistoric or historic district, site, building, structure, or object included in or eligible for inclusion in the NRHP. Under Section 106 of NHPA, the Project is required to consider ways to avoid or minimize effects from its undertakings on significant cultural resources.

## 3.8.1 Affected Environment

Phase I archaeological and historic architecture surveys were performed within the Project Site between June 6 through September 17, 2021. The cultural survey area, also referred to as the APE included more than 1,837 acres. Additionally, a half-mile buffer around the Physical APE was developed to account for visual and other non-physical effects from Project implementation (Visual APE). The Visual APE included the Project Site plus an additional half-acre buffer surrounding the Project Site. The Phase I archaeological resources survey of the Physical APE and the architectural resources survey of the Visual APE and the refined visual effects buffer were conducted between June and September 2021.

The purpose of the Phase I Survey was to identify and document both previously recorded and newly documented archaeological and historic-age non-archaeological resources, to assess their eligibility for listing in the NRHP, and to assess potential Project impacts to historic (NRHP-listed or eligible) properties as required under Section 106 of the NRHP. The surveys were conducted to professional standards and guidelines and in accordance with the *Mississippi State Historic* 

Preservation Office's (MSSHPO) Mississippi Department of Archives and History (MDAH), Mississippi Standards for Archaeological Practices (MDAH 2019), the Secretary of the Interior's Standards and Guidelines for Archaeology and Historic Preservation (48 FR 44716-44742), and in accordance with the Secretary's Standards for Identification (48 FR 44720-44723).

Prior to the cultural resource survey field effort, a desktop review was conducted of the Study Area comprising the Physical APE and a half-mile buffer to determine if any previously recorded archaeological sites, architectural resources, and NRHP-listed properties are located in the Study Area. The review involved the evaluation of previously recorded archaeological sites and cultural resources survey data maintained housed at the MDAH – Archaeological Search Room, within the Charlotte Capers Archives and History building in Jackson, Mississippi and the online MDAH Historic Resources Inventory Map Database.

The background research identified three previously recorded archaeological sites within the Study Area (Table 3-9). None of the previously recorded archaeological sites are located within the Physical APE. Two of the previously recorded archaeological sites, 22LO889 and 22LO1082, are unevaluated for listing in the NRHP. The third previously recorded archaeological site, 22LO1085, has been determined not eligible for listing in the NRHP.

Table 3-9: Previously Recorded Archaeological Sites within the Study Area

Site Number	Site Type	Site Component	Name	NRHP Eligibility	Date Recorded
22LO889	Artifact scatter	Mississippian	Tom Hardy #3	Unevaluated	09/24/1987
22LO1082	Artifact scatter	Historic-era	-	Unevaluated	07/13/2020
22LO1085	Artifact scatter/Concrete silo	Historic-era		Not eligible	07/11/2020

Source: Mississippi Department of Archives and History (MDAH), January 2020 and April 2021.

Three previous cultural resources surveys have been recorded within the 0.5-mile Study Area (Table 3-10). None of the previous cultural resource surveys cross the Project's Physical APE.

Table 3-10: Previous Archaeological Surveys within the Project Study Area

Report Title	Report Number	Report Author	Date
Cultural Resources Survey of Proposed Widening of U.S. Highway 45A from the Noxubee County Line to U.S. Highway 82 (MDOT Project No. 97-0079-01-018-10), Lowndes County, Mississippi.	95-221	Gray, Bruce J.	June 1995
Cultural Resources Investigations of the Proposed Nickoles Brothers Catfish Farm Tract, Lowndes County, Mississippi.	00-0312	Atkinson, Jerry R.	October 2000

Report Title	Report Number	Report Author	Date
Phase I Cultural Resources Survey for the Golden Triangle I Solar Project, Lowndes County, Mississippi.	-	Shaver, Douglas, et. al.	October 2021

Source: MDAH, January 2020 and April 2021.

TVA initiated consultations on a government-to-government basis with the tribes listed below regarding the Proposed Action's potential to affect historic properties that are of religious and cultural significance to federally-recognized Indian tribes. A summary of TVA's consultations with federally-recognized Tribes is included in Appendix E. None of the consulted tribes identified such properties or objected to the Proposed Action.

- Absentee Shawnee Tribe of Indians of Oklahoma,
- Alabama-Coushatta Tribe of Texas,
- The Chickasaw Nation,
- The Choctaw Nation of Oklahoma,
- Coushatta Tribe of Louisiana,
- Eastern Shawnee Tribe of Oklahoma,
- Jena Band of Choctaw Indians,
- Kialegee Tribal Town,
- Mississippi Band of Choctaw Indians,
- The Muscogee (Creek) Nation,
- Shawnee Tribe, and
- Thlopthlocco Tribal Town

#### 3.8.1.1 Architectural Resources

Review of the MDAH Historic Resources Inventory Map, and previous survey efforts for the GT2 Project adjacent to Golden Triangle I, identified 8 individual resources on 5 parcels within the Non-Physical APE that were determined not eligible for NRHP inclusion (Resources 15, 41a, 41b, 42, 43, and 44a-44c).

The historic-age, non-archaeological resources survey resulted in the documentation of 25 historic-age resources on 7 properties. Of the accessible parcels, 2 properties include resources recommended for NRHP inclusion (Resources 02a and 05a). Two additional resources associated with Resource 05a are recommended as contributing (Resources 05c and 05d). Two resources, Resource 02h (Smith Oaks Plantation cemetery) and Resource 07 (a former African American cemetery) warrant additional NRHP eligibility analysis. The remaining resources are not recommended for NRHP inclusion due to a lack of historic associations, integrity, and/or significance. The NRHP eligible (and contributing) properties and the Smith Oaks Plantation cemetery and the former African American cemetery would not be impacted directly by the Project. Specifically, the resources are removed in distance and shielded by vegetation and other

setting intrusions. As a result, construction of the Project would not affect any of the characteristics that qualify the resources for NRHP inclusion.

The remaining documented resources include primarily mid- to late-twentieth-century Minimal Traditional and Ranch-style dwellings and outbuildings, including barns and sheds, as well as two additional family cemeteries. Other than the properties recommended for NRHP inclusion, none of the remaining resources recorded during the survey, either individually or collectively, appear to meet the criteria for NRHP inclusion due to a lack of significance and/or architectural integrity.

# 3.8.1.2 Archaeological Resources

During fieldwork for the Phase I archaeological resources survey of the Physical APE, 26 newly identified archaeological sites, of which 13 are isolated finds consisting of a single artifact or a site with three or less similar artifacts within a single shovel test, were recorded. Of the 13 isolated find sites, two are precontact Native American finds and eleven are historic-era. None of the isolated find sites are recommended eligible for inclusion in the NRHP. None of the isolated find sites are recommended for avoidance by Project implementation.

The other newly identified non-isolated find archaeological sites include 10 historic-era sites, one multicomponent site consisting of a historic era artifact scatter and a single Late Archaic Ledbetter or Flint Creek Pontchartrain Plano convex lithic knife, and two precontact Native American sites. One of the newly identified precontact Native American sites, site 22LO1100, has been identified as a Late Archaic through Late Woodland period precontact Native American occupation / camp site artifact scatter. Site 22LO1100 is recommended as unevaluated for inclusion in the NRHP and will be avoided by Project implementation. The second precontact Native American site consists of a single Late Archaic Smithsonia or Flint Creek type projectile point. Two newly identified sites, 22LO1114 and 22LO1098 are historic scatters with intact features. 22LO1114 (DS4) was identified through shovel tests excavations, with one shovel test uncovering a burn feature. 22LO1098 (CW1) is an occupation site with an intact cistern and small brick lined foundation. Both sites are associated with structures depicted on a historic 1911 Soils Map of Lowndes County. Site 22LO1098 is recommended as eligible for inclusion in the NRHP under Criterion A and D and will be avoided by Project implementation. An individual archaeological assessment for site 22LO1114 recommends it as unevaluated for inclusion in the NRHP and it will be avoided by Project implementation.

Site 22LO1117 is an extant portion of the Smith Oaks harness horse racing track. The 0.5-mile dirt track was constructed by Thomas Wilburn II in the late 1940s or early 1950s. Wilburn was a professional Standardbred horse trainer and harness horse racing champion who was inducted in the Illinois Harness Horse Racing Hall of Fame, was the 1991 Mississippi Cattleman of the Year, and is a direct descendant of John McLaughlin Smith, the original plantation owner. Site 22LO1117 is recommended for inclusion in the NRHP under Criterion A and B. Site 22LO1117 will be avoided by Project implementation.

While not recommended NHRP-eligible individually, four of the newly recorded archaeological sites (22LO1114, 22LO1115, 22LO1116, and 22LO1119) are recommended as NRHP-eligible as

part of an archaeological district. The sites represent former tenant occupations associated with the Oakland Plantation and are recommended NRHP-eligible under Criterion A. The four sites reflect evidence of land use relevant to the context of tenant farming in the region, a group traditionally underrepresented in both the archaeological and archival records. Additionally, the artifact assemblages reflect potential associations with the material cultural of this group during the period of significance (1830-1930). No other information would likely be gained via additional archaeological investigations of the sites, and they are not recommended NRHP-eligible under Criterion D. These four sites, and a 100-foot buffer around each site, will be avoided by Project implementation. As they are not associated with any other built or landscape features that would contribute to the district, avoidance of physical impacts would avoid adverse effects to the district under Section 106.

The remaining six, non-isolated find, archaeological sites (22LO1099, 22LO1101, 22LO1118, 22LO1120, 22LO1121, and 22LO1122), that were not part of the archaeological district assessment, or whose NRHP recommendations have been previously discussed, have a NRHP recommendation of not eligible.

# 3.8.2 Environmental Consequences

#### 3.8.2.1 No Action Alternative

Under the No Action Alternative, existing land use would be expected to remain unchanged. Ground-disturbing industrial agricultural practices currently carried out through portions of the site would continue to have the potential to impact intact cultural resources at the surface or within the first 8 to 10 inches of soil. The impact of livestock within newly recorded archaeological sites was noted during the cultural resources survey. Livestock will continue to have a negative impact on cultural resources at the site. If the site were instead redeveloped for residential, commercial, or industrial land uses, then more significant impacts to existing or potential cultural resources would be expected.

### 3.8.2.2 Proposed Action

A total of 25 historic-age non-archaeological resources on 7 properties were recorded during the Phase I cultural resources survey, in addition to 8 individual resources on 5 parcels within the Non-Physical APE that were previously evaluated as part of the MS Solar 5 Project cultural resources survey. Of the accessible resources within the current survey, two properties (the Smith Oaks Plantation former commissary [Resource 02a] and the Oakland Plantation dwelling [Resource 05a] are recommended for NRHP inclusion, and two individual resources associated with the Oakland Plantation (Resources 05c and 05d) are recommended as contributing. Two additional resources would require additional NRHP analysis to confirm their eligibility status (the Smith Oaks Plantation Cemetery [Resource 02h] and a former African American cemetery [Resource 07]. The resources recommended for NRHP inclusion and of undetermined NRHP eligibility status would not be adversely affected by the Project. Specifically, the resources are removed in distance and shielded by vegetation and other setting intrusions. As a result,

construction of the Project would not affect any of the characteristics that qualify the resources for NRHP inclusion.

None of the remaining resources recorded during the survey, either individually or collectively, appear to meet the criteria for NRHP inclusion due to a lack of significance and/or architectural integrity. As no historic (NRHP-eligible or listed) resources would be physically or otherwise adversely affected by the proposed Project, no further consideration of effects to historic properties under Section 106 of the NHPA is recommended in connection with the Project as currently proposed.

Thirteen newly identified archaeological sites and thirteen isolated finds were encountered within the Physical APE during the Phase I cultural resource surveys. One of the thirteen newly identified sites, 22LO1117, is recommended eligible for inclusion in the NRHP.

Site 22LO1117, the Thomas Wilburn II harness horse racing track is recommended eligible for inclusion in the NRHP under Criteria A and B. Wilburn constructed the track in the late 1940s or early 1950s and trained 1,667 Standardbred racehorses at this location. Some of whom were sold to influential families, including the Hunts and Mellons. Until his retirement from horse racing in 1974, Wilburn won over 2,000 races throughout the United States and Canada, eventually being inducted into the Illinois Harness Horseracing Hall of Fame. According to local informants, when Wilburn held races at his track, which is one of the ways he marketed his horses to potential buyer, upwards of 5,000 people, both local and from far away, would attend. Wilburn was significant in the world of harness horse racing and the horse racing community; his races and influence was significant within the local and regional communities; and much can be learned from how he constructed and maintained his track.

The Wilburn racetrack represents a track whose primary use was training and marketing, as opposed to a strictly spectator event location. Roughly two-thirds of the track is currently located in the Physical APE and is under heavy vegetation. The portion that is outside of the Physical APE is maintained and covered in short grasses. The portion outside of the Physical APE has not been investigated for cultural material, though any recommendation for inclusion in the NRHP should include the entirety of the racetrack. The portion of the harness horse racing track within the Physical APE will be avoided by project implementation. A 100-foot avoidance buffer has been placed around 22LO1117.

One newly identified precontact Native American archaeological site, 22LO1100, identified as a Late Archaic and Late Woodland period precontact Native American occupation / campsite, and is located in an active agricultural field that has only been in use since 2017. The site consists of twenty-six lithic artifacts and a single sand tempered pottery sherd. Of the twenty-six lithic artifacts four are diagnostic projectile points or knives. The projectile points / knives include a likely Flint Creek projectile point, a Baker's Creek projectile point, a partial Dehli or Smithsonian projectile point, and a Variety Russel of the Nodena point type. All lithic material from 22LO1100 appears to be from heat treated Tuscaloosa Gravels. Since the location of 22LO1100 has only been utilized for agriculture for a relatively short period of time, since 2017, it is likely that there have

been minimal impacts to any potential, not discovered, subsurface features. For this reason, site 22LO1100 is recommended as unevaluated for inclusion in the NRHP. Site 22LO1100 will be avoided by direct impacts from the Project. A 100-foot avoidance buffer has been placed around 22LO1100.

Two sites with intact features, identified as historic occupations depicted on the 1911 Soils Map of Lowndes County, were identified during the current investigations. 22LO1098 is a moderate historic scatter with extant features located within a wooded area. The extant features include a cistern, that has been filled with cinder blocks, bricks and rubbish, and a small brick lined foundation. Site 22LO1114 is a subsurface historic scatter located in a mowed hayfield at the time of survey. One shovel test indicated an intact burn feature. It is unknown if the sites are associated with tenant farms, sharecroppers, or descendants of the original plantation owners. Information gained from either site could provide valuable insights into the lifeways and land use by farmers in rural Mississippi during the Reconstruction period to post-Depression era. Site 22LO1098 is recommended as eligible for inclusion in the NRHP under Criterion A and D. An individual archaeological assessment of site 22LO1114 recommends it as unevaluated for inclusion in the NRHP. Both sites will be avoided by direct impacts from the Project. A 100-foot avoidance buffer has been placed around 22LO1098 and 22LO1114.

Four archaeological sites (22LO1114, 22LO1115, 22LO1116, and 22LO1119) are recommended as NRHP-eligible as part of the newly identified Oakland Plantation Historic Archaeological District. The four sites likely represent former tenant occupations and are recommended NRHP-eligible under Criterion A. No additional information would likely be gained from additional archaeological investigations, and they are not recommended NRHP-eligible under Criterion D. These four sites, plus a 100-foot buffer, will be avoided by Project implementation. As they are not associated with any other built or landscape features that would contribute to the district, avoidance of physical impacts would avoid adverse effects to the district under Section 106.

The remaining six newly identified archaeological sites (22LO1099, 22LO1101, 22LO1118, 22LO1120, 22LO1121, and 22LO1122), not part of the Oakland Plantation Historic Archaeological District or previously discussed, and 13 isolated find locations within the Physical APE are recommended not eligible for inclusion in the NRHP. No further archaeological investigations area recommended at these locations.

Throughout the Section 106 consultation process, if it is determined that the Project would result in an adverse effect on cultural resources, the Project would be redesigned to avoid affected sites so that MS Solar 6 would not need to mitigate for impacts through an MOA process. Therefore, there would be no direct or indirect impacts to archaeological or historic resources listed eligible, potentially eligible, or undetermined for the NRHP. TVA consulted with the SHPO and federally recognized Indian tribes with an interest in the area with respect to TVA's findings of both the archaeological and architectural surveys. MDAH concurred with TVA's findings on May 27, 2022, which concluded the Section 106 consultation process. Accordingly, the requirements of Section 106 of the National Historic Preservation Act have been met.

Should previously undiscovered cultural resources be identified during Site construction or operations, construction in the affected area will be immediately stopped and the discovery location secured against further disturbance, pending completion of consultation with appropriate stakeholders. TVA and the SHPO would be consulted before any further action were taken.

#### 3.9 Utilities

This section describes an overview of existing utilities within the Project Site and the immediate surrounding area and the potential impacts on these utilities that would be associated with the No Action and Proposed Action Alternatives. Specific utility components analyzed below include telecommunications, electricity, natural gas, water, and sewer.

#### 3.9.1 Affected Environment

The Project Site is located on farmland in a rural area of Lowndes County, Mississippi. Available power sources to the county residents are electricity and natural gas. No significant renewable energy sources are currently located in this area.

#### 3.9.1.1 Telecommunications

Telecommunication services in the area are provided primarily by Franklin Telephone Company and Bellsouth Telecommunications, as well as mobile providers.

# 3.9.1.2 Electricity

The local electricity provider for most of Lowndes County as well as the adjacent counties is 4-County Electric Power Association (4CEPA), a not-for-profit electric cooperative that purchases power generated by TVA. A TVA-owned transmission line traverses the Project Site, cutting east-west across the northern portion of the site (see Figure 2-2). Distribution lines are present throughout the area, including along portions of Gilmer-Wilbum Road and other major and minor roads in the vicinity.

#### 3.9.1.3 Natural Gas

Natural gas service in the vicinity is provided by Atmos Energy, which owns a natural gas pipeline that taps the Southern Natural Gas pipeline at a meter station located on a parcel that abuts the Golden Triangle I Solar and BESS Project Site.

## 3.9.1.4 Water and Sewer

Because the Project Site is predominantly outside of incorporated municipality limits, water service is provided either by the Prairie Land Water Association or through private wells, and sewer service is provided through private septic systems. No known public service water lines or line markers servicing individual customers were observed on the Project Site. The residents located adjacent to the Project Site may have water service from Prairie Land Water Association or other public utility companies.

# 3.9.2 Environmental Consequences - Utilities

This section describes the potential impacts to utilities should the No Action or Proposed Action Alternatives be implemented.

#### 3.9.2.1 No Action Alternative

Under the No Action Alternative, there would be no Project-related impacts to utilities. Existing land use would be expected to remain a mix of agricultural and forested land, and existing on-site utilities would likely remain unchanged, with the exception of potential upgrades and maintenance.

### 3.9.2.2 Proposed Action

Under the Proposed Action, minimal temporary impacts to local electricity service could be expected when bringing GT2 on-line or during routine maintenance of the facility. Once the Project enters the operation phase, 4-County Electric Power Association (4CEPA) would provide the required back-up power for controls. Based on discussions with 4CEPA and given the low level of retail electric demand needed for the facility, no changes to the 4CEPA distribution system would be expected, and there would be no impacts to the local utility or its customers. Implementation of the Proposed Action would result in additional renewable energy resources in the region, which would augment total electricity supply and constitute an improvement to the environmental impacts associated with regional electricity generation.

Water would be needed for soil compaction and dust control during construction, and to a lesser extent for domestic use during operations (i.e., cleaning solar panels if drought conditions persisted). There would be no habitable buildings on-site and no need for potable water. Portable toilet facilities would be available on-site for the duration of the construction period; there would be no need for a septic system or connection to the closest sanitary sewer. Water in sufficient quantity and quality would be made available through use of on-site groundwater wells, or delivery via water trucks. MS Solar 6's construction contractor would determine daily water requirements based on the preliminary grading plan and size the new on-site wells accordingly. MS Solar 6's construction contractor would perform groundwater drilling and testing work prior to full construction to generate data on aquifer characteristics and develop a plan for the production well design. If existing groundwater wells are available and needed, they would be utilized for the Project; the exact location of which would be identified in the final design. The wells would be spaced around the Project Site to provide easy access for construction water and to reduce the potential for any significant water level drawdown.

Natural gas service would not be required for the Project, and the existing natural gas infrastructure present near GT2 would not be disturbed or otherwise impacted during the construction or operation of the Project.

No communication resources are anticipated to be acquired through the local providers. MS Solar 6 would have a dedicated communications system to remotely monitor the Project facility and operations.

No long-term adverse impacts to utilities would be anticipated as a result of implementation of the Proposed Action. No indirect impacts to utilities would occur under the Proposed Action. The Proposed Action would result in a new source of renewable energy for the region; thus, the Project would ultimately contribute to a beneficial impact to electrical services across the region.

# 3.10 Waste Management

This section provides an overview of existing waste management within the surrounding area of the Project Site and the potential impacts to waste management that would be associated with the No Action or Proposed Action Alternatives. Components of waste management that are analyzed include solid and hazardous waste and materials.

#### 3.10.1 Affected Environment

"Hazardous materials" and "hazardous waste" are substances which, because of their quantity, concentration, or characteristics (physical, chemical, or infectious), may present a significant danger to public health and/or the environment if released. These substances are defined by the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA; 42 U.S.C. §§ 9601 et seq.) and the Solid Waste Disposal Act, as amended by the Resource Conservation and Recovery Act ([RCRA]; 42 U.S.C. §§ 6901 et seq.). Regulated hazardous wastes under RCRA include any solid, liquid, contained gaseous, or semisolid waste or combination of wastes that exhibit one or more of the hazardous characteristics of ignitability, corrosivity, toxicity, or reactivity, or is listed as a hazardous waste under 40 CFR part 261. Storage and use of hazardous materials and wastes are regulated by local, state, and federal guidance including the Emergency Planning and Community Right-to-Know Act (42 U.S.C. §§ 116 et seq.) and RCRA.

Because the current land use of the Project Site is entirely either agricultural or undeveloped, no known hazardous waste exists on the site, and none is anticipated aside from potential petroleum, pesticides, herbicides, or fertilizers that can be removed as part of the construction process.

Collection and disposal of solid waste in Lowndes County is conducted by Golden Triangle Waste Services. Nonhazardous wastes, including construction wastes, can be hauled to an operating Class I facility.

#### 3.10.2 Environmental Consequences

This section describes the potential impacts to waste management resources should the No Action or Proposed Action Alternatives be implemented.

#### 3.10.2.1 No Action Alternative

Under the No Action Alternative, the proposed Solar Facility would not be constructed; therefore, no Project-related impacts to waste management resources would occur. Existing land use would be expected to remain a mix of agricultural and undeveloped land, and existing waste management conditions would be expected to remain as they are at present.

# 3.10.2.2 Proposed Action

Construction of the Proposed Action would result in the generation of hazardous and nonhazardous solid waste in the form of construction debris, grading spoils, packaging materials, and general construction waste. Under the Proposed Action, every effort would be made to minimize the amount of waste generated during and after construction of the Project.

Materials suitable for soil compaction are anticipated to be procured from within the Project Site as needed and off-loaded at the designated road or building location for immediate dispersion. Materials unsuitable for compaction, such as mowed debris, would be removed and loaded immediately for subsequent disposal at an acceptable off-site location. Contaminated grading and mowing materials are not anticipated; however, if any such materials are encountered during excavation, they would be disposed of at the nearest appropriate facility in accordance with applicable laws, ordinances, regulations, and standards. It is estimated that not more than 20 cubic yards of construction debris and material waste would be generated each week (during heavier periods of construction), which would be accumulated in a construction debris container and hauled off monthly. A list of acceptable waste facilities is listed in Table 3-11.

Landfill **Address Materials** Class I rubbish landfill for yard waste, 2221 Armstrong Rd, Columbus, MS 39702 Columbus Landfill construction debris, and furniture/appliances. 9778 Old West Point Rd. Household garbage, mixed building debris, Golden Triangle Regional Starkville, MS 39759 vegetative debris, commercial wastes, Solid Waste office wastes, packaging wastes and other Management Authority non-hazardous solid wastes.

Table 3-11: Waste Facilities Near the Project Site

### Hazardous Waste

Small quantities of hazardous wastes would be generated during construction, operation and maintenance, and decommissioning of the Project. Hazardous wastes generated during the construction phase would include substances such as paint and primer, thinners, and solvents. Hazardous solid and liquid waste streams that would be generated during operation of the Project include substances such as used hydraulic fluids, used oils, greases, filters, etc., as well as fluorescent light bulbs, spent cleaning solutions, and spent batteries. Hazardous wastes generated during decommissioning would include substances such as carbon dioxide, diesel fuel, hydraulic fuel, and lube oil. To the extent practicable, hazardous wastes would be recycled. Waste collection and disposal would be conducted in accordance with applicable federal, state, and local regulatory requirements to minimize health and safety effects.

MS Solar 6 (or its contractor) would report any spills to MDEQ. A sampling and cleanup report would be prepared and sent to the agency to document each spill and clean up. Each spill,

regardless of amount, would be cleaned up within 48 hours and a spill report completed. Copies of spill and cleanup reports would be kept on-site.

Minimal amounts of petroleum fuel would be kept on-site during construction. BMPs would be implemented in order to minimize the potential of a spill and to instruct on-site workers on how to contain and clean up any potential spills. The Project Site would be surrounded by security fencing during both construction and operational phases and access gates would normally remain locked. General public health and safety would not be at risk in the event of an accidental spill on-site.

During construction, hazardous materials would be stored on-site in storage tanks, vessels, or other appropriate containers specifically designed for the characteristics of the materials to be stored. The storage facilities would include secondary containment in case of tank or vessel failure. Construction- and decommissioning-related hazardous materials used for development of the Project could include gasoline, diesel fuel, oil, lubricants, and small quantities of solvents and paints. Material Safety Data Sheets for applicable materials present on-site would be made readily available to on-site personnel.

Fueling of some construction vehicles and other mobile equipment would occur primarily in the construction laydown area. Special procedures would be identified to minimize the potential for fuel spills, and spill control kits would be carried on refueling vehicles for activities such as refueling, vehicle or equipment maintenance procedures, waste removal, and tank clean-out. Fuel for construction equipment could be provided by a fuel truck or could be stored in aboveground double-walled storage tanks with built-in containment. The volume of each individual tank would not exceed 1,320 gallons, the threshold above which a Spill Prevention, Countermeasure and Control (SPCC) Plan would be required (40 CFR 112). However, because there would be fuel in reserve for diesel generators, in addition to the volume of oil contained in the main electrical transformers, the total volume of regulated materials may exceed the threshold. In that case, an SPCC Plan would be prepared.

The SPCC Plan would include procedures, methods, and equipment supplied during construction to prevent discharges from reaching navigable waters. The facility would fall under USEPA's SPCC applicability as a Tier I Qualified Facility. Because oil storage would consist of only "oil-filled operational equipment," double-walled protection would not be required [40 CFR 112.7(k)(2)] and the SPCC Plan would not have to be certified by a Professional Engineer [40 CFR 112.3(g)]. The SPCC plan would be prepared and implemented prior to and during construction to prevent oil discharges during facility operation. The administering agency is the EPA.

At the end of its useful life, the Project facilities would be decommissioned and dismantled, restoring the site. During decommissioning, above ground equipment and below ground electrical connections would be removed from the Project Site. In addition, concrete pads and foundations would be broken and removed, underground utilities would be abandoned, compacted areas would be scarified, and soils would be stabilized. The majority of decommissioned materials and equipment would be recycled. MS Solar 6 would seek a processor to recycle the solar panels to the highest degree practicable. Materials that cannot be recycled would be disposed at approved

facilities in accordance with applicable federal, state, and local laws and regulations. Alternatively, the Project facilities may be repurposed for new solar technologies available at the end of the plant lifecycle, where equipment, cabling, and foundations would be re-used where practicable. Hazardous materials that could be present during construction and decommissioning of the Proposed Action are included in Table 3-12.

Table 3-12: Summary of Special Handling Precautions for Large Quantity Hazardous Materials

Hazardous Material	Use	Relative Toxicity <sup>1</sup> and Hazard Class <sup>2</sup>	Permissible Exposure Limit	Storage Description; Capacity	Storage Practices and Special Handling Precautions
Diesel Fuel	Equipment Generator refueling and emergency diesel fire pump	Low toxicity; Hazard class – combustible liquid	PEL; none established TLV: 100 mg/m <sup>3</sup>	Carbon steel tank (3,600 gallons)	Secondary containment, overfill protection, vapor recovery, spill kit.
Hydraulic fluid (if applicable)	Tracker drive units	Low to moderate toxicity; Hazard class – Class IIIB combustible liquid	TWA (oil mist); 5 mg/m <sup>3</sup> STEL: 10 mg/m <sup>3</sup>	Hydraulic drive tank, approx. 20 gallon per tracker drive unit (if applicable) throughout solar field. Carbon steel tank, maintenance inventory in 55-gallon steel drums.	Found only in equipment with a small maintenance inventory. Maintenance inventory stored within secondary containment, alternative measures to secondary containment for equipment would be implemented at the Project.
Lube Oil	Lubricate rotating equipment (e.g., tracker drive units)	Low toxicity Hazard class – N/A	None established	Carbon steel tank, maintenance inventory in 55- gallon steel drums.	Secondary containment for tank and for maintenance inventory.

PEL = permissible exposure limit

TLV = threshold limit value

TWA = time weighted average

STEL = short-term exposure limit

<sup>1</sup>Low toxicity is used to describe materials with an NFPA Health rating of 0 or 1. Moderate toxicity is used to describe materials with an NFPA rating of 2. High toxicity is used to describe materials with an NFPA rating of 3. Extreme toxicity is used to describe materials with an NFPA rating of 4.

<sup>2</sup>N/A denotes materials that do not meet the criteria for any hazard class defined in the 1997 Uniform Fire Code.

In addition to the chemicals listed above, small quantities (less than 55 gallons, 500 pounds or 200 cubic feet) of janitorial supplies, office supplies, laboratory supplies, paint, degreasers, herbicides, pesticides, air conditioning fluids (chlorofluorocarbons), gasoline, hydraulic fluid, propane, and welding rods typical of those purchased from retail outlets may also be stored and used at the facility. Flammable materials (e.g., paints, solvents) would be stored in flammable material storage cabinet(s) with built-in containment sumps. Due to the small quantities involved and the controlled environment, a spill could be cleaned up without significant environmental consequences.

MS Solar 6 would develop and implement a variety of plans and programs to ensure safe handling, storage, and use of hazardous materials (e.g., Hazardous Material Business Plan). Facility personnel would be supplied with appropriate personal protective equipment (PPE) and would be properly trained in the use of PPE as well as the handling, use, and cleanup of hazardous materials used at the facility and the procedures to be followed in the event of a leak or spill. Adequate supplies of appropriate cleanup materials would be stored on site.

# Non-hazardous Waste Management

Construction, operation, and maintenance, and decommissioning of the Project would generate non-hazardous solid wastes. Facility-related wastes generated during all phases of the Proposed Action would include soiled rags, worn or broken metal and machine parts, defective or broken electrical materials, other scrap metal and plastic, insulation material, empty containers, cardboard, glass, wood/pallets, and other miscellaneous solid wastes including the typical refuse generated by workers. These materials would be disposed of by means of contracted refuse collection and recycling services. Waste collection and disposal would be in accordance with applicable federal, state, and local regulatory requirements to minimize health and safety effects, and no waste would be treated on site during construction or operations. Designated contractor and subcontractor personnel would be responsible for daily inspection, cleanup, and proper labeling, storage, and disposal of all refuse and debris produced. Disposal containers such as dumpsters or roll-off containers would be obtained from a proper waste disposal contractor. Records of the amounts generated would be provided to MS Solar 6. Information on universal wastes anticipated to be generated during Project construction is provided in Table 3-13.

Table 3-13: Summary of Construction Waste Streams and Management Methods

Waste Stream	Origin and Composition	Estimated Frequency of Generation	On-site Treatment	Waste Management Method / Offsite Treatment
Construction waste	Empty material containers	Intermittent	None	Return to vendor
Construction Waste	Used oil, hydraulic fluid, oily rags	Intermittent	None	Recycle wherever practicable, remove to off-site disposal location
Construction Waste	Steel, glass, plastic, wood/pallets, cardboard,paper	Intermittent	None	Recycle wherever practicable, otherwise dispose of at a class I landfill
Sanitary waste	Portable chemical toilets – sanitary waste	Periodically pumped to tanker truck by licensed contractors	None	Ship to sanitary wastewater treatment facility.

The anticipated quantities of waste produced during Project operation are summarized in Table 3-14. Universal wastes and unusable materials produced as a result of implementation of the Proposed Action would be handled, stored, and managed in accordance with Mississippi Universal Waste requirements.

Waste Stream	Origin and	Estimated	Estimated Frequency	Waste Management Method	
waste otteam	Composition	Volume	of Generation	Onsite	Offsite
Used hydraulic fluid, oils, and grease- petroleum-related wastes	Tracker drives, hydraulic equipment	1,000 gallons/year	Intermittent	Accumulate for <90 days	Recycle
Oily rags, oil absorbent, and oil filters – petroleum-related wastes	Various	One 55-gallon drum/month	Intermittent	Accumulate for <90 days	Send offsite for recovery or disposed at Class I landfill
Spent batteries	Lead acid/lithium ion	1,000	Every 10 years	Accumulate for <90 days	Recycle

Table 3-14: Summary of Operational Waste Streams and Management Methods

Waste collection and disposal would be conducted in accordance with applicable federal, state, and local regulatory requirements to minimize health and safety effects. To the extent practicable, waste would be recycled. Materials that could not be recycled would be disposed of at an approved facility to be determined by the designated contractor(s) in accordance with applicable federal, state, and local laws and regulations. No waste oil would be disposed of on the Project Site.

If necessary, MS Solar 6 or its contractor would obtain a hazardous waste generator identification number from the State of Mississippi prior to generating any hazardous waste. Any spills related to the Project would be reported to MDEQ's Emergency Response Division. A sampling and cleanup report would be prepared for the Project Site and sent to MDEQ to document each spill and clean up. Each spill, regardless of amount, would be cleaned up within 48 hours and a spill report would be completed. Copies of any spill and cleanup reports would be kept on site.

Designated contractor and subcontractor personnel would be responsible for daily inspection, cleanup, and proper labeling, storage, and disposal of all refuse and debris produced. Disposal containers such as dumpsters or roll-off containers would be obtained from a proper waste disposal contractor. Records of the amounts generated would be provided to the designated Solar Facility environmental specialist.

# 3.11 Public and Occupational Health and Safety

This section describes an overview of existing public health and safety at the Project Site and the potential impacts to public health and safety associated with the No Action and Proposed Action Alternatives. Analyzed issues include emergency response and preparedness and occupational

or worker safety in compliance with the Occupational Safety and Health Administration (OSHA) standards.

### 3.11.1 Affected Environment

The Project Site is currently private property, with a predominant land use activity of agricultural and livestock. Public emergency services in the area include fire protection services, law enforcement services, urgent care clinics, and hospitals. Fire protection services are provided by the Columbus Fire Department and the Starkville Fire Department, located approximately 15 miles (20 minutes) and 11 miles (16 minutes), respectively, from the Project Site. Law enforcement services in the City of Starkville are provided by the Starkville Police Department. Lowndes County law enforcement services are provided by the Lowndes County Sheriff's Department in Columbus, approximately 12 miles (17 minutes) northeast of the Project Site. The State Urgent Care Center, located on South Montgomery Street, approximately 11 miles (16 minutes) northwest of the Project Site, is the closest urgent care center to the Project Site. The North Mississippi Medical Center is the closest hospital, located in West Point approximately 16 miles (19 minutes) north of the Project Site. The Mississippi Emergency Management Agency has the responsibility and authority to coordinate with state and local agencies in the event of a release of hazardous materials.


### **Glint and Glare**

Glint is typically defined as a momentary flash of bright light, often caused by a reflection off a moving source. Atypical example of glint is a momentary solar reflection from a moving car. Glare is defined as a continuous source of bright light. Glare is generally associated with stationary objects, which, due to the slow relative movement of the sun, reflect sunlight for a longer duration. The difference between glint and glare is duration. Industry-standard glare analysis tools evaluate the occurrence of glare on a minute-by-minute basis; accordingly, they generally refer to solar hazards as "glare" (ForgeSolar 2021).

Due to the proposed Project's proximity to the GTR Airport, a glare analysis was performed in accordance with Federal Aviation Administration (FAA) standards. To understand and model glare in accordance with FAA's standards, Sandia National Laboratories developed the Solar Glare Hazard Analysis Tool (SGHAT). To perform the glare analysis for this study, the SGHAT, as licensed to ForgeSolar, was utilized (ForgeSolar 2021). The SGHAT allows the user to specify a site location, draw an outline of the proposed photovoltaic array, and specify observer locations. Once these points are given, the properties of the arrays such as the tracking type, tilt, module surface type, and orientation can be specified as well for each array. Latitude, longitude, and elevation for each observation point and array vertex are tracked and used for sun position and vector calculations to determine glare for that observation point. Additional information regarding reflectance, environment, and ocular factors can be altered; however, typical values were already provided.

The ocular impact of glare is visualized with the Solar Glare Ocular Hazard Plot (SGOHP). This chart displays the ocular impact as a function of glare subtended source angle and retinal

irradiance. Each minute of glare is displayed on the chart as a small circle in its respective hazard zone. For convenience, a reference point is provided which illustrates the hazard from viewing the sun without filtering, i.e., staring at the sun. Each plot includes predicted glare for one PV array and one receptor (ForgeSolar 2021).



Source: ForgeSolar 2020

Figure 3-13: Solar Glare Ocular Hazard Plot

If glare is found, the SGHAT calculates the retinal irradiance and size/distance of the glare source, defines how many minutes of "green glare," "yellow glare," and "red glare" exist at each observation point, and produces the SGOHP. Other results from the SFHAT are a plot that specifies when glare would occur throughout the year and at what times with color codes indicating the potential ocular hazard.

The FAA established an interim policy in 2013 relating to glare from solar projects. The FAA determined that for pilots, no yellow or red glare is allowable on approach, green glare is acceptable on approach, and there are no restrictions for when regularly flying the plane. See below for exact wording on page 2 of Federal Register / Vol. 78, No. 205 / Wednesday, October 23, 2013 / Notices:

"No potential for glare or "low potential for afterimage" along the final approach path for any existing landing threshold or future landing thresholds (including any planned interim phases of the landing thresholds) as shown on the current FAA-approved Airport Layout Plan (ALP). The final approach path is defined as two (2) miles from fifty (50) feet above the landing threshold using a standard three (3) degree glidepath" (FAA, 2013).

# 3.11.2 Environmental Consequences

This section describes the potential impacts to public and occupational health and safety if the No Action or Proposed Action Alternatives is implemented.

### 3.11.2.1 No Action Alternative

Under the No Action Alternative, the proposed Solar Facility would not be constructed; therefore, no Project-related impacts on public health and safety would result. Existing land use would be expected to remain a mix of agricultural and pastureland, and existing public health and safety issues would be expected to remain as they are at present.

# 3.11.2.2 Proposed Action

Based on typical hazards associated with construction, workers on the Project Site would have an increased safety risk during construction. Due to increased safety risks associated with construction, standards such as those developed by OSHA would be implemented to maintain health and safety on all construction sites. Health and safety plans in compliance with OSHA standards outline and implement BMPs for site safety management in efforts to minimize potential health and safety risks to workers. BMPs include initial site safety orientations for all personnel on-site; development of work procedures and programs for site specific activities; use of equipment safety measures, emergency stop-work procedures, lockout and tag out procedures, general site housekeeping, and personal protective equipment; safety inspections; and plans and procedures to identify and resolve potential safety hazards.

Fuel for construction vehicles may be stored on-site during construction. An SPCC plan would be developed and implemented to minimize the potential of a spill and would include detailed instructions for on-site personnel on how to contain and clean up any potential spills. Any hazardous materials stored within the Project Site during construction would be secure and not accessible to the general public. General public health and safety would not be at risk in the event of an accidental spill on site. Emergency response for any potential incidents at the Project Site would be provided by the local, regional, and state law enforcement, fire, and emergency responders, as described in the prior section.

A minor increase in potential public health and safety hazards could result from increased traffic on local roadways during construction of the Project. The few residential sites along roadways used by construction traffic to access the Project Site would experience increased commercial and industrial traffic during construction. Established traffic procedures and awareness of these residences would be implemented in the health and safety plans to minimize potential safety concerns associated with the temporary increase in traffic.

No public health or safety hazards would be anticipated from operation of the Solar Facility. Impacts to public health and safety associated with implementation of the Proposed Action would be temporary and minor.

# Results of the Glare Analysis

A Solar Glare Ocular Impact Analysis was performed for the proposed Project to demonstrate that any glare created from the Project would not adversely impact surrounding properties, vehicles traveling on roadways near the Project Site, or pilots approaching the GTR airport. As part of the analysis, observation points were identified, SGHAT Analysis was performed, Line-of-Sight Analysis was performed, View Angle Analysis was performed, and a landscape review was performed.

The intent of the analysis was not to illustrate that all potential glare has been removed, but to identify the glare that could exist and determine if the glare would adversely impact surrounding properties, vehicles traveling along nearby roadways, or pilots approaching the GTR airport. It was found that the Project would not produce any glare that could cause permanent eye damage due to retinal burn. Potential glare at all observation points was categorized as having low potential for afterimage (designated as green glare) or having potential for afterimage (designated as yellow glare). Based on the SGHAT results, onsite visual observations, a view angle analysis, and reviews of the landscaping, it was found that no observation points have or would have potential glare to adversely impact surrounding properties near the Project Site.

# 3.12 Socioeconomics

This section describes an overview of existing socioeconomic conditions near the Project Site, and the potential impacts to socioeconomic conditions that would be associated with the No Action and Proposed Action Alternatives. Components of socioeconomic resources presented include population, demographics, employment, and income.

# 3.12.1 Affected Environment

The proposed Solar Facility would be located within an unincorporated portion of Lowndes County, Mississippi. The Project Site is entirely in U.S. Census Tract (CT) 2808733.07. The total population for Lowndes County, as reported by the U.S. Census Bureau (USCB), was 59,779 in 2010 and an estimated 59,150 in 2019 (USCB, 2021a). Top employment industries for Lowndes County are education, healthcare and social services, manufacturing, and retail trade (USCB, 2021b). Lowndes County offers residents and visitors access to activities and attractions such as museums, campgrounds, biking trails, and a historic welcome center (Lowndes, 2021a).

Columbus is the county seat to Lowndes County and is the only major city in the county. Lowndes County is comprised of three towns, including Artesia, and several communities including Mayhew (Lowndes, 2021b). In 2019, Lowndes County had a labor force of approximately 27,731 with 24,392 employed and 2,142 unemployed civilians. The unemployment rate was an estimated 8.1 percent. By comparison, the unemployment rate for the state of Mississippi was an estimated 7.5%. The median household income in Lowndes County was \$50,441. Seventeen percent of

households in Lowndes County made between \$50,000 to \$74,999 in income and benefits. By comparison, the median household income in Mississippi was \$45,081 and 16.8 percent of households made between \$50,000 to \$74,999 in income and benefits (USCB, 2021b).

# 3.12.2 Environmental Consequences

This section describes the potential impacts to socioeconomic resources should the Proposed Action or No Action Alternatives be implemented. Social and economic issues considered for evaluation within the impact area include change in expenditures for goods and services and short- and long-term effects on employment and income.

# 3.12.2.1 No Action Alternative

Under the No Action Alternative, the proposed Solar Facility would not be constructed. Therefore, there would be no Project-related socioeconomic impacts within Lowndes County, including the beneficial impacts to local population, employment, and land value associated with the proposed Project.

# 3.12.2.2 Proposed Action

Under the Proposed Action, a new solar facility would be built at the Project Site. Minor adverse indirect impacts could occur on the agricultural economy of the region due to the loss of up to 495 acres of annual soybean and corn production. MS Solar 6 would be leasing the parcels for the Solar Facility which results in a higher lease payment to the landowner than agricultural production does. However, the loss of agricultural land would adversely impact the farmers working the land as well as other services that support agricultural production. These impacts would be minimal, and the economic benefit of the Project would outweigh the adverse impacts substantially.

Construction activities at the Project Site would take approximately 17 months to complete with a crew of approximately 200 to 300 workers at the site, depending on construction activities. Workers would include general laborers and electrical technicians. Work would generally occur seven days a week during daylight hours. Short-term beneficial economic impacts would result from construction activities associated with the Project, including the purchase of materials, equipment, and services and a temporary increase in employment and income. This increase would be local or regional, depending on where the goods, services, and workers were obtained. It is likely some construction materials and services would be purchased locally in Lowndes County and/or in adjacent counties. Most of the other components of the solar and transmission facilities would be acquired from outside the local area. Also, most of the construction workforce would be sought locally or within the region, while a small portion of the construction workforce might come from out of the region. The direct impact on the economy associated with construction of the Project would be short-term and beneficial.

The majority of the indirect employment and income impacts would be from expenditure of the wages earned by the workforce involved in construction activities, as well as the local workforce used to provide materials and services. Construction of the Project could have minor beneficial

indirect impacts to population and short-term impacts to employment and income levels in Lowndes County.

During operation of the Solar Facility, a full-time workforce of up to two people would be on site five days a week from 7 A.M. to 5 P.M. This workforce would manage and maintain the Solar Facility and conduct regular inspections. Grounds maintenance and some other operation and maintenance activities may be conducted by local contractors. Therefore, operation of the Solar Facility would have a small positive impact on employment and population in Lowndes County.

Overall, socioeconomic impacts for the operation of the proposed Solar Facility would be positive and long-term, but small relative to the total economy of the region. The local tax base would increase from construction of the Solar Facility and would be most beneficial to Lowndes County and the vicinity. Additionally, the local governments would not have to provide any of the traditional government services typically associated with a large capital investment, such as water, sewer, or schools.

# 3.13 Environmental Justice

This section provides an overview of environmental justice considerations within the Project Area and the potential impacts to environmental justice populations that would be associated with the No Action and Proposed Action Alternatives. Components of environmental justice that are presented include the proportions of the local population that are minority and low-income and the potential for disproportionate effects on these populations.

# 3.13.1 Affected Environment

Executive Order (E.O.) 12898 – Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations was issued in 1994 with the purpose of focusing federal attention on the environmental and human health effects of federal actions on minority and low-income populations. The E.O.'s goal is to achieve environmental protection for all communities. Per the E.O., federal agencies are directed to identify and address minority and low-income populations that are disproportionately affected by adverse human health and environmental effects to the greatest extent practicable and permitted by law (USEPA, 2020c). While not subject to this E.O., TVA routinely considers environmental justice in its NEPA review process.

Minority individuals are those who are members of the following demographics: American Indian or Alaskan Native, Asian or Pacific Islander, Black (not of Hispanic origin), or Hispanic. CEQ defines a "minority population" as a readily identifiable group of people residing in geographic proximity with a population comprised of 50 percent minority or greater or an identifiable group that has a meaningfully greater minority population than the adjacent geographic areas or may also be a geographically dispersed set of individuals such as Native Americans or migrant workers (Spies, Stine, Gravenmier, Long, & Reilly, 2018). In 2018, the minority population in Lowndes County was approximately 48.2%. By comparison, the minority population in the state of Mississippi was approximately 43.0% (USCB 2020a). Based on the USEPA's Environmental

Justice Screening and Mapping Tool (EJSCREEN), 36 percent of the population block (ID# 280870010001) that the Project is located in is considered minority population (USEPA 2020a).

A low-income population is considered a community or group of individuals that live in geographic proximity to one another, or a set of individuals such as American Indians or migrant workers who meet the standards for low income and experience common conditions of environmental exposure or effect. Low-income populations located in an affected area should be identified using the annual statistical poverty thresholds provided by the USCB's annual current population reports (Series P-60) on poverty and income. Poverty is defined as the number of individuals or families with income below a defined threshold level (typically 50 percent) (Spies et al., 2018). Lowndes County's estimated poverty rate for 2018 was 23.1%. By comparison, the state of Mississippi had a poverty rate of approximately 19.7%. (USCB 2020a). Based on the USEPA's EJSCREEN tool, 25 percent of the population block (ID#280870010001) that the Project is located in is considered low-income population (USEPA 2020).

# 3.13.2 Environmental Consequences

This section describes the potential impacts on environmental justice populations should the Proposed Action or No Action Alternative be implemented. According to the CEQ, adverse health effects to be evaluated within the context of environmental justice impacts may include bodily impairment, infirmity, illness, or death. Environmental effects may include ecological, cultural, human health, economic, or social impacts. Disproportionately high and adverse human health or environmental effects occur when the risk or rate of exposure to an environmental hazard or an impact or risk of an impact on the natural or physical environment for a minority or low-income population is high and appreciably exceeds the impact level for the general population or for another appropriate comparison group (CEQ 1997).

# 3.13.2.1 No Action Alternative

Under the No Action Alternative, there would be no changes attributable to the proposed solar Project within Lowndes County that would create disproportionately high and adverse direct or indirect impacts on minority or low-income populations.

# 3.13.2.2 Proposed Action

Based on the USEPA's online screening tool, EJSCREEN, no minority or low-income populations have been identified in the potentially affected area. Additionally, based on the analysis of impacts for all resource areas presented in this EA, it was determined that there would be no significant adverse health impacts on members of the public or significant adverse environmental impacts on the physical environment (water, air, aquatic, and terrestrial resources) and socioeconomic conditions. As there are no identified environmental justice communities in the block group within which the proposed Project is located, there would be no disproportionately high or any adverse direct or indirect impacts on minority or low-income populations due to human health or environmental effects resulting from the Proposed Action.

# 3.14 Transportation

This section describes roadways and other transportation infrastructure serving the Project Site and surrounding area, and the potential impacts on transportation resources that would be associated with the No Action and Proposed Action Alternatives. Components of transportation resources that are analyzed include roads, traffic, railroads, and airports.

# 3.14.1 Affected Environment

The area considered for transportation is located within the Northwest portion of Lowndes County, Mississippi, with the cities of Artesia and Mayhew to the north. Artesia West Point Road runs north-south along the west portion of the north site from Artesia to Mayhew. It appears to be a paved road that turns into a gravel road before it reaches the western boundary of the site. State Highway 182 is a paved road that runs east-west approximately five miles north of the Project Site. Artesia Road is a paved road that runs east-west approximately 1.5 miles north of GT2. Wilmer Gilburn Road is a county-maintained gravel road that runs east to west along the norther boundary of GT2.

Existing traffic volumes on roads in the Project Site were determined using the most recent 2018 Average Annual Daily Traffic (AADT) counts measured at existing Mississippi Department of Transportation (MDOT) stations (MDOT 2019). The 2019 AADT count for Site ID 44110, located on Artesia Road, 1.5 miles north of the Project Site, consisted of 950 vehicles. The 2018 AADT count for Site ID 44003, located on Highway 82 consisted of 20,000 vehicles. There were no AADT counts for Wilmer Gilburn Road.

The closest rail line is operated by Kansas City Southern and runs north to south approximately 1,200 feet west of GT2. The closest regional airport is the GTR Airport, located approximately 3 miles northeast of the Project Site. The airport consists of one runway that is approximately 7,860 feet long.

# 3.14.2 Environmental Consequences

This section describes the potential impacts to transportation resources should the No Action or Proposed Action Alternatives be implemented.

# 3.14.2.1 No Action Alternative

Under the No Action Alternative, the proposed solar Project would not be constructed. Therefore, no Project-related impacts to transportation resources would result. Existing land use would be expected to remain as a mix of farmland and undeveloped land, and the existing transportation network and traffic conditions would be expected to remain as they are at present.

# 3.14.2.2 Proposed Action

The construction and operation of the Project would have no effect on operation of the airports in the region. The operation of the Project would not affect commercial air passenger traffic or freight traffic in the region and would not adversely affect any aerial crop dusters operating in the vicinity of the Project Site.

During construction of the proposed Solar Facility, a crew of approximately 200 to 300 workers would be present at the Project Site between sunrise and sunset, seven days a week. A majority of these workers would likely come from the local area or region. Other workers would come from outside the region, and many would likely stay at hotels in Starkville or Columbus. It is anticipated that workers would drive personal vehicles to the Project Site. Some of the individual workers and work teams would likely visit local restaurants and other businesses during the construction phase of the Project. Additional traffic due to deliveries and waste removal would consist of approximately five vehicles per day during construction, as discussed in more detail below.

Traffic flow around the Project Site would be heaviest at the beginning of the workday, at lunch, and at the end of the workday. Deliveries and most workers would likely access the Project Site from the east along Gilmer Wilburn Road and west along Artesia Road. Should substantial traffic congestion occur, MS Solar 6 would implement staggered work shifts during daylight hours to assist traffic flow near Project Site access locations. Implementation of such mitigation measures would minimize potential adverse impacts to traffic and transportation to negligible levels.

Construction equipment and material delivery and waste removal would require approximately 20 flatbed semi-trailer trucks or other large vehicles visiting the Project Site each day during the 17-month construction period. The Project Site can be accessed via routes that do not have load restrictions. These vehicles should be easily accommodated by existing roadways; therefore, only minor impacts to transportation resources in the area surrounding the Project would be anticipated as a result of construction vehicle activity.

Several on-site access roads would be maintained within the Project Site. Access points during construction include several existing farm roads on the south side of Gilmer Wilburn Road,

The Solar Facility would be staffed by up to two full-time workers who would live in the area. The addition of vehicles for full-time staff on local roadways would be accommodated by existing infrastructure; therefore, the operation of the Project would not have a noticeable impact on the local roadways.

The overall direct impacts on transportation resources associated with implementation of the Proposed Action would be moderate during construction due to the influx of workers traveling to the job site. These impacts would be temporary and minimized through appropriate mitigation. The Proposed Action would not result in any indirect impacts on transportation.

#### 4.0 ANTICIPATED ENVIRONMENTAL IMPACTS AND CUMULATIVE IMPACTS

This chapter summarizes the anticipated adverse environmental impacts of the Project and considers the relationship between short-term uses and long-term productivity and whether the Project makes irreversible and irretrievable commitments of resources. This chapter also considers the cumulative impacts in relation to other ongoing or reasonably foreseeable proposed activities within the Project Site and the surrounding area.

#### 4.1 **Unavoidable Adverse Environmental Impacts**

As described in Section 2 (Table 2-1), the Proposed Action could result in some unavoidable adverse environmental effects. Specifically, construction activities would temporarily increase noise, traffic, and health and safety risks and temporarily affect air quality, GHG emissions, and visual aesthetics of the Project Site vicinity. Construction activities would primarily be limited to daytime hours, which would minimize noise impacts. Temporary increases in traffic would be minimized or mitigated by staggering work shifts and/or posting a flag person during the heavy commute periods. Temporary increases in health and safety risks would be minimized by implementation of the Project health and safety plan. Construction and operations of the Project would have minor, localized effects on soil erosion and sedimentation that would be minimized by placement of construction BMPs, early soil stabilization, and vegetation management measures. Selective maintenance of tree buffers and/or fence screening (existing vegetated fence rows) along the perimeter of the Solar Facility would minimize effects to visual resources, during both construction and operation. The Project would change land uses on the Project Site from primarily agricultural to solar uses, where these practices are not presently occurring.

Table 4-1 provides a list of impacts and proposed mitigation on environmental and human resources associated with the Project.

Impact Type Location Description Mitigation Measure Solar Facility Temporary construction Stabilize exposed soils to prevent erosion, and maintain any soil activities within floodplain. Permanent placement of stockpiles outside the boundaries of solar arrays and buried the floodplain. Once in operation, the collection lines within support structures would not impede floodplain. One or more floodplains or floodwaters. At full tilt, Floodplain access roads would cross all panels within the floodplain would be greater than 1 foot above the base floodplains. blood elevation. Roads would be constructed such that upstream flood elevations would not be increased by more than 1.0 foot. Throughout Project Access roads will be used to Utilize existing bridges and culverts cross streams. first. Where additional stream Site crossings are needed, keep Streams crossings as narrow as possible and maintain BMPs to keep sediment out of streams.

Table 4-1: Unavoidable Impacts and Proposed Mitigation for GT2

Impact Type	Location	Description	Mitigation Measure
Wetlands	Throughout Project Site	PEM, PFO and PSS wetlands occur within the Project Site	Avoid jurisdictional wetland and install silt fencing along the outside of the construction work area where wetlands are present.
Soils	Solar Facility areas occurring on agricultural land.	Permanent loss of prime farmland soils.	If fill or soil removal were needed, the topsoil would first be stripped and segregated. Once fill or other intense earthwork was complete, the topsoil would be reapplied to the surface.
Vegetation	Solar Facility	Removal of agricultural and forested vegetation resulting in a large area of exposed soil.	Seeding Solar Facility with native and non-invasive low-growing grasses and flowers that would attract pollinators.
Vegetation	Solar Facility	Prevent unintentional encroachment in avoidance areas.	Install signage and/or temporary construction fencing around avoidance areas. Identify avoidance areas on site plans and constraints maps.
Wildlife	Solar Facility	Temporary displacement of wildlife during construction. Potential direct impacts due to vegetation removal and grading/construction.	Enhance the existing Project Site by revegetating with native and/or naturalized non-invasive herbaceous plants maintained without the extensive use of harmful herbicides and pesticides.
Visual	Solar Facility in vicinity of single-family homes	Three homes within viewshed of Project Site	Avoid clearing mature trees that serve as natural buffers near homes.
Noise	Solar Facility located less than 500 feet fromone single-family home	Potential for increased noise levels to residents within 500 feet of inverters.	For the one single-family home that is within 500 feet of an inverter, a preconstruction sound study including an ambient survey would be conducted to quantify the existing ambient environment. After the project reaches commercial operation, MS Solar 6 would measure the sound levels at the property line and identify any equipment that generates a Ldn sound level that exceeds 55 dBA at the property line. If there are locations where noise levels exceed that threshold, MS Solar 6 would install sound buffers (walls, fences with screening, or vegetation) in order to minimize the noise levels from operating equipment
Transportation	Solar Facility	Delivery trucks and most workers would likely access the Project Site from Gilmer Wilburn Road. Potential for congestion, especially if trains are progressing northbound along the adjacent railroad track.	Should substantial traffic congestion occur, MS Solar 6, or its contractor, would implement staggered work shifts to assist traffic flow near Project Site access locations. Implementation of such mitigation measures would minimize potential adverse impacts to traffic and transportation to negligible levels.

With the installation and routine maintenance of appropriate BMPs, no unavoidable adverse effects to groundwater are expected. Long-term habitat loss would occur due to alteration of land use on the Project Site. Revegetation of the Project Site with native and/or noninvasive grasses and herbaceous vegetation would help minimize effects to open, grassy habitats. The Project is not expected to adversely affect any federal or state-listed species.

#### 4.2 Relationship of Short-Term Uses and Long-Term Productivity

Short-term uses are generally those that occur on a year-to-year basis, such as wildlife foraging, timber management, recreation, and uses of water resources. Long-term productivity is the capability of the land to provide resources, both market and nonmarket, for future generations. For this EA, long-term impacts to site productivity would be those that last beyond the life of the Project. The Proposed Action would affect short-term uses of the Project Site by converting it from agricultural and undeveloped land to solar power generation. However, the effects on long-term productivity would be minimal because the existing land uses could be readily restored on the Project Site following the decommissioning and removal of the Solar Facility.

#### 4.3 Irreversible and Irretrievable Commitments of Resources

An irreversible or irretrievable commitment of resources would occur if resources would be consumed, committed, or lost as a result of the Project. The commitment of a resource would be considered irretrievable if the Project would directly eliminate the resource, its productivity, or its utility for the life of the Project and possibly beyond. Construction and operation activities would result in an irretrievable and irreversible commitment of natural and physical resources. The implementation of the Proposed Action would involve irreversible commitment of fuel and resource labor required for the construction, maintenance, and operation of the Solar Facility. Because removal of the solar arrays and associated on-site infrastructure could be accomplished rather easily, and the facility would not irreversibly alter the site, the Project Site could be returned to its original condition or used for other productive purposes once the Solar Facility is decommissioned. Most of the Solar Facility components could also be recycled after the facility is decommissioned.

#### 4.4 Cumulative Impacts

A cumulative impact, as defined by the Council on Environmental Quality (CEQ), is described as an impact on the environment that results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions regardless of what agency or person undertakes such other actions. Cumulative impacts can result from individually minor but aggregately significant actions taking place over a period of time (40 CFR 1508.7). A cumulative impacts analysis acknowledges the effects of the proposed alternatives on the various environmental resources. The analysis also recognizes the effects of other past, present, and reasonably foreseeable future actions, and describes the cumulative or additive effects that may result. While some cumulative effects, however minimal, can be established for virtually any resource or condition, the effects described in this EA are considered to be the most applicable and representative of those associated with the Proposed Action. Cumulative impacts associated

with the Proposed Action are described below in the following resource sections. This section addresses the cumulative impacts of the Project when combined with any reasonably foreseeable future action in the vicinity and are depicted in Figure 4-1.

Desktop research of potential past, present, and future actions in the Lowndes County, Mississippi area was conducted. Resources examined included:

- Local and regional news sources.
- City of Starkville and City of Columbus website records, including planning commission meetings, city meeting minutes, and public notices; and
- Mississippi DOT website.

The proposed Project would result in minor direct impacts to land use, geological resources and farmlands, water resources, biological resources, visual resources, noise, air quality, public health and safety, and transportation.

# 4.4.1 Federal Projects

This section addresses other projects with potential effects to land use, geological resources and farmlands, water resources, biological resources, visual resources, noise, air quality, public health and safety, and transportation.

Based on a review of the above listed resources, and through discussions with local officials, seven projects were identified in Lowndes County that have or will have at least partial federal funding. These seven projects are part of the MSDOT's 5-Year Plan and will receive federal funding and additional funding from either the State or County (MDOT 2021). The projects include sign replacements, 3 rumble strip installation projects along existing highways, and three resurfacing projects along existing highways. The closest of these, a resurfacing project, is directly north of the Project Site on US Hwy 82. The resurfacing along Hwy 82 will occur along the existing roadway about one year before the Golden Triangle I Solar and BESS Project would begin construction. Given the nature of the impacts of the proposed Golden Triangle I Solar and BESS Project, the Proposed Action is unlikely to contribute towards adverse cumulative effects to the same resources affected by these MSDOT projects.

The Infinity Megasite is a pending 1,144-acre industrial development that is adjacent to the proposed Golden Triangle Solar Project to the east, on the north side of Artesia Road. The proposed Megasite is a TVA and Golden Triangle Development LINK (LINK) project. A megasite is a land development that is intended to promote business clusters. The organizations (in this case TVA and LINK) develop the land through permitting and infrastructure, so that it is "shovel ready" for big business. Development of the Infinity Megasite would result in the permanent conversion of approximately 250 acres of forested land and 894 acres of predominantly agricultural land to commercial/industrial land use. There would be minor impacts on smaller streams and riparian wetlands which would be mitigated through the purchase of wetland and stream mitigation credits or some other approved mitigation development process. The Golden

Triangle Solar Project combined with the Infinity Megasite Project would result in an adverse cumulative impact on agricultural land. However, as previously discussed in section 3 of this EA, once the Project is decommissioned, the Project Site could be returned to its pre-construction/preoperation use without significant effort. Therefore, the Project's contribution towards a long-term cumulative impact on agricultural land, when combined with impacts from the Infinity Megasite, would be minor. If the Golden Triangle Project and the Megasite were constructed at the same time, a minor adverse cumulative impact on air (from construction equipment emissions and fugitive dusts) and noise (from construction equipment operation) would be expected. This would be a short-term cumulative impact that would be most obvious during normal daytime working hours, and it would subside once construction was complete.

Golden Triangle I Solar and BESS Project (GT1) is a recently approved solar project that would be located north of GT2 (on the north side of Gilmer Wilburn Road. The GT1 Project Site is over 3,000 acres and would be expected to generate up to 200 MW of AC capacity with an additional 50 MW of battery storage capacity. Similar to GT2, GT1 will consist of multiple parallel rows of photovoltaic (PV) panels on single-axis tracking structures, along with DC and AC inverters and transformers. MS Solar 5 has entered into long-term lease agreements or land purchases on 29 parcels. Implementation of GT1 will result in the long-term conversion of up to 150 acres of forest and an additional temporary conversion of up 3,444 acres of pasture/agriculture to commercial/industrial use. The table below shows the total potential cumulative impacts from the three projects discussed above. Long-term impacts, such as the clearing of mature trees or the conversion of agricultural land to developed industrial land will inevitably overlap for all three projects, thus resulting in a long-term cumulative impact on those resources. Short-term cumulative impacts would occur on noise and local air quality only if two or more of the projects are under construction at the same time.




Figure 4-1. Cumulative Impact Map

Table 4-2: Potential Cumulative Impacts Associated with GT2

Impact Type	Project	Impact	Description	
Forested Uplands	Golden Triangle I Solar and BESS Project	150 acres	Long-term minor cumulative impacts on mature forested areas would occur if two or more of these projects occur within the similar timelines.	
	Infinity Megasite Development	250 acres		
	GT2	265 acres		
	Cumulative Total	665 acres		
Agricultural Land	Golden Triangle I Solar and BESS Project	3,444 acres	impacts on agricultural operations would occur if two or more of these projects occur within the similar timelines.	
	Infinity Megasite Development	894 acres		
	GT2	1,260 acres		
	Cumulative Total	5,598 acres		
Noise	Golden Triangle I Solar and BESS Project	Construction only	Short-term cumulative impacts from construction	
	Infinity Megasite Development	Construction only	equipment noise could occur if two or more of the projects have overlapping construction schedules.	
	GT2	Construction only		
	Cumulative Total	N/A		
Air Quality	Golden Triangle I Solar and BESS Project	Construction only	Short-term cumulative impacts from construction emissions and dust could occur if two or more of the projects have overlapping construction schedules.	
	Infinity Megasite Development	Construction only		
	GT2	Construction only		
	Cumulative Total	N/A	CONSTRUCTION SCHEDULES.	

If GT2 and one or more of the other projects were constructed at the same time, a short-term and minor adverse cumulative impact on air (from construction equipment emissions and fugitive dusts) and noise (from construction equipment operation) would be expected. Utilizing water trucks to spray exposed soil during construction would aid in reducing fugitive dusts. Maintaining all construction equipment in good working order would help in minimizing equipment emissions during construction. Even if all three projects were constructed at the same time, fugitive dust mitigation and maintenance of construction equipment and machines are standard best practices that would be expected regardless of whether there is an overlap in the construction schedules for these projects.

There is a possibility that if more than one project were constructed at the same time, nearby residents could experience a minor cumulative impact from construction noise. This would also be a short-term cumulative impact that would be most obvious during normal daytime working hours, and it would subside once construction was complete.

Although not quantified in the table, a cumulative increase in local and regional traffic could be experienced if the proposed Project and one or more of the other projects were to schedule its peak construction workforce at the same time. Similar to noise impacts, this would also be a

short-term cumulative impact that would be most obvious during normal daytime working hours, and it would subside once construction was complete.

Long-term minor temporary cumulative impacts on agricultural operations and mature forests would occur when the Project and one or both of the other projects are operational during the same time period. GT1 and GT2 will certainly be operational at the same time, thus prohibiting agricultural operations within those project sites for at least 20 years. However, under the terms of the conditional PPA between TVA and MS Solar 5, LLC (GT1) and MS Solar 6, LLC (for GT2), TVA would purchase the electric output generated by the proposed solar facility for an initial term of 20 years, subject to satisfactory completion of all applicable environmental reviews. As mentioned above, once the Project Sites are decommissioned, both Project Sites could be returned to their pre-construction/pre-operation use without significant effort. Therefore, the Project's contribution towards a long-term cumulative impact on agricultural/pastureland and loss of forested habitat, when combined with impacts from the GT1, would be minor.

# 4.4.2 State and Local Projects

The Project Site is within both the Town of Artesia and unincorporated Lowndes County but is largely rural and agricultural. Aside from the previously discussed County and State projects with associated federal funding, there are no known past, present, or reasonably foreseeable future projects near the proposed Project Site that, when combined with impacts from the Proposed Action, would result in adverse cumulative impact on the same resources.

# 5.0 LIST OF PREPARERS

Table 5-1 presents the members of the Project team and summarizes the expertise of each member and their contributions to this EA.

Table 5-1: GT2 Environmental Assessment Team

Name/Education	Experience	Project Role				
TVA						
J. Taylor Johnson M.S. Environmental Science, B.S. Biochemistry	7 years in environmental planning and policy and NEPA compliance.	NEPA Compliance and Project Management				
Adam Dattilo M.S. Forestry B.S. Natural Resource Conservation	17 years in ecological restoration and plant ecology, 10 years in botany	Vegetation				
Elizabeth B. Hamrick M.S. Wildlife and Fisheries Science, B.A. Biology	19 years conducting field biology, 13 years technical writing, 10 years NEPA and ESA compliance	Terrestrial Ecology, Threatened and Endangered Species				
A. Chevales Williams B.S., Environmental Engineering	15 years of experience in water quality monitoring and compliance; 14 years of NEPA planning and environmental services	Surface Water				
Craig Phillips B.S. and M.S. Wildland and Fisheries Science	13 years sampling and hydrologic determination for streams and wet weather conveyances, 12 years in environmental reviews	Aquatics				
Carrie Williamson, P.E., CFM M.S., Civil Engineering; B.S., Civil Engineering	9 years Floodplains, 3 years River Forecasting, 11 years compliance monitoring.	Floodplains				
Michaelyn Harle Ph.D., Anthropology; M.A. Anthropology; B.A. Anthropology	20 years in cultural resource management	Cultural and Historic Resources, Section 106 Compliance				
Burns & McDonnell						
Jesse Brown B.A. Biology M.S. Biology	12 years in NEPA documentation, environmental permitting, protected species evaluations, and wetland delineations.	Document Preparation, Field Survey Coordination, Field Survey Biologist, Habitat Assessment, Deputy PM				
John Fulmer B.A. Anthropology M.A. Anthropology	More than 20 years of experience surveying, documenting, and report writing for cultural and archeological resources	Senior Technical Reviewer				

Name/Education	Experience	Project Role
Olivia Haney B.S. Chemistry Certificate: Natural Resources and Environmental Studies	3 years of environmental permitting, wetland and waterbody surveys, and GIS support	Mapping and impact calculations
Ellen Pennington B.S. Ecological Restoration B.S. Renewable Natural Res.	4 years of environmental permitting and NEPA documentation.	Document Preparation
Claire Randall M.S. Biology	4 years of environmental permitting and NEPA documentation.	Document Preparation
Doug Shaver M.S. Environmental & Urban Geosciences Registered Professional Archeologist	13 years in cultural, archeological, and Native American Studies. Graduate Certification in Native American Studies. Graduate Certification in Historic Preservation	Principal Investigator and lead author for cultural resources.
Robyn Susemihl B.S. Zoology, Chemistry minor	18 years in NEPA documentation, project management, protected species analysis, and stream and wetland delineations	Project Manager, Document Preparation, Sr. Technical Review
David Thomas B.S. Biology M.S. Zoology	25+ years preparing NEPA documents, protected species evaluations and surveys, and field assessments	Senior Technical Review
Rebecca Torres B.S. Animal Biology M.S. Wildlife & Fisheries Science	4 years of environmental permitting and NEPA documentation with focus on aquatic ecology	Document Preparation

# 6.0 LITERATURE CITED

- Bullock, Lindsay K. 2008. Mammals of Mississippi 12:1-6. *Eastern Spotted Skunk*. Published December 5, 2008 by the Department of Wildlife and Fisheries, Mississippi State University.
- Burns and McDonnell. 2021a. Golden Triangle II Solar Project: Protected Species Habitat Assessment Report. April 2021.
- Burns and McDonnell. 2021b. Golden Triangle II Solar Project: Wetland Delineation Report. May.
- Chafin, Linda G. 2007. Field Guide to Rare Plants of Georgia. State Botanical Garden of Georgia, Athens, Georgia.
- City-Data.2021. City Data for Artesia, MS. <a href="http://www.city-data.com/city/Artesia-Mississippi.html">http://www.city-data.com/city/Artesia-Mississippi.html</a>. Accessed: December 13, 2021.
- Cornell University 2019a. The Cornell Lab. All About Birds. Painted Bunting Life History. <a href="https://www.allaboutbirds.org/guide/Painted\_Bunting/lifehistory">https://www.allaboutbirds.org/guide/Painted\_Bunting/lifehistory</a>. Accessed December 13, 2021.
- Cornell University 2019b. The Cornell Lab. All About Birds. Scissor-tail Flycatcher Life History. <a href="https://www.allaboutbirds.org/guide/Scissor-tailed\_Flycatcher/lifehistory">https://www.allaboutbirds.org/guide/Scissor-tailed\_Flycatcher/lifehistory</a>. Accessed December 13, 2021.
- Cornell University 2019c. The Cornell Lab. All About Birds. American Kestrel Life History. <a href="https://www.allaboutbirds.org/guide/American\_Kestrel/lifehistory">https://www.allaboutbirds.org/guide/American\_Kestrel/lifehistory</a>. Accessed October 21, 2020.
- Cornell University 2019d. The Cornell Lab. All About Birds. Osprey Life History. <a href="https://www.allaboutbirds.org/guide/Osprey/lifehistory">https://www.allaboutbirds.org/guide/Osprey/lifehistory</a>. Accessed December 13, 2021.
- Cornell University. 2019e. The Cornell Lab. All About Birds. Lesser Yellowlegs Life History. <a href="https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory">https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory</a>. Accessed December 213, 2021.
- Cornell University. 2019f. The Cornell Lab. All About Birds. Marbled Godwit Life History. <a href="https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory">https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory</a>. Accessed October 20, 2021.
- Cornell University 2019g. The Cornell Lab. All About Birds. Red-headed Woodpecker Life History. <a href="https://www.allaboutbirds.org/guide/Red-headed\_Woodpecker/lifehistory">https://www.allaboutbirds.org/guide/Red-headed\_Woodpecker/lifehistory</a>. Accessed October 20, 2020.
- Cornell University 2019h. The Cornell Lab. All About Birds. Rusty Blackbird Life History. <a href="https://www.allaboutbirds.org/guide/Rusty\_Blackbird/lifehistory">https://www.allaboutbirds.org/guide/Rusty\_Blackbird/lifehistory</a>. Accessed October 20, 2020.

- Cornell University 2019i. The Cornell Lab. All About Birds. Short-billed Dowitcher Life History. <a href="https://www.allaboutbirds.org/guide/Short-billed\_Dowitcher/lifehistory">https://www.allaboutbirds.org/guide/Short-billed\_Dowitcher/lifehistory</a>. Accessed October 20, 2020.
- Cornell University 2019j. The Cornell Lab. All About Birds. Willet Life History. <a href="https://www.allaboutbirds.org/quide/Willet/lifehistory">https://www.allaboutbirds.org/quide/Willet/lifehistory</a>. Accessed October 20, 2020.
- Council on Environmental Quality (CEQ). 1997. Environmental Justice Guidance Under the National Environmental Policy Act. Available at: <a href="https://ceq.doe.gov/docs/ceq-regulations-and-guidance/regs/ei/justice.pdf">https://ceq.doe.gov/docs/ceq-regulations-and-guidance/regs/ei/justice.pdf</a>. 10 December 1997.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. *Classification of Wetland and Deepwater Habitats of the United States*. Washington, D.C.: U.S. Fish and Wildlife Publication FWS/OBS-79/31. Accessed 14 September 2020.
- Federal Emergency Management Agency (FEMA). 2021a. National Flood Insurance Program. Accessed April 20, 2021. <a href="https://www.fema.gov/flood-insurance">https://www.fema.gov/flood-insurance</a>.
- Federal Emergency Management Agency (FEMA). 2021b. National Flood Hazard Layer. Lowndes County, MS. 18 February 2011. Accessed April 20, 2021. <a href="https://www.fema.gov/flood-maps/tools-resources/flood-map-products/national-flood-hazard-layer">https://www.fema.gov/flood-maps/tools-resources/flood-map-products/national-flood-hazard-layer</a>.
- Federal Aviation Administration (FAA). 2013. Interim Policy, FAA Review of Solar Energy System Projects on Federally Obligated Airports. 78 FR 63276
- Flora of North America (FNA). 2020. *Carex microdonta*. <a href="www.efloras.org/florataxon.aspx?flora\_id=1&taxon\_id=242357334">www.efloras.org/florataxon.aspx?flora\_id=1&taxon\_id=242357334</a>. Accessed August 31, 2020.
- ForgeSolar. 2020. About Glint & Glare. Website. Accessed in December 2021. Available at: https://www.forgesolar.com/help/#glare.
- GEOServices, LLC. Report of Geotechnical Exploration for the Proposed Golden Triangle II Solar Project. GEOServices.
- Hilty, John. 2018. Illinois Wildflowers: Prairie Parsely (*Polytaenia nuttallii*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/pr\_parsley.html">https://www.illinoiswildflowers.info/prairie/plantx/pr\_parsley.html</a>. Accessed April 26, 2021.
- Hilty, John. 2019a. Illinois Wildflowers: Mead's Sedge (*Carex meadii*). <a href="https://www.illinoiswildflowers.info/grasses/plants/mead\_sedge.htm">https://www.illinoiswildflowers.info/grasses/plants/mead\_sedge.htm</a>. Accessed April 26, 2021.
- Hilty, John. 2020a. Illinois Wildflowers: James' Sedge (*Carex jamesii*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm">https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020b. Illinois Wildflowers: Ohio Buckeye (*Aesculus glabra*). <a href="https://www.illinoiswildflowers.info/trees/tree\_index.htm#oh\_buckeye">https://www.illinoiswildflowers.info/trees/tree\_index.htm#oh\_buckeye</a>. Accessed April 26, 2021.

- Hilty, John. 2020c. Illinois Wildflowers: Wild Hyacinth (*Camassia scilloides*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm">https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020d. Illinois Wildflowers: Kingnut Hickory (*Carya laciniosa*). <a href="https://www.illinoiswildflowers.info/trees/plants/kn\_hickory.htm">https://www.illinoiswildflowers.info/trees/plants/kn\_hickory.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020e. Illinois Wildflowers: Canada Moonseed (*Menispermum canadense*). <a href="https://www.illinoiswildflowers.info/prairie/plants/canada\_moonseed.htm">https://www.illinoiswildflowers.info/prairie/plants/canada\_moonseed.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020f. Illinois Wildflowers: Heath Aster (*Symphyotrichum ericoides*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/hth\_asterx.htm">https://www.illinoiswildflowers.info/prairie/plantx/hth\_asterx.htm</a>. Accessed August 31, 2020.
- Lowndes County (Lowndes). 2021a. Lowndes County. Tourism. http://lowndescountyms.com/about/tourism/. December 13, 2021.
- Lowndes County (Lowndes). 2021b. Lowndes County. Cities, Towns, and Communities. <a href="http://lowndescountyms.com/about/cities-towns-and-communities/">http://lowndescountyms.com/about/cities-towns-and-communities/</a>. December 13, 2021.
- MacGown, J.; Brown, R.; and Hill, J. 2006. The Black Belt Prairie in Mississippi and Alabama. Mississippi Entomological Museum. Mississippi State University. Available at: <a href="https://mississippientomologicalmuseum.org.msstate.edu/habitats/black.belt.prairie/BlackBeltPrairie.htm">https://mississippientomologicalmuseum.org.msstate.edu/habitats/black.belt.prairie/BlackBeltPrairie.htm</a>.
- Minnesota Wildflowers. 2021. A Field Guide to Flora of Minnesota: Great Plains Ladies' tresses (*Spiranthes magnicamporum*). <a href="https://www.minnesotawildflowers.info/flower/great-plains-ladies-tresses">https://www.minnesotawildflowers.info/flower/great-plains-ladies-tresses</a>. Accessed April 27, 2021.
- Minnesota Department of Natural Resources. 2021. Species Profile: *Cycleptus elongatus*. <a href="https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AF">https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AF</a> CJC04010. Accessed December 7. 2021.
- Minnesota Department of Natural Resources. 2021. Species Profile: *Crystallaria asprella*. https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AF CQC01010. Accessed December 7, 2021.
- Mississippi Department of Environmental Quality (MDEQ). 2020. Mississippi Approved 2020 Section 303(d) List of Impaired Water Bodies Final List Version 2.0. Prepared by MDEQ Surface Water Division of the Office of Pollution Control. Jackson, MS. November 2020.
- Mississippi Department of Transportation (MDOT). 2019.Traffic Volume Maps. <a href="https://mdot.ms.gov/portal/traffic volume">https://mdot.ms.gov/portal/traffic volume</a>. Accessed December 13, 2021.
- Mississippi Department of Transportation (MDOT). 2020. Five Year State Strategic Plan for Fiscal Years 2023 2027. Available at: http://www.lbo.ms.gov/misc/strategic/FY23/941-00-plan.pdf. Accessed on December 2, 2021.

- Mississippi Department of Wildlife, Fisheries, and Parks (MDWFP). 2022. Black Prairie Wildlife Management Area. Black Prairie WMA Map. <a href="https://www.mdwfp.com/media/300636/black-prairie-map.pdf">https://www.mdwfp.com/media/300636/black-prairie-map.pdf</a>. Accessed February 10, 2022.
- Mississippi Hills Natural Heritage Area. 2021. *Who We Are*. Retrieved May 2021 from: <a href="https://mississippihills.org/who-we-are/">https://mississippihills.org/who-we-are/</a>.
- Mississippi Natural Heritage Program. 2021. Wildlife of Mississippi. Mississippi Department of Wildlife, Fisheries, and Parks. <a href="http://www.mdwfp.com/museum/seek-study/natural-heritage-program/">http://www.mdwfp.com/museum/seek-study/natural-heritage-program/</a>. Accessed: May 26, 2021.
- Mississippi Natural Heritage Program. 2018. Listed Species of Mississippi. Museum of Natural Science, Mississippi Dept. of Wildlife, Fisheries, and Parks, Jackson, MS. 6 pp. 24 September 2018.
- Mississippi Museum of Natural Science. 2014. Endangered Species of Mississippi. Mississippi Department of Wildlife, Fisheries, and Parks, Mississippi Museum of Natural Science, Jackson, MS.

  https://www.mdwfp.com/media/256526/endangered species of mississippi.pdf.
- Mississippi Natural Heritage Program, 2018b. Special Plants Tracking List. Museum of Natural Science, Mississippi Department of Wildlife, Fisheries, and Parks, Jackson, MS. 12 pp.
- Missouri Botanical Garden (MBG). 2020a. Plant Finder: *Fraxinus profunda*. <a href="https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=c987">www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=c987</a>. Accessed August 31, 2020.
- Missouri Botanical Garden (MBG). 2020b. Plant Finder. *Quercus macrocarpa*. <a href="https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=a902">https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=a902</a>. Accessed August 31, 2020.
- Missouri Department of Conservation (MDC). 2020. Discover Nature: Field Guide. Celestial Lily (Nemastylis geminiflora). <a href="https://nature.mdc.mo.gov/discover-nature/field-guide/celestial-lily">https://nature.mdc.mo.gov/discover-nature/field-guide/celestial-lily</a>. Accessed August 31, 2020
- Missouri Department of Conservation (MDC). 2021. Discover Nature: Field Guide. Eastern Spotted Skunk (*Spilogale putorius*). <a href="https://nature.mdc.mo.gov/discover-nature/field-guide/eastern-spotted-skunk">https://nature.mdc.mo.gov/discover-nature/field-guide/eastern-spotted-skunk</a>. Accessed April 25, 2021.
- Montana Natural Heritage Program. 2021. Big-head Evax *Evax Prolifera*. Montanta Field Guide. <a href="http://FieldGuide.mt.gov/speciesDetail.aspx?elcode=PDAST3S050">http://FieldGuide.mt.gov/speciesDetail.aspx?elcode=PDAST3S050</a>. Accessed. April 27, 2021.
- National Oceanic and Atmospheric Administration (NOAA). 2019. National Centers for Environmental Information. State Climate Summaries (Revised in 2019). Mississippi. Available via the internet at: <a href="https://statesummaries.ncics.org/chapter/ms/">https://statesummaries.ncics.org/chapter/ms/</a>. Accessed September 2020.
- National Park Service (NPS). 2018. Series: Physiographic Provinces Coastal Plain Province. https://www.nps.gov/articles/coastalplain.htm. Accessed August 20, 2020.

- NatureServeExplorer, 2021a, Cylinder Elimia (Elimia cylindrica).
  - https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.113362/Elimia\_cylindrace a. Accessed December 7, 2021.
- NatureServeExplorer. 2021b. Flat Pigtoe (Pleurobema marshalli).
  - https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.114612/Pleurobema\_marshalli. Accessed December 7, 2021.
- NatureServeExplorer. 2021c. Freckled Darter (Percina lenticula).
  - https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.116207/Pleurobema\_taitianum. Accessed December 7, 2021.
- NatureServeExplorer. 2021d. Heavy Pigtoe (*Pleurobema taitianum*). <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.101698/Percina\_lenticula.">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.101698/Percina\_lenticula.</a> Accessed December 7, 2021.
- NatureServeExplorer. 2021e. Southern Clubshell (*Pleurobema decisum*). <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.1066298/Pleurobema\_decisum">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.1066298/Pleurobema\_decisum</a>. Accessed December 7, 2021.
- NatureServeExplorer. 2021f. Southern Combshell (*Epioblasma penita*).

  <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.109895/Epioblasma\_penit">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.109895/Epioblasma\_penit</a>

  a. Accessed December 7, 2021.
- NatureServeExplorer. 2021g. Stirrupshell (Quadrula stapes).
  - https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.107836/Theliderma\_stape s. Accessed December 7, 2021.
- NatureServeExplorer. 2021h. Striped Bass (*Morone saxatilis*). <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.104731/Morone\_saxatilis">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.104731/Morone\_saxatilis</a>. Accessed December 7, 2021.
- NatureServeExplorer. 2021h. White Heelsplitter (*Lasmigona compressa*). <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.111988/Lasmigona\_compressa">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.111988/Lasmigona\_compressa</a>. Accessed December 7, 2021.
- New York Natural Heritage Program. 2021. Online Conservation Guide for *Carya laciniosa*. Available from: <a href="https://guides.nynhp.org/big-shellbark-hickory/">https://guides.nynhp.org/big-shellbark-hickory/</a>. Accessed April 26, 2021.
- Paleontology Portal. 2020. Paleontology and Geology of Mississippi. http://paleoportal.org/index.php?globalnav=time\_space&sectionnav=state&name=Mississippi. Accessed August 13, 2020.
- Phillips, George. 2012. *The Black Prairie of Northeast Mississippi*. Mississippi Outdoors. March April 2012. Pgs 34 35, and 37.
- Shaver, Harris, House, and Kepka. 2021. Draft *Phase I Cultural Resources Survey for the Proposed Golden Triangle I Solar Farm Project, Lowndes County, Mississippi.* Prepared for MS Solar 5, LLC, and Tennessee Valley Authority. 17 September 2020. Revised December 2021.

- Sibley, David Allen. 2017. Sibley Birds East. A Field Guide to Birds of Eastern North America. Alfred A. Knofp. New York. 2017.
- Spies, T. A., Stine, P. A., Gravenmier, R. A., Long, J. W., & Reilly, M. J. (Spies et al.) 2018. Synthesis of science to inform land management within the Northwest Forest Plan area. Gen. Tech. Rep. PNW-GTR-966. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 809-849. Accessed March 6, 2020.
- Tennessee Valley Authority (TVA). 1981. *Class Review of Repetitive Actions in the 100-Year Floodplain*. Federal Register. Vol. 46, No. 76 (22845 22846). 21 April 1981.
- Tennessee Valley Authority (TVA). 2014. Final Programmatic Environmental Assessment. TVA Solar Photovoltaic Projects. Alabama, Georgia, Kentucky, Mississippi, North Carolina, Tennessee, and Virginia. September 2014. Knoxville, TN.
- Tennessee Valley Authority (TVA). 2017a. Tennessee Valley Authority Site Clearing and Grading Specifications. Revised July 2017.
- Tennessee Valley Authority (TVA). 2017b. A Guide for Environmental Protection and Best Management Practices for Tennessee Valley Authority Construction and Maintenance Activities, Revision 3. Edited by G. Behel, S. Benefield, R. Brannon, C. Buttram, G. Dalton, C. Ellis, C. Henley, T. Korth, T. Giles, A. Masters, J. Melton, R. Smith, J. Turk, T. White, and R. Wilson. Chattanooga, TN.
- Tennessee Valley Authority (TVA). 2019a. Integrated Resource Plan. <u>Volume I Final Resource Plan</u>.
- Tennessee Valley Authority (TVA). 2019b. Integrated Resource Plan. <u>Volume II Final</u> Environmental Impact Statement.
- Tennessee Valley Authority (TVA). 2021. TVA Heritage Database Results.
- Texas Parks and Wildlife. 2021. Plant Guidance by Ecoregions. Ecoregion 4 The Blackland Prairies. Available at: https://tpwd.texas.gov/huntwild/wild/wildlife\_diversity/wildscapes/ecoregions/ecoregion\_4 .phtml. Accessed May 25, 2021.
- United States. 1977. Executive Office of the President Jimmy Carter. Executive Order 11988: Floodplain Management. 24 May 1977.
- U.S. Census Bureau (USCB). 2021a. QuickFacts. Lowndes County, Mississippi. <a href="https://data.census.gov/cedsci/table?g=0400000US28\_0500000US28087&d=ACS%205-Year%20Estimates%20Data%20Profiles">https://data.census.gov/cedsci/table?g=0400000US28\_0500000US28087&d=ACS%205-Year%20Estimates%20Data%20Profiles</a>. Accessed December 13, 2021.
- U.S. Census Bureau (USCB). 2021b. American Community Survey. Obion County, Tennessee. <a href="https://data.census.gov/cedsci/table?t=Employment&g=0400000US28\_0500000US28087&d=ACS%205-Year%20Estimates%20Data%20Profiles">https://data.census.gov/cedsci/table?t=Employment&g=0400000US28\_0500000US28087&d=ACS%205-Year%20Estimates%20Data%20Profiles</a>. Accessed December 13, 2021.

- U.S. Census Bureau (USCB). 2021c. QuickFacts; Starkville City/Columbus City. Mississippi. <a href="https://www.census.gov/quickfacts/fact/table/starkvillecitymississippi,US">https://www.census.gov/quickfacts/fact/table/starkvillecitymississippi,US</a>. Accessed December 14, 2021.
- U.S. Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS).

  ND. State Soil Data Access (SDA) Hydric Soil Ratings by Map Unit.


  <a href="https://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcseprd1389479.html">https://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcseprd1389479.html</a>.

  Accessed December 14, 2021.
- U.S. Department of Agriculture. 2018. Official Soil Series Descriptions. https://soilseries.sc.egov.usda.gov/. Accessed. August 30, 2020.
- U.S. Department of Agriculture (USDA) 2021 Natural Resources Conservation Service (NRCS). 2021. Farmland Protection Policy Act. <a href="https://www.nrcs.usda.gov/wps/portal/nrcs/detail/?cid=nrcs143\_008275">https://www.nrcs.usda.gov/wps/portal/nrcs/detail/?cid=nrcs143\_008275</a>. Accessed December 14, 2021.
- U.S. Environmental Protection Agency (USEPA). 1974. Information on Levels of Environmental Noise Requisite to Protect Public Health and Welfare with an Adequate Margin of Safety. Office of Noise Abatement and Control. March 1974. 242 pps.
- U.S. Environmental Protection Agency (USEPA). 2011. Spill Prevention, Control, and Countermeasures (SPCC) Rule. Streamlined Requirements for Tier I and II Qualified Facilities. May 2011. Available at: <a href="http://www.epa.gov/emergencies">http://www.epa.gov/emergencies</a>.
- U.S. Environmental Protection Agency (USEPA). 2021b. Mississippi Nonattainment / Maintenance Status for Each County by Year for All Criteria Pollutants. <a href="https://www3.epa.gov/airquality/greenbook/anayo\_ms.html">https://www3.epa.gov/airquality/greenbook/anayo\_ms.html</a>. Accessed December 13, 2021.
- U.S. Environmental Protection Agency (USEPA). 2021c. Laws & Regulations. Summary of Executive Order 12898 Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations. <a href="https://www.epa.gov/laws-regulations/summary-executive-order-12898-federal-actions-address-environmental-justice">https://www.epa.gov/laws-regulations/summary-executive-order-12898-federal-actions-address-environmental-justice</a>. Accessed December 13, 2021.
- U.S. Environmental Protection Agency (USEPA). 2021d. 2017 National Emissions Inventory (NEI) Data. https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-data. Accessed December 14, 2021.
- U.S. Environmental Protection Agency (USEPA). 2020e. Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2018. Accessed August 27, 2020.
- U.S. Environmental Protection Agency (USEPA). 2021a. Report on the Environment. Land Use. <a href="https://www.epa.gov/report-environment/land-use">https://www.epa.gov/report-environment/land-use</a>. Accessed December 13, 2021.
- U.S. Environmental Protection Agency (USEPA). 2020a. EPA's Environmental Justice Screening and Mapping Tool. Minority Populations. ID # 280870070001. Available at ejscreen.epa.gov/mapper/.

- U.S. Environmental Protection Agency (USEPA). 2020b. EPA's Environmental Justice Screening and Mapping Tool. Low Income Populations. ID # 280870070001. Available at ejscreen.epa.gov/mapper/.
- U.S. Fish and Wildlife Service (USFWS). 2015. Midwest Region Endangered Species Northern Long-eared Bat *Myotis septentrionalis*. <a href="https://www.fws.gov/midwest/Endangered/mammals/nleb/nlebFactSheet.html">https://www.fws.gov/midwest/Endangered/mammals/nleb/nlebFactSheet.html</a>. Accessed October 21, 2020.
- U.S. Fish and Wildlife Service (USFWS). 2016. Price's potato-bean (*Apios priceana*) 5-Year Review: Summary and Evaluation. Updated on 2 September 2016. Available at https://ecos.fws.gov/docs/tess/species\_nonpublish/2363.pdf
- U.S. Fish and Wildlife Service (USFWS). 2019. Range-wide Indiana Bat Survey Guidelines. April 2019. Available via the internet at: https://www.fws.gov/arkansas-es/docs/FINAL%202019%20Range-wide%20IBat%20Survey%20Guidelines%204.10.19.pdf.
- U.S. Fish and Wildlife Service (USFWS). 2019. Midwest Region Endangered Species Price's Potato Bean (*Apios priceana*). <a href="https://www.fws.gov/midwest/endangered/plants/pricesp.html">https://www.fws.gov/midwest/endangered/plants/pricesp.html</a>. Accessed October 21, 2020.
- U.S. Fish and Wildlife Service (USFWS). 2021. IPaC Report for Golden Triangle Solar Project. Mississippi Ecological Services Field Office. Jackson, MS. April 21, 2020.
- U.S. Fish and Wildlife Service (USFWS) 2020b. Wood Stork Mycteria americana Species Account. North Florida Ecological Services Field Office.

  <a href="https://www.fws.gov/northflorida/Species-Accounts/Wood-stork-2005.htm">https://www.fws.gov/northflorida/Species-Accounts/Wood-stork-2005.htm</a>. Accessed October 21, 2020.
- U.S. Forest Service. 2020. Plant of the Week: Indian Paintbrush (*Castilleja coccinea*). <a href="https://www.fs.fed.us/wildflowers/plant-of-the-week/Castilleja-coccinea.shtml">https://www.fs.fed.us/wildflowers/plant-of-the-week/Castilleja-coccinea.shtml</a>. Accessed August 31, 2020.
- U.S. Forest Service (USFS) 2021. Fire Effects Information System (FEIS): *Quercus macrocarpa*. https://www.fs.fed.us/database/feis/plants/tree/quemac/all.html. Accessed April 26, 2021.
- U.S. Geological Survey (USGS). 1998. Ground Water Atlas of the United States, Segment 5 Arkansas, Louisiana, and Mississippi. <a href="https://pubs.usgs.gov/ha/730f/report.pdf">https://pubs.usgs.gov/ha/730f/report.pdf</a>.
  Accessed December 13, 2021.
- U.S. Geological Survey (USGS). 2014. Seismic Risk Map. <a href="https://earthquake.usgs.gov/static/lfs/nshm/conterminous/2014/2014pga2pct.pdf">https://earthquake.usgs.gov/static/lfs/nshm/conterminous/2014/2014pga2pct.pdf</a>. Accessed 8/24/2020.
- U.S. Geological Survey (USGS). 2018. Water Usage for Mississippi; Lowndes County. https://waterdata.usgs.gov/ms/nwis/water\_use?format=html\_table&rdb\_compression=file &wu area=County&wu year=2015&wu county=087&wu category=ALL&wu county n

- ms=Lowndes%2BCounty&wu\_category\_nms=--ALL%2BCategories--. Accessed December 13, 2021.
- U.S. Geological Survey (USGS). 2019a. Volcano Hazards Program. U.S. Volcanoes and Current Activity Alerts. https://volcanoes.usgs.gov/index.html. Accessed September 24, 2021.
- U.S. Geological Survey (USGS). 2019b. Earthquakes 101 The Basics. https://www.usgs.gov/natural-hazards/earthquake-hazards/science/earthquake-hazards-101-basics?qt-science\_center\_objects=0#qt-science\_center\_objects. Accessed December 13, 2021.
- U.S. Geological Survey (USGS). 2020c. Plants of Louisiana *Agalinis oligophylla*. <a href="https://warcapps.usgs.gov/PlantID/Species/Details/1985">https://warcapps.usgs.gov/PlantID/Species/Details/1985</a>. Accessed October 21, 2020.
- U.S. Global Change Research Program (USGCRP). 2018. Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II. Rev. 2020. <a href="https://nca2018.globalchange.gov/downloads/NCA4\_2018\_FullReport.pdf">https://nca2018.globalchange.gov/downloads/NCA4\_2018\_FullReport.pdf</a>. Accessed August 27, 2020.
- U.S. Water Resources Council (USWRC). 1978. Guidelines for Implementing Executive Order 11988, Floodplain Management. FR Vol. 43, No. 29—Friday, February 10, 1978. pp. 6030-6054.



# Tennessee Valley Authority Site Clearing and Grading Specifications

1. General - The project manager with the clearing and/or grading contractor(s) shall review the environmental evaluation documents for the project or proposed activity (categorical exclusion checklist, environmental assessment, or environmental impact statement) along with all clearing and construction appendices, conditions in applicable general and/or site-specific permits, the storm water pollution prevention plan, open burning or demolition notification requirements, and any Tennessee Valley Authority (TVA) commitments to property owners. The contractor shall then plan and carry out operations using techniques consistent with good engineering and storm water management practices as outlined in TVA's best management practices (BMPs) manual. The contractor will protect areas that are to be left unaffected by access or clearing work at and adjacent to all work sites. In sensitive areas and their buffers, the contractor will retain as much native ground cover and other vegetation as possible. BMPs shall be installed before general site clearing or grading, with progressive stabilization BMPs applied from the perimeter toward the interior work areas as grading is completed. Any stabilized area that must be disturbed in subsequent steps shall have temporary BMPs installed until work is completed and the area is restabilized.

If the contractor fails to use BMPs or to follow environmental expectations discussed in the prebid, prework meeting or present in contract specifications, TVA will order corrective changes and additional work, as deemed necessary in TVA's judgment, to meet the intent of environmental laws and regulations or other guidelines. Major violations or continued minor violations will result in work suspension until correction of the situation is achieved or other remedial action is taken at the contractor's expense. Penalty clauses may be invoked as appropriate.

- 2. Regulations The clearing contractor shall comply with all applicable federal, state, and local environmental and antipollution laws, regulations, and ordinances, including without limitation, all air, water, solid and hazardous waste, noise, and nuisance laws, regulations, and ordinances. He or she shall secure, or ensure that TVA has secured, all necessary permits and authorizations and made all appropriate notifications to conduct work on the acres shown on the drawings and plan and profile for the contract. The contractor's designated project manager will actively seek to prevent, control, monitor, and safely abate all commonly recognized forms of workplace and environmental pollution. Permits or authorizations and any necessary certifications of trained employees knowledgeable of environmental requirements shall be documented with copies submitted to TVA's project manager or environmental specialist before work begins. The contractor and subcontractors will be responsible for meeting all conditions specified in permits. Permit conditions shall be reviewed in prework discussions.
- 3. <u>Land and Landscape Preservation</u> The contractor shall exercise care to preserve the condition of cleared soils by avoiding as much compacting and deep scarring as possible in areas not to be developed for buildings, structures, or foundations. As soon as possible after initial disturbance of the soil and in accordance with any permit(s) or other state or local environmental regulatory requirements, cover material shall be placed to prevent erosion and sedimentation of water bodies or conveyances to

surface water or groundwater. The placement of erosion/sediment controls shall begin at the perimeter and work progressively to the interior of the site. Repeated work in an area will require establishment of a ground cover immediately after each disturbance is completed. In areas outside the clearing, borrow, fill, or use and access areas, the natural vegetation shall be protected from damage. The contractor and his or her employees and subcontractors must not deviate from delineated access routes or use areas and must enter the site(s) at designated areas that will be marked. Clearing operations shall be conducted to prevent any unnecessary destruction, scarring, or defacing of the remaining natural vegetation and adjacent surroundings in the vicinity of the work. In sensitive public or environmental areas, appropriate buffer zones shall be observed by modifying the methods of clearing or reclearing, grading, borrow, or fill so that the buffer and sensitive area are protected. Some areas may require planting native low-growing plants or grasses to meet the criteria of regulatory agencies, executive orders, or commitments to special program interests.

- 4. Streamside Management Zones The clearing and/or grading contractor(s) must leave as many rooted ground cover plants as possible in buffer zones along streams and other bodies of water or wet-weather conveyances thereto. In such streamside management zones (SMZs), tall-growing tree species (trees that would interfere with TVA's National Electrical Safety Code clearances) shall be cut, and the stumps may be treated to prevent resprouting. Low-growing trees identified by TVA as marginal electrical clearance problems may be cut and then the stump treated with growth regulators to allow low, slow-growing canopy development and active root growth. Only approved herbicides shall be used, and herbicide application shall be conducted by certified applicators from the Transmission Operations and Maintenance (TOM) organization after initial clearing and construction. Cutting of trees within SMZs must be accomplished by using either hand-held equipment or other appropriate clearing equipment, such as a feller-buncher. The method will be selected based on sitespecific conditions and topography to minimize soil disturbance and impacts to the SMZ and surrounding area. Disturbed soils in SMZs must be stabilized by appropriate methods immediately after the access or site is cleared. Stabilization must occur within the time frame specified in applicable storm water permits or regulations. Stumps within SMZs may be cut close to the ground but must not be removed or uprooted. Trees, limbs, and debris shall be prevented from falling into water bodies or immediately removed from streams, ditches, ponds, and wet areas using methods that will minimize dragging or scarring the banks or stream bottom. No debris will be left in the water or watercourse. Equipment will cross streams, ditches, or wet areas only at locations designated by TVA after the application of appropriate erosion-control BMPs and consistent with permit conditions or regulatory requirements.
- 5. Wetlands In forested wetlands, tall trees will be cut near the ground, leaving stumps and roots in place. The cambium may be treated with herbicides applied by certified applicators from the TOM organization to prevent regrowth. Understory trees that must be initially cut and removed may be allowed to grow back or may be treated with tree growth regulators selectively to slow growth and increase the reclearing cycle. The decision will be situationally made based on existing ground cover, wetland type, and tree species, since tall tree removal may "release" understory species and allow them to quickly grow to "electrical clearance problem" heights. In many circumstances, herbicides labeled for water and wetland use may be used in reclearing.

At substation, switching stations, and communications sites, wetlands are avoided unless there is no feasible alternative.

- 6. Sensitive Area Preservation If prehistoric or historic artifacts or features that might be of archaeological or historical significance are discovered during clearing, grading, borrow, or fill operations, the activity shall immediately cease within a 100-foot radius, and a TVA project manager, an environmental specialist, and the TVA Cultural Resources program manager shall be notified. The site shall be protected and left as found until a determination about the resources, their significance, and site treatment is made by TVA's Cultural Resources Program. Work may continue beyond the finding zone and the 100-foot radius beyond its perimeter.
- 7. Water Quality Control The contractor's clearing, grading, borrow and fill, and/or disposal activities shall be performed using BMPs that will prevent erosion and entrance of spillage, contaminants, debris, and other pollutants or objectionable materials into drainageways, surface waters, or groundwater. Special care shall be exercised in refueling equipment to prevent spills. Fueling areas shall be remote from any sinkhole, crevice, stream, or other water body. Open burning debris shall be kept away from streams and ditches and shall be incorporated into the soil. Only materials allowed to be burned under an open burning permit may be incorporated into the soil.

The clearing and grading contractor(s) and subcontractors will erect and (when TVA or contract construction personnel are unable) maintain BMPs, such as silt fences, on steep slopes and adjacent to any steam, wetland, or other water body. BMPs will be inspected by the TVA field engineer or other designated TVA or contractor personnel routinely and at least as frequently as required by the permit or good management practices and during periods of high runoff; any necessary repairs will be made as soon as practicable. BMP runoff sampling will be conducted in accordance with permit requirements. Records of all inspections and sampling will be maintained on site, and copies of inspection forms and sampling results will be forwarded to the TVA environmental specialist.

8. Turbidity and Blocking of Streams - If temporary clearing, grading, borrow, or fill activities must interrupt natural drainage, appropriate drainage facilities and erosion/sediment controls shall be provided to avoid erosion and siltation of streams and other water bodies or water conveyances. In Tennessee, conditions of an Aquatic Resource Alteration Permit shall be met. Turbidity levels in receiving waters or at storm water discharge points shall be monitored, documented, and reported if required by the applicable permit. Erosion and sediment control measures such as silt fences, water bars, and sediment traps shall be installed as soon as practicable after initial access, site, borrow, fill, or right-of-way disturbance and after sequential disturbance of stabilized areas due to stepwise construction requirement in accordance with applicable permit or regulatory requirements.

On rights-of-way, mechanized equipment shall not be operated in flowing water except when approved and then only to construct necessary stream crossings under direct quidance of TVA.

Construction of stream fords or other crossings will only be permitted at approved locations and to current TVA design or construction access road standards. At any construction site, material shall not be deposited in watercourses or within stream bank

areas where it could be washed away by high stream flows. Any clearing debris that enters streams or other water bodies shall be removed immediately. Appropriate U.S. Army Corps of Engineers and state permits shall be obtained for stream or wetland crossings.

- 9. <u>Air Quality Control</u> The clearing or grading contractor shall take appropriate actions to limit the amount of air emissions created by clearing and disposal operations to be well within the limits of clearing or burning permits and/or forestry or local fire department requirements. All operations must be conducted in a manner that prevents nuisance conditions or damage to adjacent land, crops, dwellings, highways, or people. If building renovation or demolition is involved, the required air quality organization shall be notified the minimum 10 days in advance, and if the start date is delayed, renotified to start the clock again.
- 10. <u>Dust and Mud Control</u> Clearing, grading, borrow, fill, or transport activities shall be conducted in a manner that minimizes the creation of fugitive dust. This may require limitations as to type of equipment, allowable speeds, and routes utilized. Control measures such as water, gravel, etc., or similar measures may be used subject to TVA approval. On new construction sites and easements, the last 100 feet before an access road approaches a county road or highway shall be graveled to prevent transfer of mud onto the public road.
- 11. <u>Burning</u> The contractor shall obtain applicable permits and approvals to conduct controlled burning. The contractor will comply with all provisions of the permit, notification or authorization including burning site locations, controlled draft, burning hours, and such other conditions as stipulated. If weather conditions such as wind speed or wind direction change rapidly, the contractor's burning operation may be temporarily stopped by TVA's field engineer. The debris to be burned shall be kept as clean and dry as possible and stacked and burned in a manner that produces the minimum amount of smoke. Residue from burning will be disposed of according to permit stipulations. No fuel starters or enhancements other than kerosene will be allowed.
- 12. <u>Smoke and Odors</u> The contractor will properly store and handle combustible and volatile materials that could create objectionable smoke, odor, or fumes. The contractor shall not burn oil or refuse that includes trash, rags, tires, plastics, or other manufactured debris.
- 13. Vehicle Exhaust Emissions The contractor shall maintain and operate equipment in a manner that limits vehicle exhaust emissions. Equipment and vehicles will be kept within the manufacturer's recommended limits and tolerances. Excessive exhaust gases will be eliminated, and inefficient operating procedures will be revised or halted until corrective repairs or adjustments are made.
- 14. <u>Vehicle Servicing</u> Routine maintenance of vehicles will not be performed on the site, right-of-way, or access route. However, if emergency or "have to" situations arise, minimal/temporary maintenance to vehicles will occur in order to mobilize the vehicle to an off-site maintenance shop. Some heavy equipment may have to be serviced on the right-of-way, site, or access route, except in designated sensitive areas. The clearing, grading, borrow, or fill contractor will properly maintain these vehicles with approved spill protection controls and countermeasures. If emergency maintenance in a

- sensitive or questionable area arises, the Area Environmental Program Administration or project manager will be consulted. All wastes and used oils will be properly recovered, handled, and disposed/recycled. Equipment shall not be temporarily stored in stream floodplains, whether overnight or on weekends or holidays.
- 15. <u>Noise Control</u> The contractor shall take steps to avoid the creation of excessive sound levels for employees, the public, or the site and adjacent property owners. Concentration of individual noisy pieces as well as the hours and locations of operation should be considered.
- 16. <u>Noise Suppression</u> All internal combustion engines shall be properly equipped with mufflers. The equipment and mufflers shall be maintained at peak operating efficiency.
- 17. <u>Sanitation</u> A designated representative of TVA or the clearing, grading, borrow, fill, or construction contractor shall contract a sanitary contractor who will provide sanitary chemical toilets convenient to all principal points of operation for every working party and at each construction step. The facilities shall comply with applicable federal, state, or local health laws and regulations. They shall not be located closer than 100 feet to any stream or tributary or to any wetland. The facilities shall be required to have proper servicing and maintenance, and the waste disposal contractor shall verify in writing that the waste disposal will be in state-approved facilities. Employees shall be notified of sanitation regulations and shall be required to use the toilet facilities.
- 18. <u>Refuse Disposal</u> The clearing, grading, borrow, fill, or construction contractor and subcontractor(s) shall be responsible for daily cleanup and proper labeling, storage, and disposal of all refuse and debris on the site produced by his or her operations and employees. Facilities that meet applicable regulations and guidelines for refuse collection will be required. Only approved transport, storage, and disposal areas shall be used. Records of waste generation shall be maintained for a site and shall be provided to the project manager and environmental specialist assigned to the project.
- 19. <u>Brush and Timber Disposal (Initial Clearing)</u> For initial clearing, trees are commonly part of the contractor's contract to remove as they wish. Trees may be removed from the site for lumber or pulpwood, or they may be chipped or stacked and burned. All such activities must be coordinated with the TVA field engineer and the open burning permits; notifications and regulatory requirements must be met. On rights-of-way, trees may be cut and left in place only in areas specified by TVA and approved by appropriate regulatory agencies. These areas may include sensitive wetlands or SMZs where tree removal would cause excessive ground disturbance or in very rugged terrain where windrowed trees are used as sediment barriers along the edge of the right-of-way, site, or access.

Trees that have been cut may not be left on a substation, switching station, or communications site.


20. <u>Restoration of Site</u> - All disturbed areas, with the exception of farmland under cultivation and any other areas as may be designated by TVA's specifications, shall be stabilized in the following manner unless the property owner and TVA's engineer specify a different method:

- A. The subsoil shall be loosened to a minimum depth of 6 inches if possible and worked to remove unnatural ridges and depressions.
- B. If needed, appropriate soil amendments will be added.
- C. All disturbed areas will initially be seeded with a temporary ground cover such as winter wheat, rye, or millet, depending on the season. Perennials may also be planted during initial seeding if proper growing conditions exist. Final restoration and final seeding will be performed as line, site, or communications facilities construction is completed. Final seeding will consist of permanent perennial grasses such as those outlined in TVA's A Guide for Environmental Protection and Best Management Practices for Tennessee Valley Authority Construction and Maintenance Activities (TVA, 2017). Exceptions would include those areas designated as native grass planting areas. Initial and final restoration will be performed by the clearing contractor with emphasis on using landscaping materials provided in guidelines for low maintenance native vegetation use.
- D. TVA holds the option, depending upon the time of year and weather condition, to delay or withdraw the requirement of seeding until more favorable planting conditions are certain. In the meantime, other stabilization techniques must be applied.
- E. Vegetation designated by the Federal Invasive Species Council must be eliminated at the work site, and equipment being transported from location to location must be inspected to ensure removal and destruction of live material.

### References

Tennessee Valley Authority. 2017. A Guide for Environmental Protection and Best Management Practices for Tennessee Valley Authority Construction and Maintenance Activities, Revision 3. Edited by G. Behel, S. Benefield, R. Brannon, C. Buttram, G. Dalton, C. Ellis, C. Henley, T. Korth, T. Giles, A. Masters, J. Melton, R. Smith, J.Turk, T. White, R. Wilson. Chattanooga, TN.: Retrieved from <a href="https://www.tva.com/Energy/Transmission-System/Transmission-System-Projects">https://www.tva.com/Energy/Transmission-System/Transmission-System-Projects</a> (n.d.).

Revision July 2017





# Golden Triangle II Solar Project Wetland and Waterbody Delineation Report



MS Solar 6, LLC

**Golden Triangle Solar Project** 

Revision 2.0 10/28/2021

# Golden Triangle II Solar Project Wetland and Waterbody Delineation Report

prepared for

MS Solar 6, LLC
Golden Triangle Solar Project
Artesia, MS

Revision 2.0 10/28/2021

prepared by

Burns & McDonnell Engineering Company, Inc. Atlanta, Georgia

COPYRIGHT © 2021 BURNS & McDONNELL ENGINEERING COMPANY, INC.

# **EXECUTIVE SUMMARY**

MS Solar 6, LLC, is evaluating a site in Lowndes County, Mississippi (MS) for potential development of a 150 megawatt (MW) solar energy facility (Golden Triangle Solar II Project or the Project). Burns & McDonnell Engineering Company, Inc. (Burns & McDonnell) was contracted by MS Solar 6, LLC, to provide wetland delineation services for the proposed Project, specifically potential parcels upon which solar arrays and other appurtenant facility components may be installed, in Lowndes County, MS (**Figure 1, Appendix A**). The Survey Area consists of approximately 1,525 acres of land that is predominantly used for pasture, hay production, and agriculture with fragmented forested areas throughout. The Project is east of the US Hwy Alt 45, south and adjacent to Gilmer Wilburn Road, North of Swedenburg Road, and west of Hardy Billups Road in Artesia, MS, which is approximately 8 miles east of Starkville, MS and 10 miles west of Columbus, MS. The Project was surveyed for ecological resources from March 8 through March 13, 2021.

The Survey Area is composed of four distinct vegetative/land use communities: active agriculture, bottomland hardwood forest, upland forest, and pasture. A total of 100 aquatic resources were identified within the Survey Area for the Project including 58 ephemeral, intermittent, or perennial streams, 20 ponds, and 22 wetlands. These features, if impacted by the Project, may be subject to regulatory requirements under Section 404 of the Clean Water Act (CWA).

MS Solar 6, LLC i Burns & McDonnell

# **TABLE OF CONTENTS**

EXEC	UTIVI	E SUMMARY <u>Page No.</u>
1.0	INTR	ODUCTION1-1
2.0	MET	HODOLOGY2-1
	2.1 2.2 2.3	Existing Data Review
3.0	RES	ULTS AND DISCUSSION3-1
	3.1	Existing Data Review
	3.2	3.1.3Previous Delineations and Jurisdictional Determinations3-2Wetland Delineation Survey3-23.2.1Vegetation and Land Use Communities3-33.2.2Waters of the US3-5
	3.3	Conclusions 3-10
4.0	REF	ERENCES4-1
APPE APPE APPE	NDIX NDIX NDIX	A - FIGURES B - ANTECEDENT PRECIPITATION TOOL ANALYSIS C - REPRESENTATIVE PHOTOGRAPHS D - REPRESENTATIVE WETLAND DETERMINATION DATA FORMS E - TVA RAPID ASSESSMENT METHOD (RAM) FIELD FORM
		LIST OF TABLES
Table :		Waterbodies within the Survey Area

## LIST OF ABBREVIATIONS

Abbreviation Term/Phrase/Name

1987 Manual 1987 Corps of Engineers Wetlands Delineation Manual

° F Degrees Fahrenheit

APT Antecedent Precipitation Tool

Burns & McDonnell Engineering Company, Inc.

CWA Clean Water Act

FEMA Federal Emergency Management Agency

GPS Global Positioning Systems

HUC Hydrologic Unit Code

NAIP National Agriculture Imagery Program

NFHL National Flood Hazard Layer

NHD National Hydrography Dataset

NRCS Natural Resources Conservation Service

NWI National Wetland Inventory

PAB Palustrine Aquatic Bed

PEM Palustrine Emergent Wetland

PFO Palustrine Forested Wetland

PSS Palustrine Scrub-shrub Wetland

PUB Palustrine Unconsolidated Bottom - Pond

Regional Supplement Regional Supplement to the Corps of Engineers Wetland Delineation

Manual: Atlantic and Gulf Coast Plain Region – Version 2.0

SSURGO Soil Survey Geographic

Survey Area The approximately 1,525 acres that were evaluated during field surveys

USACE U.S. Army Corps of Engineers

USDA U.S. Department of Agriculture

USEPA U.S. Environmental Protection Agency

USFWS U.S. Fish and Wildlife Service

USGS U.S. Geological Survey

WETS NRCS's Climate Analysis for Wetlands Table

MS Solar 6, LLC i Burns & McDonnell

## 1.0 INTRODUCTION

MS Solar 6, LLC plans to construct a new utility scale solar farm and associated infrastructure on five parcels in Lowndes County, Mississippi. The Project is east of the US Hwy Alt 45, south and adjacent to Gilmer Wilburn Road, North of Swedenburg Road, and west of Hardy Billups Road in Artesia, MS, which is approximately three miles southeast of Artesia, eight miles east of Starkville, and 10 miles west of Columbus, MS. The Project has the potential to impact wetlands or other water bodies that may be under the jurisdiction of the US Army Corps of Engineers (USACE) as designated by Section 404 of the Clean Water Act (CWA). Burns and McDonnell conducted a wetland delineation for the Project to evaluate the presence of wetlands and other water bodies, including streams, drainages, and ponds. The delineation was conducted within the five parcels being considered for the proposed Project (Survey Area) as identified by MS Solar 6, LLC. The Survey Area included in the wetland delineation totaled approximately 1,525 acres.

#### 2.0 METHODOLOGY

The following discussions summarize the methods used for the review of existing data and the wetland delineation.

# 2.1 Existing Data Review

Burns & McDonnell reviewed available background information for the proposed Project prior to conducting a site visit. This available background information included:

- U.S. Geological Survey (USGS) 7.5-minute topographic maps (Artesia and Bent Oak, MS quadrangles),
- USGS National Hydrography Dataset (NHD),
- U.S. Fish & Wildlife Service (USFWS) National Wetland Inventory (NWI) maps,
- National Agriculture Imagery Program (NAIP) aerial photography (2021),
- Federal Emergency Management Agency (FEMA) 2021 National Flood Hazard Layer (NFHL),
- U.S. Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS) 2021
   Soil Survey Geographic (SSURGO) digital data for Lowndes County, Mississippi,
- Any wetland delineations performed in adjacent or abutting areas within the last five years, and
- NRCS Climate Analysis for Wetlands Table (WETS Table).

**Figure sets 2 and 3 in Appendix A** depict this data. A summary of historic and recent rainfall data is provided in Section 3, below.

# 2.2 Environmental Field Survey

A wetland delineation was completed March 8 - March 13, 2021. The delineation was conducted in accordance with the 1987 Corps of Engineers Wetlands Delineation Manual (1987 Manual) and the 2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Atlantic and Gulf Coast Plain Region – Version 2.0 (Regional Supplement).

Systematic quadrants were used as part of the field delineation methodology in conjunction with desktop analyzed target areas due to characteristics of the Study Area. Wetland data points were established as verification of the known or suspected wetland areas or other Waters of the U.S. (WOTUS) and to confirm the primary non-wetland habitat areas. All wetland points and potential WOTUS were identified on Project mapping using Global Positioning Systems (GPS) with submeter accuracy.

Wetland (and non-wetland) sample points were established using the 1987 Manual and Regional Supplement based on observations of vegetation, topographic and hydrologic features, transitions in the field, and soils. Soil samples were taken using either a soil probe, hand auger, or shovel to a minimum depth of 12 inches. Munsell Color Charts were used to reference soil matrix, mottle and chroma. Observations were documented on the USACE Atlantic and Gulf Coastal Plain Regional Wetland Determination Data Forms from the Regional Supplement (**Appendix D**) as well as through digital photographs representative of each area (**Appendix B**).

# 2.3 Wetland and Waterbody Classifications

Under typical conditions, wetlands are defined by three key criteria: vegetation, hydrology, and soils. Wetlands in the Survey Area that are anticipated to be jurisdictional are considered part of a palustrine wetland system within either a forested class (PFO), scrub-shrub class (PSS), unconsolidated bottom (PUB), aquatic bed (PAB) and/or an emergent class (PEM).

Palustrine forested wetlands (PFO) consist of an overstory dominated by deciduous broad-leaved tree species such as maples, elms, and oaks, and an assortment of herbaceous plants and vines in the understory. The dominant tree and sapling species found in the Survey Area included water oaks (Quercus nigra), green ash (Fraxinus pennsylvanica), sugarberry (Celtis laevigata), osage orange (Maclura pomifera), black willow (Salix nigra), water locust (Gleditsia aquatica), red maple (Acer rubrum), American sycamore (Platanus occidentalis), American elm (Ulmus americana), Chinese privet (Ligustrum sinense), and willow oak (Quercus phellos). Common vegetation comprising the understory includes poison ivy (Toxicodendron radicans), butterweed (Packera glabella), roundleaf greenbrier (Smilax rotundifolia), Virginia creeper (Parthenocissus quinquefolia), yellow nutsedge (Cyperus esculentus), and soft rush (Juncus effusus). Primary hydrological indicators observed in PFO wetlands included surface water, saturation in the upper 12 inches, water marks and/or water-stained leaves, drainage patterns, oxidized rhizospheres on living roots, and geomorphic position.

Palustrine scrub-shrub wetlands (PSS). There were two PSS wetlands within the Survey Area. PSS were generally identified within the southern portion of the Survey Area near the lower elevations and relief associated with the floodplain. Typical, dominant scrub shrub species encountered during surveys included green ash, black willow (*Salix nigra*), butterweed, bulbous bittercress (*Cardamine bulbosa*), and soft rush. Primary hydrological indicators observed in PSS wetlands included surface water, saturation in the upper 12 inches, water marks and/or water-stained leaves, and sediment deposits.

Palustrine emergent wetlands (PEM). Species commonly found in the palustrine emergent wetlands within the Survey Area included shallow sedge (*Carex lurida*), black willow, alligator weed (*Alternanthera philoxeroides*), rough cockleburr (*Xanthium strumarium*), butterweed, , annual ragweed (*Ambrosia artemisiifolia*), great ragweed (*Ambrosia trifida*) ,woolgrass (*Scirpus cyperinus*), curly dock (*Rumex crispus*), blackberries (*Rubus* spp.), spike rush (*Eleocharis palustris*), yellow nutsedge, and soft rush. Primary hydrological indicators observed in PEM wetlands include surface water, saturation in the upper 12 inches, algal mats or crust, drainage patterns, geomorphic position, and a positive FAC-neutral test.

**Perennial streams** are typically characterized by a well-defined channel that contains waterflow all year.

**Intermittent streams** also have a well-defined channel, with little to no vegetation through the middle; however, these streams may not flow year-round. Under typical conditions, the intermittent streams within the Survey Area likely only contain water flow in the late winter and early spring when ground water levels are higher.

**Ephemeral streams** exist within open agricultural fields, pastures, and forested areas throughout the Survey Area. These areas have been used for row crops, hay production, and livestock grazing, and over the past century stormwater drainage improvements have been made. Some ephemeral features mapped are manmade drainages, intended to aid in agricultural production, and have altered the historical drainage patterns that would have existed prior to farming in this region.

# 3.0 RESULTS AND DISCUSSION

The following sections describe the results of the desktop data review and the completed wetland delineation survey.

# 3.1 Existing Data Review

The initial phase of this study included a comprehensive review and assessment of all available information related to the Survey Area and adjacent properties.

# 3.1.1 Online Mapping and Databases

The existing USGS topographic maps were reviewed to familiarize Burns & McDonnell wetland personnel with the topography and potential locations of wetlands and other water bodies (**Figure Set 2**, **Appendix A**). The USGS topographic maps indicate the Survey Area crosses open fields and forested areas with gentle slopes. Elevation within the Survey Area ranges from approximately 217 feet to 357 feet above mean sea level (asl)

A review of FEMA's NFHL indicates the southern portion of the Survey Area is within a 100-year floodplain (Figure Set 2, Appendix A). The floodplain is predominately associated with the north branch of Magowah Creek, which flows along the southern boundary of the Survey Area.

The NWI data identified a variety of wetland types, primarily associated with riparian areas along perennial streams. The NHD data identified perennial and intermittent streams (Figure Set 2, Appendix A). Basing the presence or absence of wetlands on only NWI maps cannot be assumed as an accurate assessment of potentially occurring jurisdictional wetlands. Wetland identification criteria differ between the USFWS and the USACE. As a result, wetlands shown on an NWI map may not be under the jurisdiction of the USACE; likewise, all USACE-jurisdictional wetlands are not always identified on NWI maps. Therefore, a detailed field survey was conducted to identify any wetlands or other water bodies that may be present.

The USDA NRCS SSURGO digital data indicate that portions of more than 12 soil map units are within the Survey Area (**Figure Set 3, Appendix A**). Approximately 605 acres of the mapped soils (or approximately 40 percent of the Survey Area) are on the hydric soil lists (Map Unit Soil Key, Appendix A). The Survey Area is located within MLRA-135A (Alabama and Mississippi Blackland Prairie). Therefore, upland soils will exhibit a low chroma matrix, which is characteristic of the native parent material and is not necessarily the result of extensive soil saturation.

The 2021 NAIP aerial photography indicates the Survey Area consists largely of maintained pastureland, actively cultivated agriculture, and fragmented forested areas. (Figure Set 4, Appendix A).

## 3.1.2 Climate Information

The Antecedent Precipitation Tool (APT) – developed by the USACE – was used to assess the climate conditions in the months leading up to and during the March 2021 pedestrian survey. The APT provides a standardized method for evaluating precipitation conditions relative to a climate normal, the presence of drought conditions, and the approximate dates of wet and dry seasons for a given location.

At the time of the March 2021 pedestrian survey, conditions on site appeared to be dry, with little to no rainfall immediately before field mobilization. The APT results (**Appendix C**) confirm that conditions on site were normal, and the overall drought index (PDSI) was listed as mild wetness.

Weather conditions during the field surveys varied from clear to overcast and temperatures ranged from a low of 31 degrees Fahrenheit (°F) to a high of 80°F.

#### 3.1.3 Previous Delineations and Jurisdictional Determinations

Burns & McDonnell was provided with a copy of the 2016 Wetland Delineation and Determination Report for the GTRA Mega Site in Lowndes County (Headwaters, Inc., 2016). The Mega Site, which is the future location for an industrial and technology park, was first surveyed in 2009/2010 and is approximately four to five miles north northeast from the Survey Area. Similar results were obtained during the 2015/2016 study. The Mega Site Report provides a more baseline look at a site conditions during a typical year in terms of normal rainfall levels. Additionally, in 2020, a wetland delineation was conducted for a proposed solar facility on approximately 4000 acres immediately north of the Survey Area. Results for that wetland delineation were similar, however, record rainfall for the year, in addition to several heavy rain events, were encountered during that delineation, resulting in wetter than normal conditions in 2020 (Burns & McDonnell 2020).

# 3.2 Wetland Delineation Survey

From March 8 through March 13, 2021, a team of Burns and McDonnell wetland scientists conducted a wetland delineation over 1,525 acres of land that is being considered for the siting of the new Golden Triangle II Solar Project. The field surveys documented average hydrologic conditions, and evaluated multiple parameters in addition to hydrology, including hydric soils, hydrophytic vegetation, and a variety of stream morphological characteristics in making resource determinations. Sample plots were established at multiple locations, and wetland determination data forms from the Regional Supplement were completed to characterize the Survey Area (**Appendix D**). Vegetation, soil conditions, and hydrologic

indicators were recorded at each of these sample plots. Locations of sample plots and other identified features were surveyed using a sub-meter accurate GPS unit. Natural color photographs depicting water bodies, streams, and representative field conditions were taken and are included in **Appendix B**. Additional representative photographs were taken during the wetland delineation to document onsite conditions where sample plots were not collected. These additional photographs are not included in Appendix B but can be provided upon request. Land cover and delineated wetlands from field surveys are discussed in detail below.

# 3.2.1 Vegetation and Land Use Communities

The Project is in the US Environmental Protection Agency (USEPA) Blackland Prairie Ecoregion (Level 4) and is in the Middle Tombigbee River [Hydrologic Unit Code (HUC) 03160106] and Tibbee Creek (HUC 03160104) watersheds.

Four distinct vegetative cover/land use communities were observed in the Survey Area, including active agriculture and pasture, bottomland hardwood forest, upland forest, and right-of-way. Descriptions of these communities are provided below. Soils within the Survey Area, as identified in the USDA NRCS geospatial data for Lowndes County, included Brooksville silty clay, Catalpa silty clay, Demopolis-Binnsville complex, Leeper silty clay, Okolona silty clay, Sumter silty clay loam, Sumter-Demopolis-Chalk outcrop complex, and Vaiden silty clay. Catalpa silty clay, Demopolis-Binnsville complex, Leeper silty clay, Sumter silty clay loam, Sumter-Demopolis-Chalk outcrop complex, and Vaiden silty clay are identified by the NRCS as having hydric components. (Figure Set 3, Appendix A).

#### Active Agriculture and Pasture

Active agricultural and pasture is the primary land use community found in the Survey Area and composed approximately 73.64 percent (1,123 acres) of the Survey Area. Areas identified as active agriculture include cattle pasture, hay production, and row crop fields. Vegetation in these communities is maintained in an early successional state due to crop growth/harvesting, and regular management/mowing. Corn is planted in late spring and cover the row crop fields on approximately 125 acres. Vegetation observed in pastures consists of primarily tall fescue grass (*Festuca arundinacea*), Johnson grass (*Sorghum halepense*), annual bluegrass (*Poa annua*), couch grass (*Elymus repens*), bermuda (*Cynodon dactylon*), annual ragweed, cheatgrass (*Bromus tectorum*), perennial ryegrass (*Lolium perenne*), green foxtail (*Setaria viridis*), butterweed, bulbous bittercress (*Cardamine bulbosa*), soft rush (*Juncus effusus*), Cherokee sedge (*Carex cherokeensis*), fox sedge (*Carex vulpinoidea*), red sorrel (*Rumex acetosella*), curly dock (*Rumex crispus*), prairie fleabane (*Erigeron strigosus*), jimsonweed (*Datura*)

stramonium), Carolina horsenettle (Solanum carolinense), bull thistle (Cirsum vulgare), wild violet (Viola sororia), field garlic (Allium vineale), and sensitive partridge pea (Chamaecrista nictitans).

The soil matrix color throughout these areas ranged from a 2/1 (black) to a 5/4 (yellowish brown) on the 10YR chart; however, little to no redox concentrations or depletions were present. The Survey Area is located within MLRA-135A (Alabama and Mississippi Blackland Prairie). Therefore, upland soils will exhibit a low chroma matrix, which is characteristic of the native parent material and is not caused by wetness of the soil. Since redox concentrations were not present within at least 4-inches of the first 12-inches (F6) or 6-inches of redox within the first 10-inches of soil samples collected, the soils in the active agricultural fields and pasturelands do not meet the requirements for hydric soil indicators.

#### **Bottomland Hardwood Forest**

Approximately 17.6 percent (268 acres) of the Survey Area is composed of forested areas. Bottomland hardwood forest communities in the Survey Area are composed of a canopy age ranging from approximately 20 to 70+ years old. Dominant vegetation observed consisted of water hickory (*Carya aquatica*), willow oak, cherrybark oak (*Quercus pagoda*), swamp chestnut oak (*Quercus michauxii*), silky dogwood (*Cornus amomum*), osage orange, green ash, eastern red cedar (*Juniperus virginiana*), water locust, shagbark hickory (*Carya ovata*), box elder (*Acer negundo*), red maple, American sycamore (*Platanus occidentalis*), sugarberry, Cherokee sedge (*Carex cherokeensis*), wild petunia (*Ruellia humilis*), tall fescue, poison ivy, greenbrier, Virginia spiderwort (*Tradescantia virginiana*), Virginia creeper, prairie ironweed (*Vernonia fasciculata*), hairy buttercup (*Ranunculus sardous*), and resurrection fern (*Pleopeltis polypodioides*),

# **Upland Forest**

Upland forest communities in the Survey Area and are composed of a canopy age ranging from approximately 20 to 70 years old. Dominant vegetation observed consisted of white oak (*Quercus alba*), southern red oak (*Quercus falcata*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), mockernut hickory (*Carya tomentosa*), shagbark hickory, pignut hickory (*Carya glabra*), loblolly pine (*Pinus taeda*), eastern red cedar, American elm (*Ulmus americana*), honey locust (*Gleditsia triacanthos*), osage orange, Chinese privet, Devil's walkingstick (*Aralia spinosa*), Christmas fern (*Polystichum acrostichoides*), Japanese honeysuckle (*Lonicera japonica*), multiple greenbrier species (*Smilax spp.*), muscadine (*Vitus rotundifolia.*), Virginia creeper, blackberry (*Rubus spp.*), wooly panic grass (*Dichanthelium acuminatum*), and wild violet.

# 3.2.2 Waters of the US

Jurisdictional water(s) of the US are defined by 33 CFR Part 328.3 (b) and are protected by Section 404 of the CWA (33 USC 1344). An assessment of potential water(s) of the US that were identified within the Survey Area was performed using USGS topographic maps, NWI maps, county soil survey maps, and then refined during field investigations. Wetland locations were determined using the 1987 Corps of Engineers Wetlands Delineation Manual and the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: 2010 Atlantic and Gulf Coastal Plain. The multi-parameter approach requires that areas be considered jurisdictional wetlands if they exhibit evidence of all three wetland classification parameters. During the field surveys, a total of 59 potential ephemeral, intermittent, or perennial streams, 20 open waters, and 19 wetlands were identified within the Survey Area of the Project. Additionally, the USACE Mobile district has not made any official jurisdictional determinations on aquatic resources within the Survey Area for this project at this time (Table 3-2 and Figure Set 4, Appendix A).

#### **Streams**

Fifty-nine stream channels, consisting of three stream types (perennial, intermittent, and ephemeral) and totaling 50,858 linear feet were delineated within the Survey Area (**Photographs, Appendix C**). The different stream types are summarized below.

Thirty-one ephemeral stream channels, totaling 12,195 feet were delineated in the Survey Area. Ephemeral streams were characterized by a defined bed and bank, with some water but no flow during the survey, indicating that these streams largely carry water only during and after precipitation events. Ephemeral streams ranged from approximately 1 to 2 feet in width at the ordinary high-water mark (OHWM) (if present) and with bank heights ranging from 0.25 to 2 feet. At the time of the delineation, water was not observed flowing in features identified as ephemeral streams. The substrates of the ephemeral streams were comprised of vegetation and silt with limited gravel. These streams were in upland fields and wooded riparian areas.

Twenty-two intermittent stream channels, totaling 13,138 feet were delineated in the Survey Area. Intermittent streams were characterized by the presence of a limited volume of flow at the time of the site visit. This is a likely indicator that the stream is partially influenced by groundwater, but it may not flow during dry periods. Intermittent streams were one to four feet in width at the OHWM with bank heights ranging from one to three feet. At the time of the delineation, water was observed at a depth of 0.25 inches to six inches. The substrates of intermittent streams were comprised of silt with limited gravel. These streams flowed through upland fields, agricultural fields, and wooded riparian areas.

Six perennial streams, totaling 25,525 feet were delineated within the Survey Area. Perennial streams were characterized by the presence of a substantial volume of flow at the time of the site visit as well as secondary characteristics such as observance of fish and rooted aquatic fauna, indicating that water flows year-round. Perennial streams were approximately 6 to 25 feet in width at the OHWM with bank heights ranging from 3 to 25 feet. At the time of the delineation, the depth of water observed was 0.5 to 2.5 feet. The substrates of the perennial streams were likely comprised of silt, gravel, and cobble although this could not be confirmed at all streams due to turbidity. Perennial streams flowed through upland fields, agricultural fields, and wooded riparian areas.

Common riparian vegetation associated with streams in the Survey Area consisted of Osage orange, eastern red cedar, sugarberry, tall fescue, deadnettle henbit, white clover, Japanese honeysuckle, and curly dock.

There are no streams within the Survey Area listed on the final 2018 version of Mississippi's 303(d) list as "non-supporting" streams. Further, there are no streams located within one linear mile upstream of, and within the same watershed as a "non-supporting" biota impaired system.

Table 3-1: Waterbodies within the Survey Area

Waterbody ID	Waterbody Type	Length of Stream (feet) in Survey Area	Latitude (N°)	Longitude (W°)	Anticipated Jurisdictional Status <sup>a</sup>
S-01	Intermittent	1,113	33.398430	-88.615520	Yes
S-02	Intermittent	156	33.398198	-88.616934	Yes
S-03	Intermittent	151	33.398200	-88.617606	Yes
S-04	Ephemeral	776	33.397542	-88.616120	Yes
S-05	Perennial	4,243	33.392894	-88.595945	Yes
S-05	Intermittent	695	33.398762	-88.600374	Yes
S-06	Intermittent	135	33.395984	-88.598576	Yes
S-07	Intermittent	606	33.395746	-88.598127	Yes
S-08	Ephemeral	54	33.393545	-88.596627	Yes
S-09	Intermittent	429	33.391849	-88.594758	Yes
S-10	Ephemeral	36	33.391766	-88.594767	Yes
S-11	Intermittent	523	33.391423	-88.594635	Yes
S-12	Intermittent	284	33.390741	-88.594455	Yes
S-13	Ephemeral	700	33.400203	-88.598452	Yes
S-14	Ephemeral	1,675	33.396808	-88.593018	Yes
S-14	Intermittent	550	33.394039	-88.592681	Yes
S-15	Ephemeral	217	33.397628	-88.607215	Yes

Waterbody ID	Waterbody Type	Length of Stream (feet) in Survey Area	Latitude (N°)	Longitude (W°)	Anticipated Jurisdictional Status <sup>a</sup>
S-16	Perennial	11,766	33.386522	-88.601192	Yes
S-16a	Ephemeral	54	33.383941	-88.605115	Yes
S-17	Ephemeral	582	33.390062	-88.601587	Yes
S-18	Intermittent	303	33.384060	-88.602421	Yes
S-19	Perennial	664	33.386621	-88.608580	Yes
S-20	Ephemeral	122	33.383512	-88.608229	Yes
S-21	Intermittent	378	33.385406	-88.607775	Yes
S-22	Ephemeral	98	33.384430	-88.601223	Yes
S-23	Perennial	2,790	33.382229	-88.611714	Yes
S-24	Ephemeral	74	33.383015	-88.606066	Yes
S-25	Intermittent	267	33.382520	-88.606954	Yes
S-26	Ephemeral	125	33.382196	-88.609409	Yes
S-27	Ephemeral	26	33.382123	-88.610159	Yes
S-28	Intermittent	281	33.383820	-88.610585	Yes
S-29	Intermittent	3,430	33.385584	-88.617176	Yes
S-30	Ephemeral	592	33.388403	-88.616447	Yes
S-30	Intermittent	812	33.386607	-88.616824	Yes
S-31	Perennial	1,225	33.387178	-88.624430	Yes
S-32	Ephemeral	743	33.388455	-88.625646	Yes
S-33	Perennial	4,837	33.382752	-88.625294	Yes
S-34	Ephemeral	403	33.382488	-88.630895	Yes
S-34	Intermittent	569	33.383384	-88.630041	Yes
S-35	Ephemeral	314	33.382634	-88.625302	Yes
S-35	Intermittent	940	33.382848	-88.627052	Yes
S-36	Ephemeral	608	33.383681	-88.626213	Yes
S-37	Ephemeral	89	33.382964	-88.626732	Yes
S-38	Ephemeral	88	33.383651	-88.628071	Yes
S-39	Ephemeral	30	33.384041	-88.629223	Yes
S-40	Ephemeral	294	33.385219	-88.630229	Yes
S-40	Intermittent	881	33.386650	-88.629530	Yes
S-41	Ephemeral	496	33.385017	-88.630584	Yes
S-42	Intermittent	951	33.385432	-88.632391	Yes
S-43	Ephemeral	267	33.385721	-88.632310	Yes
S-44	Intermittent	273	33.384673	-88.632255	Yes
S-45	Ephemeral	1,119	33.385135	-88.634096	Yes
S-46	Ephemeral	438	33.385327	-88.634724	Yes
S-47	Ephemeral	573	33.384824	-88.633379	Yes
S-48	Ephemeral	47	33.384964	-88.634193	Yes

MS Solar 6, LLC 3-7 Burns & McDonnell

Waterbody ID	Waterbody Type	Length of Stream (feet) in Survey Area	Latitude (N°)	Longitude (W°)	Anticipated Jurisdictional Status <sup>a</sup>
S-49	Ephemeral	148	33.384355	-88.633272	No
S-50	Intermittent	677	33.383387	-88.635013	Yes
S-51	S-51 Ephemeral		33.383112	-88.634557	Yes
S-52	Ephemeral	249	33.383282	-88.633892	Yes
S-53	Ephemeral	96	33.382538	-88.623974	Yes
S-54	Ephemeral	63	33.384107	-88.630742	Yes
S-55	S-55 Ephemeral		33.385893	-88.611832	Yes
S-56	Ephemeral	191	33.386057	-88.627318	Yes
S-57	Intermittent	23	33.381979	-88.609655	Yes
S-58	Intermittent	31	33.382007	-88.608449	Yes

<sup>(</sup>a) An official Jurisdictional Determination can only be provided by the USACE. Anticipated jurisdictional status of the ephemeral streams within the Study Area are based on pre-2015 definition of Waters of the U.S.

#### Palustrine Unconsolidated Bottom - PUB

Palustrine unconsolidated bottom wetlands accounted for approximately 14.15 acres (less than one percent) of the total Survey Area. Common vegetation associated with the banks and riparian areas of the PUBs consisted of tall fescue, white clover, tall buttercup, little blue stem, annual bluegrass, scutch grass, curly dock, spike rush, Unknown sedge (*Carex sp.*), *Rubus sp.*, osage orange, and black willow.

#### Palustrine Forested Wetlands - PFO

Palustrine forested wetlands accounted for less than one percent (2.8 acres) of the total Survey Area. The common overstory and understory vegetation consists of sugarberry, Osage orange, green ash, American sycamore, red maple and Chinese privet; and common vines, shrubs, and herbaceous vegetation consists of poison ivy, saw-tooth blackberry, butterweed, alligator weed, Virginia creeper, Virginia wild rye, and soft rush.

# Palustrine Emergent Wetlands - PEM

Palustrine emergent wetlands accounted for less than one percent (11.89 acres) of the total Survey Area. Common vegetation observed within the emergent wetlands included black willow, butterweed, shallow sedge, rough cockleburr, woolgrass (*Scirpus cyperinus*), curly dock, yellow nutsedge, blackberries, soft rush, and alligator weed.

#### Palustrine Scrub/Shrub - PSS

Palustrine scrub/shrub wetlands accounted for less than one percent (2.49 acres) of the total Survey Area. Common vegetation observed within the scrub/shrub wetlands included, green ash, water locust, butterweed, curly dock, and soft rush.

Table 3-2: Wetlands Identified in the Survey Area

Waterbody #	Waterbody Type	Area of Wetland (acre) in Survey Area	Latitude (N°)	Longitude (W°)	Anticipated Jurisdictional Status
W-PUB-01	PUB	0.82	33.390615	-88.622578	No
W-PUB-02	PUB	1.30	33.398449	-88.618331	Yes
W-PUB-03	PUB	0.98	33.400909	-88.621106	Yes
W-PUB-04	PUB	0.54	33.398434	-88.617285	Yes
W-PUB-05	PUB	0.12	33.398039	-88.614666	Yes
W-PUB-06	PUB	2.33	33.392168	-88.617827	Yes
W-PUB-07	PUB	0.64	33.400902	-88.604401	Yes
W-PUB-08	PUB	1.26	33.394151	-88.603418	Yes
W-PUB-09	PUB	1.26	33.396837	-88.597633	Yes
W-PUB-10	PUB	1.70	33.392798	-88.593924	Yes
W-PUB-11	PUB	0.20	33.401778	-88.598020	Yes
W-PUB-12	PUB	1.04	33.384690	-88.620443	Yes
W-PUB-13	PUB	0.52	33.387199	-88.609356	Yes
W-PUB-14	PUB	0.35	33.385786	-88.607932	Yes
W-PUB-15	PUB	0.01	33.385725	-88.626303	Yes
W-PUB-16	PUB	0.02	33.387018	-88.625912	Yes
W-PUB-17	PUB	0.01	33.382151	-88.610252	Yes
W-PUB-18	PUB	0.21	33.388520	-88.626905	Yes
W-PUB-19	PUB	0.74	33.386950	-88.632622	Yes
W-PUB-20	PUB	0.13	33.383228	-88.613899	Yes
W-PAB-01	PAB	0.05	33.392337	-88.594335	Yes
W-PEM-01	PEM	0.04	33.398553	-88.620092	No
W-PEM-02	PEM	0.12	33.396014	-88.598877	Yes
W-PEM-03	PEM	2.69	33.382459	-88.614429	Yes
W-PEM-04	PEM	1.91	33.384396	-88.606745	Yes
W-PEM-05	PEM	0.37	33.389049	-88.616437	Yes
W-PEM-06	PEM	0.13	33.387225	-88.624340	Yes
W-PEM-07	PEM	0.40	33.384263	-88.620243	Yes
W-PEM-08	PEM	5.95	33.385407	-88.617114	Yes
W-PEM-09	PEM	0.09	33.389411	-88.623204	Yes
W-PEM-10	PEM	0.20	33.385923	-88.602325	Yes
W-PSS-01	PSS	0.09	33.382495	-88.613855	Yes
W-PSS-02	PSS	0.02	33.383829	-88.628843	Yes
W-PFO-01	PFO	0.10	33.399329	-88.593939	Yes
W-PFO-02	PFO	0.49	33.393151	-88.593141	Yes
W-PFO-03	PFO	0.10	33.386080	-88.611841	Yes
W-PFO-04	PFO	0.27	33.392033	-88.595561	Yes

MS Solar 6, LLC 3-9 Burns & McDonnell

Waterbody #	Waterbody Type	Area of Wetland (acre) in Survey Area	Latitude (N°)	Longitude (W°)	Anticipated Jurisdictional Status
W-PFO-05	PFO	0.40	33.382596	-88.620742	Yes
W-PSS/PFO- 01	PSS/PFO	3.83	33.383761	-88.603730	Yes

<sup>(</sup>a) An official Jurisdictional Determination can only be provided by the USACE. The anticipated jurisdictionality of the ephemeral streams within the Study Area are based on the pre-2015 definition of Waters of the U.S.

#### 3.3 Conclusions

Based on the review of past studies, available online databases and mapping, and the detailed field assessment performed on the Survey Area, it is concluded that the Survey Area is generally consistent with the surrounding geography. Topography and land use are similar with adjacent properties and land use activities have been primarily related to agricultural and hay production.

Based upon the field surveys, it was determined that approximately 1,402 acres of the Survey Area are classified as dry land (uplands). There are approximately 25,525 linear feet of perennial streams, 13,138 linear feet of intermittent streams, and 12,295 feet of ephemeral streams. Approximately 14.2 acres are composed of PUB ponds. Table 3-4 shows different habitat types, approximate acreage, and length of each feature present in the Survey Area.

Factors considered in determining jurisdictional waters of the U.S. included criteria as defined under the recent U.S. District Court's relinquishment of the 2020 Navigable Waters Protection Rule and reinstatement of the pre-2015 definition of Waters of the U.S. Conditions observed during the wetland delineation survey determined that 39 wetlands and 59 streams within the Survey Area meet the definition of waters of the U.S. (Tables 1 and 2). The features indicated as "Yes" in Tables 1 and 2 are presumed to be under the jurisdiction of the USACE; however, an official Jurisdictional Determination can only be made by the USACE.

TVA is obligated under Executive Order 11990 to avoid the destruction, loss, or degradation to wetlands to the extent practicable. This obligation applies to all wetland features, including those wetlands that do not fall under USACE jurisdiction. Project siting should demonstrate wetland avoidance and minimization of wetland impacts for all wetland features on site, regardless of USACE jurisdictional status. If permanent impacts to jurisdictional waters of the U.S. cannot be completely avoided, they should be minimized to the extent practicable, and a Section 404 permit from the USACE may be required. Depending on the size, location, and purpose of the permanent or temporary impacts, a variety of Nationwide Permits may be used for access roads, road crossings, and land-based renewable energy generation facilities. Depending on the type and extent of impacts to waters of the U.S., permitting

requirements may range from a self-certifying Nationwide Permit, Nationwide Permit requiring a formal Pre-Construction Notification (PCN) submittal, or an individual permit may be required. Regardless of which Nationwide Permit(s) is applicable to the Project, the regional and general conditions of the Nationwide Permit(s) would apply and would need to be followed during Project construction.

# 4.0 REFERENCES


- Burns & McDonnell. 2020. Golden Triangle Solar Project: Wetland and Waterbody Delineation Report.


  Prepared for Origis Energy MS Solar 5. June 2020.
- Headwaters, Inc., 2016. Wetland Delineation and Determination for the GTRA Mega Site. Prepared for Calvert-Spradling Engineers, Inc. January 2016.
- U.S. Army Corps of Engineers. 1987. 1987 Corps of Engineers Wetland Delineation Manual (1987 Manual).
- U.S. Army Corps of Engineers. 2010. Regional Supplement to the Army Corps of Engineers Wetland


  Delineation Manual: Atlantic and Gulf Coast Plain Region Version 2.0 (Regional Supplement).

MS Solar 6, LLC 4-1 Burns & McDonnell


**APPENDIX A - FIGURES** 











Source: Esri, NRCS SSURGO Soils, Origis, Burns & McDonnell





















# Antecedent Precipitation vs Normal Range based on NOAA's Daily Global Historical Climatology Network



Coordinates	33.3908901, -88.6113077
Observation Date	2021-03-08
Elevation (ft)	248.73
Drought Index (PDSI)	Mild wetness
WebWIMP H <sub>2</sub> O Balance	Wet Season

30 Days Ending	30 <sup>th</sup> %ile (in)	70 <sup>th</sup> %ile (in)	Observed (in)	Wetness Condition	Condition Value	Month Weight	Product
2021-03-08	3.719685	8.05315	5.019685	Normal	2	3	6
2021-02-06	4.066536	6.616536	3.11811	Dry	1	2	2
2021-01-07	4.337402	6.242914	4.889764	Normal	2	1	2
Result							Normal Conditions - 10



Weather Station Name	Coordinates	Elevation (ft)	Distance (mi)	Elevation Δ	Weighted Δ	Days (Normal)	Days (Antecedent)
STATE UNIV	33.4692, -88.7822	185.039	11.242	63.691	5.775	10602	88
CRAWFORD 5 W	33.2783, -88.7061	252.953	9.511	4.223	4.32	607	0
STARKVILLE 4.7 SE	33.4033, -88.773	288.058	9.366	39.328	4.583	38	0
STARKVILLE 6.3 SSE	33.3673, -88.7975	305.118	10.866	56.388	5.502	54	0
STARKVILLE 5.3 S	33.3808, -88.8091	314.961	11.432	66.231	5.902	52	2





Photograph C-1: View of intermittent stream (S)-01, looking west.



Photograph C-2: View of intermittent S-02, looking east.





Photograph C-3: View of intermittent S-03, looking south.



Photograph C-4: View of ephemeral S-04, looking north.



Photographs
March 8 – 13, 2021
Lowndes County, Mississippi



Photograph C-5: View of perennial S-05, facing north.



Photograph C-6: View of intermittent S-06, facing east.





Photograph C-7: View of intermittent S-07, facing south.



Photograph C-8: View of ephemeral S-08, facing south.



Photographs
March 8 – 13, 2021
Lowndes County, Mississippi



Photograph C-9: View of intermittent S-12, facing east.



Photograph C-10: View of ephemeral S-13, facing northeast.





Photograph C-11: View of ephemeral portion of S-14, facing north.



Photograph C-12: View of intermittent portion of S-14, facing south.





Photograph C-13: View of ephemeral S-15, facing south.



Photograph C-14: View of perennial S-16, facing south.





Photograph C-15: View of culvert outlet from ephemeral S-16a into perennial S-16, facing southwest.



Photograph C-16: View of ephemeral S-17, facing northwest.





Photograph C-17: View of intermittent S-18, facing north.



Photograph C-18: View of intermittent S-19, facing south.





Photograph C-19: View of ephemeral S-20, facing northwest.



Photograph C-20: View of intermittent S-21, facing south.





Photograph C-21: View of ephemeral S-22, facing northwest.



Photograph C-22: View of perennial S-23, facing east.





Photograph C-23: View of ephemeral S-24, facing northwest.



Photograph C-24: View of intermittent S-25, facing northeast.





Photograph C-25: View of ephemeral S-26, facing southeast.



Photograph C-26: View of intermittent S-28, facing north.





Photograph C-27: View of intermittent S-29, facing southeast.



Photograph C-28: View of ephemeral S-30, facing south.





Photograph C-29: View of perennial S-31, facing northeast.



Photograph C-30: View of ephemeral S-32, facing northwest.





Photograph C-31: View of perennial S-33, facing northwest.



Photograph C-32: View of ephemeral portion of S-34, facing north.





Photograph C-33: View of intermittent portion of S-34, facing north.



Photograph C-34: View of intermittent S-35, facing west.





Photograph C-35: View of ephemeral S-36, facing south.



Photograph C-36: View of ephemeral S-37, facing south.





Photograph C-37: View of ephemeral S-38, facing south.



Photograph C-38: View of ephemeral S-39, facing east.





Photograph C-39: View of ephemeral portion of S-40, facing north.



Photograph C-40: View of intermittent portion of S-40, facing west.





Photograph C-41: View of ephemeral portion of S-41, facing northwest.



Photograph C-42: View of intermittent S-42, facing south.





Photograph C-43: View of ephemeral S-43, facing south.



Photograph C-44: View of intermittent S-44, facing southeast.





Photograph C-45: View of ephemeral S-45, facing southeast.



Photograph C-46: View of ephemeral S-46, facing southeast.





Photograph C-47: View of ephemeral S-47, facing southeast.



Photograph C-48: View of ephemeral S-48, facing north.





Photograph C-49: View of ephemeral S-49, facing northeast.



Photograph C-50: View of intermittent S-50, facing east.





Photograph C-51: View of ephemeral S-51, facing north.



Photograph C-52: View of ephemeral S-52, facing north.





Photograph C-53: View of ephemeral S-53, facing north.



Photograph C-54: View of ephemeral S-54, facing north.





Photograph C-55: View of ephemeral S-55, facing northeast.



Photograph C-56: View of ephemeral S-56, facing northwest.



 $\begin{array}{c} Photographs \\ March \ 8-13, \ 2021 \\ Lowndes \ County, \ Mississippi \end{array}$ 



Photograph C-57: View of intermittent S-57, facing north.



Photograph C-58: View of intermittent S-58, facing south.





Photograph C-59: View of wetland (W)-PAB-1, facing northeast.



Photograph C-60: View of W-PUB-01, facing north.





Photograph C-61: View of W-PUB-02, facing east.



Photograph C-62: View of W-PUB-03, facing northwest.





Photograph C-63: View of W-PUB-04, facing southeast.



Photograph C-64: View of W-PUB-05, facing southwest.





Photograph C-65: View of W-PUB-06, facing southwest.



Photograph C-66: View of W-PUB-07, facing southwest.





Photograph C-67: View of W-PUB-08, facing southwest.



Photograph C-68: View of W-PUB-09, facing east.





Photograph C-69: View of W-PUB-10, facing northeast.



Photograph C-70: View of W-PUB-11, facing southeast.





Photograph C-71: View of W-PUB-12, facing southwest.



Photograph C-72: View of W-PUB-13, facing south.





Photograph C-73: View of W-PUB-14, facing north.



Photograph C-74: View of W-PUB-15, facing southeast.





Photograph C-75: View of W-PUB-16, facing northwest.



Photograph C-76: View of W-PUB-17, facing northwest.





Photograph C-77: View of W-PUB-18, facing west.



Photograph C-78: View of W-PUB-19, facing north.





Photograph C-79: View of W-PUB-20, facing northeast.



Photograph C-80: View of sample plot in W-PEM-01, facing east.





Photograph C-81: View of upland sample plot adjacent to W-PEM-01, facing south.



Photograph C-82: View of sample plot in W-PEM-02, facing northwest.





Photograph C-83: View of upland sample plot adjacent to W-PEM-02, facing north.



Photograph C-84: View of flooded portion of W-PEM-03, facing west.





Photograph C-85: View of upland sample plot adjacent to W-PEM-03, facing northwest.



Photograph C-86: View of sample plot in W-PEM-04, facing northwest.



 $\begin{array}{c} Photographs\\ March~8-13,~2021\\ Lowndes~County,~Mississippi \end{array}$ 



Photograph C-87: View of upland sample plot adjacent to W-PEM-04, facing south.



Photograph C-88: View of sample plot in W-PEM-05, facing west.





Photograph C-89: View of upland sample plot adjacent to W-PEM-05, facing south.



Photograph C-90: View of W-PEM-06, facing southwest.





Photograph C-91: View of upland area adjacent to W-PEM-06, facing north.



Photograph C-92: View of W-PEM-07, facing east.





Photograph C-93: View of upland fallow field adjacent to W-PEM-07, facing northeast.



Photograph C-94: View of W-PEM-08, facing south.





Photograph C-95: View of upland field adjacent to W-PEM-08, facing east.



Photograph C-96: View of W-PEM-09, facing northwest.





Photograph C-97: View of W-PEM-10, facing southeast.



Photograph C-98: View of W-PSS-01, facing south.





Photograph C-99: View of upland area adjacent to W-PSS-01, facing southeast.



Photograph C-100: View of W-PSS-02, facing southeast.





Photograph C-101: View of W-PFO-01, facing west.



Photograph C-102: View of upland sample plot adjacent to W-PFO-01, facing southwest.





Photograph C-103: View of flooded W-PFO-02, facing southeast.



Photograph C-104: View of flooded W-PFO-03, facing northeast.





Photograph C-105: View of upland area adjacent to W-PFO-03, facing northwest.



Photograph C-106: View of W-PFO-04, facing southeast.





Photograph C-107: View of W-PFO-05, facing south.



Photograph C-108: View of upland area adjacent to W-PFO-05, facing northeast.





Photograph C-109: View of W-PSS/PFO-1, facing northwest.



Photograph C-110: View of upland sample plot adjacent to W-PSS/PFO-1, facing north.





Photograph C-111: View of upland confirmation sample plot, SP-UPL-01, facing southeast.



Photograph C-112: View of upland confirmation sample plot, SP-UPL-02, facing north.





Photograph C-113: View of upland confirmation sample plot, SP-UPL-03, facing southwest.



Photograph C-114: View of upland confirmation sample plot, SP-UPL-04, facing north.





Photograph C-115: View of upland confirmation sample plot, SP-UPL-05, facing west.



Photograph C-116: View of upland confirmation sample plot, SP-UPL-06, facing south.





Photograph C-117: View of upland confirmation sample plot, SP-UPL-07, facing south.



Photograph C-118: View of upland confirmation sample plot, SP-UPL-08, facing southwest.



APPENDIX D -	REPRESENTA	TIVE WETLANI	D DETERMINAT	TION DATA FORMS

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2 Cit	y/County: Lowndes County Sampling Date: 2021	-03-08
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-U	
	ction, Township, Range: S28 T18N R16E	
Landform (hillslope, terrace, etc.): Upland Loc		2
Subregion (LRR or MLRA): P 135A Lat: 33.3928		
Soil Map Unit Name: Brooksville silty clay, 0 to 1 percent slopes		
Are climatic / hydrologic conditions on the site typical for this time of year?	·	
Are Vegetation, Soil, or Hydrology significantly dis		lo.
Are Vegetation, Soil, or Hydrology naturally proble  SUMMARY OF FINDINGS – Attach site map showing sa		es etc
Hydrophytic Vegetation Present? Yes No	Is the Sampled Area	
Hydric Soil Present?  Yes No V	within a Wetland? Yes No	
Wetland Hydrology Present? Yes No✓ Remarks:		
Upland confirmation sample plot. The APT sh	owed normal conditions at the time leading	g up
to and during the field survey.		
HYDROLOGY		
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two re	equired)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)	
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface	e (B8)
High Water Table (A2)  Harl Deposits (B15) (L		
☐ Saturation (A3) ☐ Hydrogen Sulfide Odo	-	
	s along Living Roots (C3) Dry-Season Water Table (C2)	
Sediment Deposits (B2)  Presence of Reduced  Presence of Reduced	— ` · · · · · · · · · · · · · · · · · ·	(00)
Drift Deposits (B3)  Recent Iron Reduction  This Must Surface (C)		(C9)
☐ Algal Mat or Crust (B4) ☐ Thin Muck Surface (C	<i>'</i> = ' ` <i>'</i>	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)	
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)	
Field Observations:		
Surface Water Present? Yes No Depth (inches): _		
Water Table Present? Yes No _ ✓ Depth (inches): _		
Saturation Present? Yes No Depth (inches): _	Wetland Hydrology Present? Yes No	<u> </u>
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos,		
Remarks:		
No indicators are met.		

## **VEGETATION (Four Strata)** – Use scientific names of plants.

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
<u>Tree Stratum</u> (Plot size: 30 ft r		Species?		Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Dercent of Deminant Charles
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 50 (A/B)
6				
7.				Prevalence Index worksheet:
8.				Total % Cover of: Multiply by:
		= Total Co	/er	OBL species $0 \times 1 = 0$
50% of total cover:				FACW species <u>5</u> x 2 = <u>10</u>
Sapling/Shrub Stratum (Plot size: 30 ft r )	20 /0 01	total cover		FAC species 60 x 3 = 180
				FACU species <u>40</u> x 4 = <u>160</u>
1				UPL species $5$ $x = 25$
2				Column Totals: 110 (A) 375 (B)
3				(2)
4				Prevalence Index = $B/A = 3.4$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
	=	= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 30 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Schedonorus arundinaceus	60		FAC	be present, unless disturbed or problematic.
2. Schizachyrium scoparium	30	<b>~</b>	FACU	Definitions of Four Vegetation Strata:
3. Elymus hystrix	5		UPL	To a Manda de de contrata estada de contrata de contra
4. Elymus repens	5		FACU	<b>Tree</b> – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
5 Poa annua	5		FACU	height.
6 Ranunculus acris	5		FACW	Carling (Charles Was donal and a supporting size of land
~·	· <u> </u>			Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	<u>110%</u> :	= Total Co	ver	
50% of total cover: 55	20% of	total cover	: 22	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4.				
5				Hydrophytic Vegetation
<b></b>		= Total Co		Present? Yes No
50% of total cover:		total cover	:	
Remarks: (If observed, list morphological adaptations below	ow).			
No test is met.				

Sampling Point: SP-UPL-01

SOIL Sampling Point: SP-UPL-01

Profile Desc	ription: (Describe	to the dep	th needed to docur	nent the	indicator	or confir	m the absence of in	ndicators.)
Depth	Matrix			x Feature		. 2	<u>.</u> .	
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	Texture	Remarks
0 - 10	10YR 3/2	100					Clay Loam	
10 - 18	10YR 3/2	80	5YR 4/6	20	<u> </u>	<u>M</u>	Clay Loam	
-								
-						-		
			-	-		-		
						-		
<u> </u>								
				-				
			Reduced Matrix, MS			rains.		=Pore Lining, M=Matrix.
1		cable to all	LRRs, unless other		•	I DD C T		Problematic Hydric Soils <sup>3</sup> :
Histosol (	ipedon (A2)		Polyvalue Be				. —	(A9) <b>(LRR O)</b> (A10) <b>(LRR S)</b>
Black His			Loamy Muck					/ertic (F18) (outside MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-	. , .	,	7 7	Floodplain Soils (F19) (LRR P, S, T)
	Layers (A5)		Depleted Ma	trix (F3)			Anomalous	s Bright Loamy Soils (F20)
	Bodies (A6) (LRR I		Redox Dark		•		(MLRA 1	
_	cky Mineral (A7) <b>(L</b>		= :					t Material (TF2) ow Dark Surface (TF12)
_	esence (A8) <b>(LRR</b> ( ck (A9) <b>(LRR P, T)</b>		Redox Depre		-8)			ow Dark Surface (1F12) Ilain in Remarks)
_	Below Dark Surfa		Depleted Ocl	-	) <b>(MLRA</b> 1	151)	<u> </u>	nam m remarks)
Thick Da	rk Surface (A12)	, ,	Iron-Mangan				P, T) <sup>3</sup> Indicator	s of hydrophytic vegetation and
	airie Redox (A16) (		· <b>—</b>					l hydrology must be present,
	ucky Mineral (S1) (	(LRR O, S)	Delta Ochric					disturbed or problematic.
	leyed Matrix (S4) edox (S5)		Reduced Ver					
_	Matrix (S6)						RA 149A, 153C, 153	3D)
_	face (S7) (LRR P,	S, T, U)		g =	,	(· ) <b>(</b>		,
Restrictive L	ayer (if observed)	):						
Type:								
Depth (inc	:hes):						Hydric Soil Pres	sent? Yes No
Remarks:								
No indic	ators met.							

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-08
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-UPL-02
••	ion, Township, Range: S28 T18N R16E
Landform (hillslope, terrace, etc.): Upland, Hillslope  Loca	
Subregion (LRR or MLRA): P 135A Lat: 33.39326	
Soil Map Unit Name: Brooksville silty clay, 1 to 3 percent slopes	NWI classification: N/A
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distu	
Are Vegetation, Soil, or Hydrology naturally problem  SUMMARY OF FINDINGS – Attach site map showing sai	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	Is the Sampled Area
Hydric Soil Present? Yes No	within a Wetland? Yes No
Remarks:	
	awad narmal canditions at the time leading up
Upland confirmation sample plot. The APT sho	wed normal conditions at the time leading up
to and during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1)  Aquatic Fauna (B13)  Mad Banasite (B45) (1.5)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Saturation (A3)  High Water Table (A2)  Hydrogen Sulfide Odor of the Action (A3)	
Water Marks (B1)  Oxidized Rhizospheres	` '
Sediment Deposits (B2)  Presence of Reduced In	
Drift Deposits (B3) Recent Iron Reduction is	_ ` ` · · · · · · · <del>_</del> · · · · · · · · · · · · · · · · · · ·
Algal Mat or Crust (B4) Thin Muck Surface (C7)	Geomorphic Position (D2)
☐ Iron Deposits (B5) ☐ Other (Explain in Remai	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:  Surface Water Present?  Yes No _  Depth (inches):	
Water Table Present?  Yes No _ Depth (inches):	
Saturation Present? Yes No Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	evious inspections), if available:
Remarks:	
No indicators are met.	

## **VEGETATION (Four Strata)** – Use scientific names of plants.

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species
1. Maclura pomifera	60		FACU	That Are OBL, FACW, or FAC: $\frac{2}{}$ (A)
2. Celtis laevigata	10		FACW	Total Number of Dominant
3. Juniperus virginiana	10		FACU	Species Across All Strata: 3 (B)
Gleditsia triacanthos	5		FAC	Percent of Dominant Species That Are OBL, FACW, or FAC: 66.7 (A/B)
6				Prevalence Index worksheet:
7				
8				
	85% =	Total Cov	er	FACW species $\frac{3}{10}$ $\frac{1}{20}$ $\frac{1}{20}$
50% of total cover: <u>43</u>	20% of	total cover:	17	00 070
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species 90 x 3 = 270  FACU species 70 x 4 = 280
1				
2				470
3				Column Totals: 170 (A) 570 (B)
4				Prevalence Index = $B/A = 3.35$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
		Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover:		
Herb Stratum (Plot size: 5 ft r )  Schedonorus arundinaceus	60	~	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must
2. Ligustrum sinense	20		FAC	be present, unless disturbed or problematic.
				Definitions of Four Vegetation Strata:
3. Rumex crispus			FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4 5				more in diameter at breast height (DBH), regardless of height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				Herb – All herbaceous (non-woody) plants, regardless
9.				of size, and woody plants less than 3.28 ft tall.
10				
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12.				
	85% =	Total Cov	er	
50% of total cover: 43				
Woody Vine Stratum (Plot size: 30 ft r )				
1	_			
2				
3				
4				
5				Hydrophytic
		Total Cov	er	Vegetation
50% of total cover:	20% of	total cover:		Present? Yes No
Remarks: (If observed, list morphological adaptations below				
· · · · · · · · · · · · · · · · · · ·	,-			
Dominance test is met.				

Sampling Point: SP-UPL-02

SOIL Sampling Point: SP-UPL-02

Profile Desc	ription: (Describe	to the dept	h needed to docur	nent the	indicator	or confir	rm the absence	of indicators.)	
Depth	Matrix	0/		x Feature		Loc. 2	_ Texture	Domorko	
(inches) 0 - 6	Color (moist) 10YR 2/2	<u>%</u> 98	Color (moist) 10YR 3/3	<u>%</u> 2	Type <sup>1</sup>	Loc <sup>2</sup>	Clay Loam	Remarks	
								Ocument	
6 - 18	10YR 3/3	60	10YR 2/2	35	_ <u>C</u>	<u>M</u>	Clay Loam	Compact	
6 - 18			7.5YR 4/6	5	<u>C</u>	М			
					_				
						_			
-									
<sup>1</sup> Type: C=C	oncentration, D=Dep	oletion, RM=	Reduced Matrix, MS	S=Maske	ed Sand G	rains.	<sup>2</sup> Location:	PL=Pore Lining, M=Matrix.	
	Indicators: (Applic							for Problematic Hydric Soils <sup>3</sup> :	
Histosol			Polyvalue Be		. , .		. —	/luck (A9) <b>(LRR O)</b>	
_	oipedon (A2)		Thin Dark Su					Muck (A10) (LRR S)	
	stic (A3) en Sulfide (A4)		Loamy Muck Loamy Gleye	-	. , .	R O)		ed Vertic (F18) <b>(outside MLRA 150</b> / ont Floodplain Soils (F19) <b>(LRR P, S</b>	
	d Layers (A5)		Depleted Ma		(1 2)			alous Bright Loamy Soils (F20)	, ',
	Bodies (A6) (LRR F	P, T, U)	Redox Dark		(F6)			RA 153B)	
	ıcky Mineral (A7) <b>(L</b>		Depleted Dai					arent Material (TF2)	
_	esence (A8) (LRR L	J)	Redox Depre		F8)			hallow Dark Surface (TF12)	
_	ıck (A9) <b>(LRR P, T)</b> d Below Dark Surfac	e (A11)	Marl (F10) <b>(L</b> Depleted Ocl		) (MLRA 1	51)	Other	(Explain in Remarks)	
I = '	ark Surface (A12)	(, , , , ,	☐ Iron-Mangan				P, T) <sup>3</sup> Indic	eators of hydrophytic vegetation and	
	rairie Redox (A16) <b>(</b>		_					land hydrology must be present,	
	Mucky Mineral (S1) (	LRR O, S)	Delta Ochric					ess disturbed or problematic.	
	Gleyed Matrix (S4) Redox (S5)		Reduced Ver						
_	Matrix (S6)						.RA 149A, 153C	, 153D)	
Dark Su	rface (S7) (LRR P,		_	Ü	,	`	•	•	
Restrictive	Layer (if observed)	:							
Type:								_	
	ches):		<u></u>				Hydric Soil	Present? Yes No	_
Remarks:									
No indic	ators are me	et.							
İ									

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2	City/County: Lowndes County Sampling Date: 2021-03-08
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-UPL-03
	Section, Township, Range: S27 T18N R16E
Landform (hillslope, terrace, etc.): Upland, Flat	Local relief (concave, convex, none): None Slope (%): 0
Subregion (LRR or MLRA): P 135A Lat: 33	3.394241 Long: -88.615904 Datum: NAD 83
Soil Map Unit Name: Vaiden silty clay, 0 to 2 percent slop	
Are climatic / hydrologic conditions on the site typical for this time of	of year? Yes No (If no, explain in Remarks.)
Are Vegetation, Soil, or Hydrology significa	antly disturbed? Are "Normal Circumstances" present? YesNo
Are Vegetation, Soil, or Hydrology naturall	ly problematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map show	ving sampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No _ ✔	,
Hydrophytic Vegetation Present? Yes No No	Is the Sampled Area
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	
Upland confirmation sample plot. The AF	PT showed normal conditions at the time leading up
	To show a frontial containents at the time reading ap
to and during the field survey.	
LIVEROLOGY	
HYDROLOGY	Occasional and in discharge (as in insurance of the constraint)
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that ap	
Surface Water (A1)  Aquatic Fauna	
	(B15) (LRR U) Drainage Patterns (B10)
Saturation (A3)  Hydrogen Sulfi	
	ospheres along Living Roots (C3) Dry-Season Water Table (C2)
1 <del></del>	educed Iron (C4)  Crayfish Burrows (C8)  eduction in Tilled Soils (C6)  Saturation Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)  Thin Muck Sur	
Iron Deposits (B5)  Other (Explain	<del>_</del>
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes No _ ✓ _ Depth (inc	ches):
Water Table Present? Yes No ✓ Depth (inc	ches):
Saturation Present? Yes No _ ✓ Depth (inc	
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial p	photos, provious inspections), if available:
Describe Necorded Data (Stream gauge, monitoring well, dental p	onotos, previous inspections), ii avaliable.
Remarks:	
No indicators are met.	
indicators are met.	

### **VEGETATION (Four Strata)** – Use scientific names of plants.

Sampling Point: SP-UPL-03
Dominance Test worksheet:
Number of Dominant Species
That Are OBL, FACW, or FAC: 1 (A)
Total Number of Dominant
Species Across All Strata: 4 (B)
Percent of Dominant Species
That Are OBL, FACW, or FAC: 25 (A/B)
Prevalence Index worksheet:
Total % Cover of: Multiply by:
OBL species $0 \times 1 = 0$
FACW species $\frac{0}{7}$ $x = \frac{0}{21}$
· — — — —
FACU species $62$ $x 4 = 248$
UPL species $0 \times 5 = 0$
Column Totals: <u>69</u> (A) <u>269</u> (B)
Prevalence Index = B/A = 3.9
Hydrophytic Vegetation Indicators:
1 - Rapid Test for Hydrophytic Vegetation
2 - Dominance Test is >50%
Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
1
<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
Definitions of Four Vegetation Strata:
Definitions of Four Vegetation Strata.
Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
more in diameter at breast height (DBH), regardless of height.
Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
than 3 in. DBH and greater than 3.20 it (1 in) tail.
Herb – All herbaceous (non-woody) plants, regardless
of size, and woody plants less than 3.28 ft tall.
Woody vine – All woody vines greater than 3.28 ft in
height.
Hydrophytic
Vegetation
Present? Yes No
Veg

SOIL Sampling Point: SP-UPL-03

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the	indicator	or confirm	the absence	of indicators.)	
Depth	Matrix			x Feature		. 2	<b>-</b> .		
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 18	10YR 4/4	_ <u>80 _</u> _					Clay Loam	Mixed Matrix	
0 - 18	5PB 4/1	20							
				_					
-									
_					·				
	-								-
				-	- ——				<del>-</del>
1							2		
	oncentration, D=De Indicators: (Application)					ains.		PL=Pore Lining, M=Matri for Problematic Hydric 9	
Histosol		cable to all L	Polyvalue Be		•	DD C T I		Muck (A9) (LRR O)	Jolis .
	oipedon (A2)		Thin Dark Su					Muck (A9) (LRR S)	
Black Hi			Loamy Muck					ed Vertic (F18) (outside I	MLRA 150A,B)
	en Sulfide (A4)		Loamy Gleye	-		,		ont Floodplain Soils (F19)	
	d Layers (A5)		Depleted Ma				·	alous Bright Loamy Soils (	F20)
	Bodies (A6) (LRR I		Redox Dark	•	•		`	RA 153B)	
_	icky Mineral (A7) <b>(L</b> esence (A8) <b>(LRR</b> (		Depleted Da					arent Material (TF2) Shallow Dark Surface (TF1	2)
	ick (A9) <b>(LRR P, T)</b>		Marl (F10) (L		0)			(Explain in Remarks)	2)
	d Below Dark Surfa		Depleted Oct		(MLRA 1	51)		(Explain in Normanio)	
	ark Surface (A12)		Iron-Mangan	ese Mass	ses (F12) <b>(</b>	LRR O, P,	T) <sup>3</sup> Indic	cators of hydrophytic vege	tation and
	rairie Redox (A16) (		_			, U)		tland hydrology must be pr	
	Mucky Mineral (S1)	(LRR O, S)	Delta Ochric			OA 450D)		ess disturbed or problema	tic.
	Gleyed Matrix (S4) Redox (S5)		Reduced Ver						
	Matrix (S6)						A 149A, 153C	. 153D)	
	rface (S7) (LRR P,	S, T, U)			, (	, (	,	,,	
Restrictive	Layer (if observed)	):							
Type:			<u> </u>						
Depth (in	ches):		<u></u>				Hydric Soil	Present? Yes	No <u>′</u>
Remarks:									
No indic	ators are me	et.							

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2	City/County: Lowndes Cour	nty	Sampling Date: 2021-03-09
Applicant/Owner: TVA			Sampling Point: SP-UPL-04
• • • • • • • • • • • • • • • • • • • •	Section, Township, Range: S2		
Landform (hillslope, terrace, etc.): Upland	· · · · · ·		Slope (%): 1
Subregion (LRR or MLRA): P 135A Lat: 33.3			Datum: NAD 83
Soil Map Unit Name: Vaiden silty clay, 2 to 5 percent slopes,		NWI classifica	
Are climatic / hydrologic conditions on the site typical for this time of y			
Are Vegetation, Soil, or Hydrology significantly			
Are Vegetation, Soil, or Hydrology naturally processed in the second	oblematic? (If needed, e	xplain any answer	rs in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing	g sampling point locatio	ns, transects	, important features, etc.
Hudombutia Vanatatian Brasanta Van			
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area		,
Wetland Hydrology Present? Yes No _ ✓	within a Wetland?	Yes	No
Remarks:			
Upland confirmation sample plot. The APT	showed normal cor	ditions at t	he time leading un
	Showed Horrial Con	iditions at t	ne time leading up
to and during the field survey.			
HYDROLOGY			
Wetland Hydrology Indicators:		Secondary Indica	tors (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)		Surface Soil (	Cracks (B6)
Surface Water (A1) Aquatic Fauna (B	3)	Sparsely Veg	etated Concave Surface (B8)
High Water Table (A2)  Harl Deposits (B1)		Drainage Pat	terns (B10)
Saturation (A3) Hydrogen Sulfide	, ,	Moss Trim Li	` '
	neres along Living Roots (C3)		Vater Table (C2)
Sediment Deposits (B2)  Presence of Redu  Draw (P8)	` '	Crayfish Burr	
	ction in Tilled Soils (C6)		sible on Aerial Imagery (C9)
☐ Algal Mat or Crust (B4) ☐ Thin Muck Surface ☐ Iron Deposits (B5) ☐ Other (Explain in	` '	Geomorphic Shallow Aqui	` ,
Inundation Visible on Aerial Imagery (B7)	(emarks)	FAC-Neutral	` '
Water-Stained Leaves (B9)		=	oss (D8) <b>(LRR T, U)</b>
Field Observations:			, , ,
Surface Water Present? Yes No Depth (inches	3):		
Water Table Present? Yes No _ ✓ Depth (inches	3):		
Saturation Present? Yes No Depth (inches	s): Wetland H	ydrology Presen	t? Yes No <u></u>
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial pho	os previous inspections) if avai	ilahle <sup>.</sup>	
2000/150 Notice Pulse (officially gauge, monitoring won, defining pro-	so, providuo iriopositorioj, ir avai	indolo.	
Remarks:			
No indicators are met.			

### **VEGETATION (Four Strata)** – Use scientific names of plants.

Dominance Test worksheet:   Number of Dominant Species   That Are OBL, FACW, or FAC:   1
That Are OBL, FACW, or FAC: 1 (A)  Total Number of Dominant Species Across All Strata: 3 (B)  Percent of Dominant Species That Are OBL, FACW, or FAC: 33.3 (A/B)  Prevalence Index worksheet:  Total % Cover of: Multiply by:  OBL species 0 x 1 = 0  FACW species 0 x 2 = 0  FAC species 70 x 3 = 210  FACU species 55 x 4 = 220  UPL species 0 x 5 = 0  Column Totals: 125 (A) 430 (B)  Prevalence Index = B/A = 3.44  Hydrophytic Vegetation Indicators:
Species Across All Strata: 3 (B)   Percent of Dominant Species That Are OBL, FACW, or FAC: 33.3 (A/B)   Prevalence Index worksheet:
Species Across All Strata: 3 (B)   Percent of Dominant Species That Are OBL, FACW, or FAC: 33.3 (A/B)   Prevalence Index worksheet:
That Are OBL, FACW, or FAC: 33.3 (A/B)    Prevalence Index worksheet:
That Are OBL, FACW, or FAC: 33.3 (A/B)    Prevalence Index worksheet:
Total % Cover of:         Multiply by:           OBL species         0         x 1 = 0           FACW species         0         x 2 = 0           FAC species         70         x 3 = 210           FACU species         55         x 4 = 220           UPL species         0         x 5 = 0           Column Totals:         125         (A)         430         (B)           Prevalence Index         = B/A = 3.44           Hydrophytic Vegetation Indicators:
Total % Cover of:         Multiply by:           OBL species         0         x 1 = 0           FACW species         0         x 2 = 0           FAC species         70         x 3 = 210           FACU species         55         x 4 = 220           UPL species         0         x 5 = 0           Column Totals:         125         (A)         430         (B)           Prevalence Index         = B/A = 3.44           Hydrophytic Vegetation Indicators:
OBL species 0
FACW species 0
FAC species 70
FACU species $55$ $x = 4$ $220$ UPL species $0$ $x = 5$ $0$ Column Totals: $125$
UPL species $0$ $x = 0$ $430$ (B)  Prevalence Index $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$
Column Totals: 125 (A) 430 (B)  Prevalence Index = B/A = 3.44  Hydrophytic Vegetation Indicators:
Prevalence Index = B/A = 3.44  Hydrophytic Vegetation Indicators:
Hydrophytic Vegetation Indicators:
Hydrophytic Vegetation Indicators:
1 - Rapid Test for Hydrophytic Vegetation
2 - Dominance Test is >50%
3 - Prevalence Index is ≤3.0 <sup>1</sup>
Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
<sup>1</sup> Indicators of hydric soil and wetland hydrology must
be present, unless disturbed or problematic.
Definitions of Four Vegetation Strata:
Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
more in diameter at breast height (DBH), regardless of
height.
Sapling/Shrub – Woody plants, excluding vines, less
than 3 in. DBH and greater than 3.28 ft (1 m) tall.
Herb – All herbaceous (non-woody) plants, regardless
of size, and woody plants less than 3.28 ft tall.
Woody vine – All woody vines greater than 3.28 ft in
height.
Heater wheets
Hydrophytic
—— Hydrophytic Vegetation Present? Yes No

SOIL Sampling Point: SP-UPL-04

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	ndicator	or confirn	n the absence of	indicato	rs.)	
Depth Matrix Color (moist) %		Redox Features			Loc <sup>2</sup>	Toyturo		Remarks		
(inches)			Color (moist)	<u>%</u>	Type <sup>1</sup>	LOC	<u>Texture</u>		Remarks	
0 - 18	10YR 3/2	_ <u>100</u> _					Clay Loam			
-										
-					'					
<sup>1</sup> Type: C=Co	oncentration, D=De	pletion, RM=R	educed Matrix, MS	S=Masked	Sand Gr	ains.	<sup>2</sup> Location: Pl	L=Pore Li	ning, M=Matr	rix.
Hydric Soil	Indicators: (Appli	cable to all LF	RRs, unless other	wise note	ed.)		Indicators fo	r Probler	matic Hydric	Soils <sup>3</sup> :
Histosol	(A1)		Polyvalue Be	low Surfa	ce (S8) <b>(L</b>	.RR S, T, l	<b>U)</b> 📙 1 cm Mud	ck (A9) <b>(L</b>	.RR O)	
	pipedon (A2)		Thin Dark Su				2 cm Mud			
Black Hi			Loamy Muck			R O)			, .	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye		F2)					) (LRR P, S, T)
	d Layers (A5)	D T 11)	Depleted Mat		.0)			_	Loamy Soils	(F20)
· = ·	Bodies (A6) (LRR I cky Mineral (A7) (L		Redox Dark S Depleted Dar	`	,		(MLRA	a 153B) ent Materi	al (TE2)	
	esence (A8) <b>(LRR</b>		Redox Depre		, ,				Surface (TF	12)
_	ick (A9) (LRR P, T)	•	Marl (F10) (L		5)			xplain in F		12)
	d Below Dark Surfa		Depleted Och	-	(MLRA 1	51)	(		,	
Thick Da	ark Surface (A12)		Iron-Mangan				, <b>T)</b> <sup>3</sup> Indicate	ors of hyd	rophytic vege	etation and
_	rairie Redox (A16)		Umbric Surfa	ce (F13) (	LRR P, T	', U)	wetlar	nd hydrolo	ogy must be p	oresent,
_	lucky Mineral (S1)	(LRR O, S)	Delta Ochric					s disturbe	d or problema	atic.
_	Gleyed Matrix (S4)		Reduced Ver							
	ledox (S5)		Piedmont Flo					505)		
_	Matrix (S6)	C T II)	Anomalous E	right Loar	ny Soils (	F20) (MLR	RA 149A, 153C, 1	53D)		
	rface (S7) (LRR P, _ayer (if observed						<del>                                     </del>			
Type:	Layer (II Observed	<i>,</i> .								
l	aboa):		_				Usalaio Coil Da		Yes	No 🗸
	ches):		<u> </u>				Hydric Soil Pr	resent?	res	_ NO <u>*</u>
Remarks:		_								
No indic	ators are m	et.								

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2 City	County: Lowndes County Sampling Date: 2021-03-09					
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-UPL-05					
Investigator(s): J. Brown, O. Haney Section, Township, Range: S27 T18N R16E						
Landform (hillslope, terrace, etc.): Upland, Flat  Local relief (concave, convex, none): None  Slope (%): 0						
Subregion (LRR or MLRA): P 135A Lat: 33.398183 Long: -88.602480 Datum: NAD 83 Soil Map Unit Name: Vaiden silty clay, 0 to 2 percent slopes NWI classification: NA						
Are climatic / hydrologic conditions on the site typical for this time of year?						
Are Vegetation, Soil, or Hydrology significantly distr						
Are Vegetation, Soil, or Hydrology naturally probler	natic? (If needed, explain any answers in Remarks.)					
SUMMARY OF FINDINGS - Attach site map showing sa	mpling point locations, transects, important features, etc.					
Hydrophytia Vagetation Propert?						
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area					
Wetland Hydrology Present? Yes No	within a Wetland? Yes No					
Remarks:						
Unland confirmation sample plot. The APT sho	owed normal conditions at the time leading up					
	wed normal conditions at the time leading ap					
to and during the field survey.						
HYDROLOGY						
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)					
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)					
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)					
High Water Table (A2)  Marl Deposits (B15) (LI						
Saturation (A3) Hydrogen Sulfide Odor	` '					
	along Living Roots (C3) Dry-Season Water Table (C2)					
Sediment Deposits (B2)  Drift Deposits (B3)  Presence of Reduced II  Recent Iron Reduction	<del>-</del> '					
Algal Mat or Crust (B4)  Thin Muck Surface (C7						
Iron Deposits (B5)  Other (Explain in Rema	· · · · · · · · · · · · · · · · · · ·					
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)					
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)					
Field Observations:						
Surface Water Present? Yes No Depth (inches):						
Water Table Present? Yes No Depth (inches):						
Saturation Present? Yes No Depth (inches):	Wetland Hydrology Present? Yes No					
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos, p.	revious inspections), if available:					
Remarks:						
No indicators are met.						

imes of pi	arito.		Sampling Point: SP-UPL-0
			Dominance Test worksheet:
			Number of Dominant Species
			That Are OBL, FACW, or FAC: 2 (A)
- ——			Total Number of Dominant
			Species Across All Strata: 4 (B)
			Percent of Dominant Species
			That Are OBL, FACW, or FAC: 50 (A/B)
5		FAC	Prevalence Index worksheet:
80% =	Total Cov	er	OBL species $0 \times 1 = 0$ FACW species $5 \times 2 = 10$
20% of	total cover:	16	
			17to species x o
			FACU species $\frac{35}{25}$ $\times 4 = \frac{140}{135}$
			UPL species $35$ $x = 175$
			Column Totals: <u>87</u> (A) <u>361</u> (B)
			Prevalence Index = B/A = 4.15
			Hydrophytic Vegetation Indicators:
			1 - Rapid Test for Hydrophytic Vegetation
			2 - Dominance Test is >50%
			☐ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
			☐ Problematic Hydrophytic Vegetation¹ (Explain)
20% 01	total cover.		
5	./	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2			Definitions of Four Vegetation Strata:
			Definitions of Four Vegetation Strata:
			Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
			more in diameter at breast height (DBH), regardless of height.
			neight.
			Sapling/Shrub – Woody plants, excluding vines, less
			Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
			than 3 in. DBH and greater than 3.28 ft (1 m) tall. <b>Herb</b> – All herbaceous (non-woody) plants, regardless
			than 3 in. DBH and greater than 3.28 ft (1 m) tall.
			than 3 in. DBH and greater than 3.28 ft (1 m) tall. <b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
			than 3 in. DBH and greater than 3.28 ft (1 m) tall. <b>Herb</b> – All herbaceous (non-woody) plants, regardless
			than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
			than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
	- Total Cov	er	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% =	- Total Cov	er	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% =	Total Covers	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% = 20% of	Total Cover	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% = 20% of	= Total Cov total cover	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% = 20% of	= Total Covers	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in
7% = 20% of	= Total Cov total cover:	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in height.
7% = 20% of	= Total Cov total cover:	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in height.  Hydrophytic
7% = 20% of	= Total Covers	er 1	than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in height.
	Absolute % Cover 30 30 5 5 5 5 20% of 5 20% of	Absolute	30    V

since most species in the area are FAC or wetter.

SOIL Sampling Point: SP-UPL-05

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)										
Depth <u>Matrix</u>		Redox Feature								
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	<u>Texture</u>	Remarks		
0-6	10YR 3/4	50					Clay Loam			
0-6	10YR 5/3	50								
6 - 18	10YR 6/6	50	7.5YR 4/6	30	<u>C</u>	М	Silty Clay Loam			
6 - 18			2.5Y 6/1	20	D	М				
-										
					_	_	<u> </u>			
1 <sub>Type:</sub> C=C	oncontration D=Da	nlotion DM	-Poduced Matrix M	S=Mooks	d Sand C	roine	2l coation: DI =D	Joro Lining M-Matrix		
Type: C=Concentration, D=Depletion, RM=Reduced Matrix, MS=Masked Sand Grains.  Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)  Location: PL=Pore Lining, M=Matrix.  Indicators for Problematic Hydric Soils³:										
Histosol					•	LRR S. T.				
Histosol (A1) Polyvalue Below Surface (S8) (LRR S, T, U Thin Dark Surface (S9) (LRR S, T, U)							. –	A10) (LRR S)		
Black H	istic (A3)		Loamy Muck			R O)	Reduced Vertic (F18) (outside MLRA 150A,B)			
<del>     </del>	en Sulfide (A4)		Loamy Gleye		(F2)		☐ Piedmont Floodplain Soils (F19) (LRR P, S, T)			
	d Layers (A5)	D T 11)	Depleted Ma	, ,	(EC)		Anomalous Bright Loamy Soils (F20)			
	Bodies (A6) (LRR I ucky Mineral (A7) (L		Redox Dark Depleted Da		,		(MLRA 153B) Red Parent Material (TF2)			
	resence (A8) (LRR		Redox Depre				Very Shallow Dark Surface (TF12)			
	uck (A9) <b>(LRR P, T)</b>		Marl (F10) <b>(L</b>		,		Other (Explain in Remarks)			
	d Below Dark Surfa	ce (A11)	Depleted Oc				•			
_	ark Surface (A12)		Iron-Mangan					of hydrophytic vegetation and		
	rairie Redox (A16) ( lucky Mineral (S1) (		A) Umbric Surfa Delta Ochric				wetland hydrology must be present, unless disturbed or problematic.			
_	Bleyed Matrix (S4)	(LKK 0, 3)	Reduced Ve					sturbed or problematic.		
	Redox (S5)		Piedmont Flo							
Stripped	Matrix (S6)		Anomalous E	Bright Loa	amy Soils	(F20) <b>(MLI</b>	RA 149A, 153C, 153D	))		
	rface (S7) (LRR P,									
	Layer (if observed)	):								
Type:			<del></del>							
	ches):						Hydric Soil Prese	ent? Yes No		
Remarks:										
No indic	ators are m	et.								

# WETLAND DETERMINATION DATA FORM – Atlantic and Gulf Coastal Plain Region

Project/Site: Golden Triangle Phase 2	City/County: Lown	des County	Sampling Date: 2021-03-11					
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-UPL-06							
Investigator(s): J. Brown, O. Haney	Section, Township, Range: S26 T18N R16E							
Landform (hillslope, terrace, etc.): Upland, Flat			Slope (%): 0					
Subregion (LRR or MLRA): P 135A Lat: 33.39			Datum: NAD 83					
Soil Map Unit Name: Vaiden silty clay, 0 to 2 percent slopes NWI classification: PFO1A								
Are climatic / hydrologic conditions on the site typical for this time of ye								
Are Vegetation, Soil, or Hydrology significantly								
Are Vegetation, Soil, or Hydrology naturally pr								
SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.								
Hydrophytic Vegetation Present?  Hydric Soil Present?  Wetland Hydrology Present?  Remarks:  Yes No  V  Remarks:	Is the Samp within a We		No					
Upland confirmation sample plot. The APT showed normal conditions at the time leading up to and during the field survey.								
HYDROLOGY								
Wetland Hydrology Indicators:		Secondary Indicate	tors (minimum of two required)					
Primary Indicators (minimum of one is required; check all that apply)	)	Surface Soil (	Cracks (B6)					
Surface Water (A1) Aquatic Fauna (B1)	13)	Sparsely Veg	etated Concave Surface (B8)					
High Water Table (A2)  Harl Deposits (B1)	5) <b>(LRR U)</b>	<u></u> Drainage Pat	terns (B10)					
Saturation (A3) Hydrogen Sulfide	, ,	Moss Trim Lin	, ,					
☐ Water Marks (B1) ☐ Oxidized Rhizosph	heres along Living Ro	oots (C3) 🔲 Dry-Season V	Water Table (C2)					
Sediment Deposits (B2)	• •	☐ Crayfish Burn						
	ction in Tilled Soils (C	_	sible on Aerial Imagery (C9)					
Algal Mat or Crust (B4)	• •	Geomorphic I	` ,					
☐ Iron Deposits (B5) ☐ Other (Explain in F	Remarks)	☐ Shallow Aquit						
Inundation Visible on Aerial Imagery (B7)		FAC-Neutral	` '					
Water-Stained Leaves (B9)		<u></u> Sphagnum m	oss (D8) <b>(LRR T, U)</b>					
Field Observations:								
Surface Water Present? Yes No Depth (inches								
Water Table Present? Yes No Depth (inches								
Saturation Present? Yes No Depth (inches (includes capillary fringe)	s):	Wetland Hydrology Presen	t? Yes No					
Describe Recorded Data (stream gauge, monitoring well, aerial phot	tos, previous inspection	ons), if available:						
Remarks:								
Indicator B10 is met. There were pockets of	of standing wa	ater (<2") in some	microdepressions					
within the forested area. The standing wat	er was not as	sociated with a hi	gh water table					
meaning it can not be used as an indication			_					
met.		.,	a, a. e.e.g, .ee.					
illet.								

20.4		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species
1. Carya ovata	50		FACU	That Are OBL, FACW, or FAC: 3 (A)
2. Quercus pagoda	20		FACW	Total Number of Dominant
3. Quercus stellata	10		UPL	Species Across All Strata: 6 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 50 (A/B)
6				
7				Prevalence Index worksheet:
8				Total % Cover of: Multiply by:
	80% =	Total Cov	er	OBL species $0 \times 1 = 0$
50% of total cover: 40				FACW species 40 x 2 = 80
Sapling/Shrub Stratum (Plot size: 15 ft r )	20 70 01	10101 00101	· ———	FAC species <u>45</u> x 3 = <u>135</u>
1. Celtis laevigata	20	~	FACW	FACU species <u>70</u> x 4 = <u>280</u>
2. Carya ovata	10		FACU	UPL species 10 x 5 = 50
3. Ulmus alata	10		FACU	Column Totals: <u>165</u> (A) <u>545</u> (B)
1.1			FAC	
4. Ligustrum sinense	- —			Prevalence Index = $B/A = 3.3$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
	45% =	Total Cov	er	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover: <u>23</u>	20% of	total cover:	9	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Ligustrum sinense	30	<b>~</b>	FAC	be present, unless disturbed or problematic.
2. Carex sp.*	5		FAC	Definitions of Four Vegetation Strata:
3. Viola sororia	5		FAC	
**	· <u> </u>			Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4				more in diameter at breast height (DBH), regardless of height.
5				1.59.11
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	40% =	Total Cov	er	
50% of total cover: 20	20% of	total cover:	8	
Woody Vine Stratum (Plot size: 30 ft r )				
1.				
2.				
3.				
4				
5				Hydrophytic
		Total Cov		Vegetation Present? Yes No
50% of total cover:		total cover:		11000mi
Remarks: (If observed, list morphological adaptations below	ow).			
No test is met.				
140 test is met.				

Sampling Point: SP-UPL-06

SOIL Sampling Point: SP-UPL-06

Depth	ription: (Describe Matrix	·	needed to docum Redox	Features		the absence of	of indicators.)	
(inches)	Color (moist)	%	Color (moist)	% Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 24	10YR 3/2	100				Clay Loam		
-								
_								
¹Type: C=Cd	ncentration D=De	nletion RM=F	Reduced Matrix, MS	=Masked Sand G	rains	<sup>2</sup> Location:	PL=Pore Lining, M=Matrix.	
			RRs, unless other		Taillo.		or Problematic Hydric Soils <sup>3</sup> :	
☐ Histosol	(A1)		Polyvalue Bel	ow Surface (S8) (	LRR S, T, U	) 🔲 1 cm M	uck (A9) (LRR O)	
. <del></del>	pipedon (A2)			face (S9) (LRR S			uck (A10) (LRR S)	
Black Hi	stic (A3) n Sulfide (A4)		Loamy Mucky	Mineral (F1) (LR	R O)		d Vertic (F18) (outside MLRA 150 nt Floodplain Soils (F19) (LRR P, \$	-
_	I Layers (A5)		Depleted Mat				ous Bright Loamy Soils (F20)	), I <i>)</i>
	Bodies (A6) (LRR I	P, T, U)	Redox Dark S				A 153B)	
	cky Mineral (A7) <b>(L</b>		= '	k Surface (F7)			rent Material (TF2)	
	esence (A8) (LRR I ck (A9) (LRR P, T)		Redox Depre	, ,			nallow Dark Surface (TF12) Explain in Remarks)	
_	Below Dark Surfa		_ ` ` ` `	ric (F11) <b>(MLRA</b> 1	151)	U Other (i	Explain in Remarks)	
ı <del>=</del>	ark Surface (A12)	,		ese Masses (F12)		<b>T)</b> <sup>3</sup> Indica	ators of hydrophytic vegetation and	
_	rairie Redox (A16) (			ce (F13) <b>(LRR P,</b>			and hydrology must be present,	
	lucky Mineral (S1) ( leyed Matrix (S4)	(LRR O, S)		F17) <b>(MLRA 151</b> ) ic (F18) <b>(MLRA 1</b>		unle	ss disturbed or problematic.	
	edox (S5)			odplain Soils (F19		9A)		
_	Matrix (S6)			right Loamy Soils			153D)	
	face (S7) (LRR P,					1		
	_ayer (if observed)	):						
Type:	ches):					Hydric Soil I	Present? Yes No	,
Remarks:	<u> </u>		<u> </u>			Hyuric 30ii i	resent: res No	
	ators are me	o.t						
INO IIIGIC	ators are mi	<b>5</b> 1.						

Project/Site: Golden Triangle Phase 2 City/C	county: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: SP-UPL-07
	on, Township, Range: S27 T18N R16E
Landform (hillslope, terrace, etc.): Depression Local	
Subregion (LRR or MLRA): P 135A Lat: 33.395583	
Soil Map Unit Name: Vaiden silty clay, 2 to 5 percent slopes, erode	
Are climatic / hydrologic conditions on the site typical for this time of year? Y	
Are Vegetation, Soil, or Hydrology significantly distur	
Are Vegetation, Soil, or Hydrology naturally problems	
SUMMARY OF FINDINGS – Attach site map showing san	
Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?  Remarks:  Upland confirmation sample plot. The APT sho	Is the Sampled Area within a Wetland? Yes No  wed normal conditions at the time leading up
to and during the field survey.	are the time reading up
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  High Water Table (A2)  High Water Table (A2)  High Water Table (A2)	
□       Saturation (A3)       □       Hydrogen Sulfide Odor (0)         □       Water Marks (B1)       □       Oxidized Rhizospheres a	
Sediment Deposits (B2)  Presence of Reduced Iro	
Drift Deposits (B3)  Recent Iron Reduction in	
Algal Mat or Crust (B4) Thin Muck Surface (C7)	Geomorphic Position (D2)
Iron Deposits (B5) Uther (Explain in Remark	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:  Surface Water Present?  Yes No _ ✓ Depth (inches):	
Water Table Present? Yes No _ ✓ Depth (inches):	
Saturation Present?  Yes No _ C Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pre	vious inspections), if available:
Remarks:	
Wetland hydrology indicators are met. Standin	g water was present within microdepressions
within the forested area. The standing water w	as not associated with a high water table, so it
can not be used as an indicator of wetland hyd	rology.

Sampling Point	: SP-UPL-07

Tree Stratum (Plot size: 30 ft r		Dominant		Dominance Test worksheet:	
1. Maclura pomifera	% Cover 30	Species? ✓	FACU	Number of Dominant Species	(4)
2. Celtis laevigata	20		FACW	That Are OBL, FACW, or FAC: 5	(A)
	20		FACU	Total Number of Dominant	
3. Fagus grandifolia				Species Across All Strata: 7	(B)
4				Percent of Dominant Species	
5				That Are OBL, FACW, or FAC: 71.4	(A/B)
6				Prevalence Index worksheet:	
7					
8				Total % Cover of:         Multiply by:           OBL species         60         x 1 = 60	_
	<u>70%</u> :	= Total Cov	er		_
50% of total cover: 35	20% of	total cover:	14	FACW species 35 x 2 = 70	_
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species $30 \times 3 = 90$	=
1. Celtis laevigata	15	<b>✓</b>	FACW	FACU species $50$ $x 4 = 200$	_
2. Ligustrum sinense	15	~	FAC	UPL species $0   x 5 = 0$	_
3.				Column Totals: <u>175</u> (A) <u>420</u>	_ (B)
4.				5	
				Prevalence Index = B/A = 2.4	_
5				Hydrophytic Vegetation Indicators:	
6				1 - Rapid Test for Hydrophytic Vegetation	
7				2 - Dominance Test is >50%	
8				3 - Prevalence Index is ≤3.0¹	
		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain	n)
· · · · · · · · · · · · · · · · · · ·	20% of	total cover	6		
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology m	nust
1. Packera glabella	60		OBL	be present, unless disturbed or problematic.	
2. Rumex crispus	15		FAC	Definitions of Four Vegetation Strata:	
3				Tree Meady plants avaluding vince 2 in (7.0)	\
4				Tree – Woody plants, excluding vines, 3 in. (7.6 of more in diameter at breast height (DBH), regardle	
5.				height.	
6.				Sanling/Shrub Woody plants evaluding vines	logo
				Sapling/Shrub – Woody plants, excluding vines, than 3 in. DBH and greater than 3.28 ft (1 m) tall.	1622
7				, ,	
8				<b>Herb</b> – All herbaceous (non-woody) plants, regar of size, and woody plants less than 3.28 ft tall.	dless
9				of size, and woody plants less than 3.20 it tall.	
10				Woody vine – All woody vines greater than 3.28	ft in
11				height.	
12					
		= Total Cov			
50% of total cover: 38	20% of	total cover	15		
Woody Vine Stratum (Plot size: 30 ft r )					
1					
2					
3.					
4.					
5					
5				Hydrophytic Vegetation	
		Total Cov		Present? Yes No	
50% of total cover:		total cover			
Remarks: (If observed, list morphological adaptations bel	ow).				
Dominance test is met.					
Dominance test is met.					

SOIL Sampling Point: SP-UPL-07

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 4/2	100		<u> </u>			Clay Loam		
									_
-									
				· ——					
	-								
-									
¹Type: C=Cd	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> Location: PL:	=Pore Lining, M=Matr	iy
	Indicators: (Appli					anio.		Problematic Hydric	
Histosol			Polyvalue Be			RR S. T. I		k (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-				Floodplain Soils (F19	
Stratified	l Layers (A5)		Depleted Ma	trix (F3)			Anomalou	s Bright Loamy Soils	(F20)
	Bodies (A6) (LRR I		Redox Dark	Surface (F	<del>-</del> 6)		(MLRA 1	•	
_	cky Mineral (A7) (L		Depleted Dai					nt Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			ow Dark Surface (TF	12)
	ck (A9) (LRR P, T)		Marl (F10) <b>(L</b>		(MI DA 4	E4)	U Other (Exp	olain in Remarks)	
	d Below Dark Surfa ark Surface (A12)	ce (ATT)	Depleted Ocl				T) <sup>3</sup> Indicator	rs of hydrophytic vege	station and
	rairie Redox (A16) (	MLRA 150A)						d hydrology must be p	
	lucky Mineral (S1)		Delta Ochric			, •,		disturbed or problema	
	leyed Matrix (S4)	,,	Reduced Ver			0A, 150B)			
	edox (S5)		Piedmont Flo						
Stripped	Matrix (S6)		Anomalous E	Bright Loar	my Soils (	F20) <b>(MLR</b>	RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,								
Restrictive I	_ayer (if observed	):							
Type:			<u> </u>						
Depth (inc	ches):		<u></u>				Hydric Soil Pre	esent? Yes	No 🔽
Remarks:							1		
No indic	ators are m	et.							
110 1110110									

Project/Site: Golden Triangle Phase 2	ty/County: Lowndes County	Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississi	Sampling Point: SP-UPL-08
Investigator(s): J. Brown, O. Haney	ection, Township, Range: S26 T18N R16	E
	ocal relief (concave, convex, none): Conc	
Subregion (LRR or MLRA): P 135A Lat: 33.399	,	
Soil Map Unit Name: Catalpa silty clay		fication: PFO1A
Are climatic / hydrologic conditions on the site typical for this time of yea		
Are Vegetation, Soil, or Hydrology significantly d		
Are Vegetation, Soil, or Hydrology naturally probability and a second secon		
SUMMARY OF FINDINGS – Attach site map showing		
Hydrophytic Vegetation Present? Yes No _ ✔		
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area	No. V
Wetland Hydrology Present? Yes No	within a Wetland? Yes _	No
Remarks:		
Upland confirmation sample plot. The APT s	nowed normal conditions a	t the time leading up
to and during the field survey.		3
to and during the neid survey.		
HYDROLOGY		
Wetland Hydrology Indicators:	Secondary In-	dicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)		Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)		Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15)		Patterns (B10)
Saturation (A3) Hydrogen Sulfide Oc	or (C1) Moss Trir	n Lines (B16)
Water Marks (B1) — Oxidized Rhizospher	es along Living Roots (C3) 🔲 Dry-Seas	on Water Table (C2)
Sediment Deposits (B2)	<del>-</del>	Burrows (C8)
Drift Deposits (B3)  Recent Iron Reduction		n Visible on Aerial Imagery (C9)
☐ Algal Mat or Crust (B4) ☐ Thin Muck Surface (☐ Iron Deposits (B5) ☐ Other (Explain in Re	<i>'</i>	hic Position (D2) Aquitard (D3)
Inundation Visible on Aerial Imagery (B7)		tral Test (D5)
Water-Stained Leaves (B9)	=	m moss (D8) <b>(LRR T, U)</b>
Field Observations:		· · · · · · · · · · · · · · · · · · ·
Surface Water Present? Yes No Depth (inches):		
Water Table Present? Yes No Depth (inches):		
Saturation Present? Yes No Depth (inches):	Wetland Hydrology Pre	sent? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos	previous inspections), if available:	
Remarks:		
Geomorphic position present. Wetland hydro	ology not met.	

5. Ulmus americana 5

Sapling/Shrub Stratum (Plot size: 15 ft r )

1. Juniperus virginiana

2. Ligustrum sinense

Herb Stratum (Plot size: 5 ft r )

3. Ligustrum sinense

5. Plantago virginica

Woody Vine Stratum (Plot size: 30 ft r )

1. Carex sp.\*

<sub>2.</sub> Poa annua

4 Claytonia virginica

50% of total cover: 38

50% of total cover: 4

Tree Stratum (Plot size: 30 ft r

1. Celtis laevigata

2. Maclura pomifera

3. Juniperus virginiana4. Quercus pagoda

Absolute Dominant Indicator

**FACW** 

FACU

FACU

**FACW** 

% Cover Species? Status

75% = Total Cover

7% = Total Cover

65

25

15

5

50% of total cover: 58 20% of total cover: 23

50% of total cover: \_\_\_\_ 20% of total cover: \_\_\_

20% of total cover: 1

✓ FAC

FACU

FACU

FAC

**FACU** 

\_\_ 20% of total cover: 15

40

5

20 \_\_\_\_

	Sa	ampling Po	int: SP-L	JPL-08
Dominance Te	est workshee	t:		
Number of Don That Are OBL,				(A)
Total Number of Species Across		6		(B)
Percent of Don That Are OBL,				(A/B)
Prevalence Inc	dex workshee	et:		
Total % Co	over of	Multi	ply by:	
OBL species	0	x 1 = 0	<u>⊳., ⊳, </u>	_
FACW species	45	x 2 = 9	0	_
FAC species	87	x 3 = 2	61	_
FACU species	65	x 4 = 2		_
UPL species	0	x 5 = 0		_
Column Totals:	197	(A) 6	11	_ _ (B)
Prevalence Inc	dex = B/A	A = 3.1		_
Hydrophytic V	egetation Inc	dicators:		
	Test for Hydro		etation	
_	ance Test is >	. , .		
	ence Index is ≤			
=	ic Hydrophytic		n <sup>1</sup> (Explai	۵)
i iobicinati		vegetatio	II (Explain	11)
<sup>1</sup> Indicators of h	ydric soil and	wetland hy	/drology m	
<sup>1</sup> Indicators of h	ydric soil and ess disturbed	wetland hy or problen	/drology matic.	
<sup>1</sup> Indicators of h be present, unl	ydric soil and ess disturbed Four Vegetat plants, exclud	wetland hy or problen ion Strata	vdrology m natic. : 3 in. (7.6 d	nust cm) or
<sup>1</sup> Indicators of h be present, unl <b>Definitions of</b> <b>Tree</b> – Woody more in diamet	ydric soil and ess disturbed  Four Vegetat plants, excluder at breast he	wetland hy or problen ion Strata ing vines, eight (DBH	vdrology matic. : 3 in. (7.6 c), regardle	em) or
<sup>1</sup> Indicators of h be present, unl <b>Definitions of</b> <b>Tree</b> – Woody more in diamet height. <b>Sapling/Shrub</b>	ydric soil and ess disturbed Four Vegetat plants, excluder at breast he o – Woody plants and greater the paceous (non-	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) pla	vdrology matic. : 3 in. (7.6 c), regardeling vines, (1 m) tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH Herb – All herb of size, and wo Woody vine –	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH Herb – All herb of size, and wo Woody vine –	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH Herb – All herb of size, and wo Woody vine –	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH Herb – All herb of size, and wo Woody vine –	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl Definitions of Tree – Woody more in diamet height. Sapling/Shrub than 3 in. DBH Herb – All herb of size, and wo Woody vine –	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) plats s than 3.2	odrology matic.  3 in. (7.6 colors), regarded (1 m) tall.  ants, regar 8 ft tall.	em) or ess of less
¹Indicators of h be present, unl  Definitions of  Tree – Woody more in diamet height.  Sapling/Shrub than 3 in. DBH  Herb – All herb of size, and wo	ydric soil and ess disturbed Four Vegetat plants, exclud er at breast he — Woody pland greater the paceous (non-ody plants less	wetland hy or problen ion Strata ing vines, eight (DBH nts, exclude nan 3.28 ft woody) pla s than 3.2 es greater	vdrology matic. : 3 in. (7.6 c), regarded ing vines, (1 m) tall. unts, regar 8 ft tall.	em) or ess of less

Remarks: (If observed, list morphological adaptations below).

No test is met. \*The Carex could not be identified to species. It was assumed to be FAC since most Carex in the area are FAC or wetter.

\_\_\_\_\_= Total Cover

115% = Total Cover

SOIL Sampling Point: SP-UPL-08

	cription: (Describe	to the depth				or confirm	n the absence of	indicators.)
Depth (inches)	Matrix Color (moist)	%	Redox Color (moist)	K Features	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
0 - 18	10YR 3/2		10YR 3/4	2	C	M	Clay Loam	romano
	10111 0/2							
-								
-			_					
	-				-	·	·	
	oncentration, D=De					ains.		=Pore Lining, M=Matrix.
l <u> </u>	Indicators: (Appli	cable to all L						r Problematic Hydric Soils <sup>3</sup> :
Histosol	. ,		Polyvalue Bel		. , .		. –	k (A9) (LRR O)
	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b> Vertic (F18) <b>(outside MLRA 150A,B)</b>
	stic (A3) en Sulfide (A4)		Loamy Mucky Loamy Gleye			R ()	777	Floodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)		Depleted Mat	•	1 2)			us Bright Loamy Soils (F20)
	Bodies (A6) (LRR I	P. T. U)	Redox Dark S	. ,	6)		(MLRA	
· = ·	ıcky Mineral (A7) (L		Depleted Dar	•	,			nt Material (TF2)
_	esence (A8) (LRR		Redox Depre				Very Shal	low Dark Surface (TF12)
1 cm Μι	ıck (A9) <b>(LRR P, T)</b>		Marl (F10) (L	RR U)			U Other (Ex	plain in Remarks)
I <b>=</b> '	d Below Dark Surfa	ce (A11)	Depleted Och				2	
	ark Surface (A12)	(141 D 4 450 A)	Iron-Mangane					ors of hydrophytic vegetation and
_	rairie Redox (A16) (							d hydrology must be present,
_	lucky Mineral (S1) ( Bleyed Matrix (S4)	(LKK U, S)	Delta Ochric ( Reduced Veri					disturbed or problematic.
_	Redox (S5)		Piedmont Flo					
	Matrix (S6)						RA 149A, 153C, 15	53D)
_	rface (S7) (LRR P,	S, T, U)			,	()	,,	,
	Layer (if observed)							
Type:								
Depth (in	ches):						Hydric Soil Pre	esent? Yes 🗸 No
Remarks:	·							
Hydric	oil indicator	F6 is ma	<b>a</b> †					
i iyanc s	on malcator	1 0 13 1110						

Project/Site: Golden Triangle Phase 2	ty/County: Lowndes County Sampling Date: 2021-03-08
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-01
• • • • • • • • • • • • • • • • • • • •	ection, Township, Range: S28 T18N R16E
	pocal relief (concave, convex, none): Concave Slope (%): 2
	573 Long: -88.620082 Datum: NAD 83
Soil Map Unit Name: Leeper silty clay, 0 to 2 percent slopes, oc	ccasionally flooded NW/ classification: N/A
Are climatic / hydrologic conditions on the site typical for this time of year'	
Are Vegetation, Soil, or Hydrology significantly dis	
Are Vegetation, Soil, or Hydrology naturally problem.	ematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing s	campling point locations, transects, important features, etc.
Livedison budies Veneratedison December 2	
Hydrophytic Vegetation Present? Yes V No No No No	Is the Sampled Area
Wetland Hydrology Present? Yes   ✓ No	within a Wetland? Yes No
Remarks:	
Sample plot in W-DFM-1. The APT showed no	ormal conditions at the time leading up to and
	offilial conditions at the time leading up to and
during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15) (	LRR U) Prainage Patterns (B10)
Saturation (A3) Hydrogen Sulfide Odd	- ` ' · · · · · · · · · <del></del> · · · · · · · · · · · · · · · · · ·
	es along Living Roots (C3)
Sediment Deposits (B2)	<del>_</del> '
☐ Drift Deposits (B3) ☐ Recent Iron Reduction	
Algal Mat or Crust (B4)  Thin Muck Surface (C	· · · · · · · · · · · · · · · · · · ·
☐ Iron Deposits (B5) ☐ Other (Explain in Rem	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	2
Surface Water Present? Yes No Depth (inches): 2  Water Table Present? Yes No Depth (inches): 2	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos,	previous inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	

<b>'EGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PEM-01
7 0 4 (D) ( 20 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1		Species?	Status	Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8		Total Cov		OBL species <u>35</u> x 1 = <u>35</u>
50% of total cover:				FACW species <u>50</u> x 2 = <u>100</u>
Sapling/Shrub Stratum (Plot size: 30 ft r			·	FAC species <u>0</u> x 3 = <u>0</u>
1				FACU species $0 \times 4 = 0$
2.				UPL species $0 \times 5 = 0$
3.				Column Totals: <u>85</u> (A) <u>135</u> (B)
4				Prevalence Index = B/A = <u>1.6</u>
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				<b>2</b> 3 - Prevalence Index is ≤3.0 <sup>1</sup>
		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 30 ft r )  1 Andropogon glomeratus	40	V	FACW	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Alternanthera philoxeroides	35		OBL	
2. Lysimachia nummularia	10		FACW	Definitions of Four Vegetation Strata:
A				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
r				more in diameter at breast height (DBH), regardless of height.
5				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				Herb – All herbaceous (non-woody) plants, regardless
9.				of size, and woody plants less than 3.28 ft tall.
10.				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	85% :	= Total Cov	ver .	
50% of total cover: 43	20% of	total cover	<u> 17</u>	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic Vegetation
E09/ of total cover:		= Total Cov		Present? Yes No
50% of total cover: Remarks: (If observed, list morphological adaptations bel		lotal cover	•	
, ,	OW).			
Rapid test is met.				

SOIL Sampling Point: W-PEM-01

Profile Desc	cription: (Describe to the depti	needed to docu	ment the	indicator of	or confirm	the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
_							
·						· · · · · · · · · · · · · · · · · · ·	
						<u> </u>	
-							
			_				_
<sup>1</sup> Type: C=C	oncentration, D=Depletion, RM=I	Reduced Matrix, Ma	S=Maske	d Sand Gra	ains.	<sup>2</sup> Location: PL=F	Pore Lining, M=Matrix.
	Indicators: (Applicable to all L						roblematic Hydric Soils <sup>3</sup> :
Histosol	(A1)	Polyvalue Be	elow Surfa	ace (S8) <b>(L</b>	RR S. T. L	J) 🔲 1 cm Muck (	(A9) (LRR O)
I —	pipedon (A2)	Thin Dark Su					(A10) <b>(LRR S)</b>
_	istic (A3)	Loamy Muck					ertic (F18) (outside MLRA 150A,B)
	en Sulfide (A4)	Loamy Gleye	-	. , .	σ,		oodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	Depleted Ma		()			Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark		F6)		(MLRA 15	• • • • •
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da				_ `	Material (TF2)
_	resence (A8) (LRR U)	Redox Depre					w Dark Surface (TF12)
_	uck (A9) (LRR P, T)	Marl (F10) (L		٠,			ain in Remarks)
_	d Below Dark Surface (A11)	Depleted Oc		) (MLRA 15	51)	Culci (Expid	
	ark Surface (A12)	Iron-Mangan				T) <sup>3</sup> Indicators	of hydrophytic vegetation and
_	rairie Redox (A16) (MLRA 150A)	=				•	hydrology must be present,
	Mucky Mineral (S1) (LRR O, S)	Delta Ochric			, -,		sturbed or problematic.
	Gleyed Matrix (S4)	Reduced Ve			0A 150B)		otarboa or problematic.
	Redox (S5)	☐ Piedmont Flo					
	d Matrix (S6)					A 149A, 153C, 153I	וח
	urface (S7) (LRR P, S, T, U)	/tilomalous t	ongni Loc	arriy Cono (i	20) (IIILIX	A 140A, 1000, 1001	2,
	Layer (if observed):						
Type:							
, , <u> </u>							
	ches):					Hydric Soil Pres	ent? Yes No No
Remarks:							
Soil san	nple was not taken d	ue to inunda	ation.				
	•						

Project/Site: Golden Triangle Phase 2 Cit	ty/County: Lowndes County	Sampling Date: 2021-03-08
Applicant/Owner: TVA		Sampling Point: W-PEM-01_UPL
• • • • • • • • • • • • • • • • • • • •	ection, Township, Range: S26 T18N R16E	
Landform (hillslope, terrace, etc.): Hillslope Lo		Slone (%)· 5
Subregion (LRR or MLRA): P 135A Lat: 33.3986		
Soil Map Unit Name: Leeper silty clay, 0 to 2 percent slopes, oc		
Are climatic / hydrologic conditions on the site typical for this time of year?		·
Are Vegetation, Soil, or Hydrology significantly dis		
Are Vegetation, Soil, or Hydrology naturally proble	ematic? (If needed, explain any answe	ers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing s	ampling point locations, transects	s, important features, etc.
Hydrophytic Vegetation Present?  Hydric Soil Present?  Wetland Hydrology Present?  Remarks:  Yes No  Yes No	Is the Sampled Area within a Wetland? Yes	No
	o ADT showed normal cond	itions at the time
Upland sample plot adjacent to W-PEM-1. Th	ie APT Showed normal cond	itions at the time
leading up to and during the field survey.		
HYDROLOGY		
Wetland Hydrology Indicators:		ators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)		Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)		getated Concave Surface (B8)
High Water Table (A2)  High Water Table (A2)  High Water Table (A2)		
☐ Saturation (A3) ☐ Hydrogen Sulfide Odd ☐ Water Marks (B1) ☐ Oxidized Rhizosphere	· · ·	Water Table (C2)
Sediment Deposits (B2)  Sediment Deposits (B2)  Presence of Reduced		
Drift Deposits (B3)  Recent Iron Reduction	· · · · · <del>_</del> · ·	isible on Aerial Imagery (C9)
Algal Mat or Crust (B4) Thin Muck Surface (C		Position (D2)
☐ Iron Deposits (B5) ☐ Other (Explain in Rem	narks) Shallow Aqu	itard (D3)
☐ Inundation Visible on Aerial Imagery (B7)	FAC-Neutra	Test (D5)
Water-Stained Leaves (B9)	Sphagnum r	moss (D8) <b>(LRR T, U)</b>
Field Observations:		
Surface Water Present? Yes No Depth (inches): _		
Water Table Present? Yes No V Depth (inches):		
Saturation Present? Yes No Depth (inches): _ (includes capillary fringe)	Wetland Hydrology Prese	nt? Yes No
Describe Recorded Data (stream gauge, monitoring well, aerial photos,	previous inspections), if available:	
Remarks:		
No indicators are met.		

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 1 (B)
4				Dancart of Dancie and On a de-
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B)
6.				That Ale OBE, I AOW, OI I AO.
7				Prevalence Index worksheet:
				Total % Cover of: Multiply by:
8				OBL species <u>0</u> x 1 = <u>0</u>
		= Total Cov		FACW species $0   x 2 = 0$
50% of total cover:	20% of	total cover	:	FAC species 40 x 3 = 120
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 0 x 4 = 0
1				
2				UPL species $0 \times 5 = 0$
3				Column Totals: <u>40</u> (A) <u>120</u> (B)
4.				Prevalence Index = B/A = 3.0
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0¹
		= Total Cov	/er	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Setaria pumila	40	<b>✓</b>	FAC	be present, unless disturbed or problematic.
2.				Definitions of Four Vegetation Strata:
3.				
				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4				more in diameter at breast height (DBH), regardless of height.
5				g.m
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				<b>Herb</b> – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Weedy vine All woody vines greater than 2.29 ft in
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12.				g.m
12.	40%	- Total Co	·or	
50% of total cover: 20				
	20% 01	total cover		
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5.				Hydrophytic
		Total Cov	/er	Vegetation
E09/ of total cover:				Present? Yes No
50% of total cover:		total cover	•	
Remarks: (If observed, list morphological adaptations below	OW).			
Dominance test is met.				

Sampling Point: W-PEM-01\_UPL

SOIL Sampling Point: W-PEM-01\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	indicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 18	10YR 3/3	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Co	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masked	Sand Gr	ains	<sup>2</sup> l ocation: Pl	=Pore Lining, M=Matri	<u> </u>
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RR S. T. I		k (A9) <b>(LRR O)</b>	
_	oipedon (A2)		Thin Dark Su				. –	k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside N	MLRA 150A,B)
Hydroge	n Sulfide (A4)		Loamy Gleye	ed Matrix (	(F2)		Piedmont	Floodplain Soils (F19)	(LRR P, S, T)
	l Layers (A5)		Depleted Ma	trix (F3)			Anomalou	s Bright Loamy Soils (	F20)
	Bodies (A6) (LRR I		Redox Dark		•		(MLRA	,	
_	cky Mineral (A7) (L		Depleted Dai					nt Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			low Dark Surface (TF1	2)
	ick (A9) <b>(LRR P, T)</b> d Below Dark Surfa		Marl (F10) <b>(L</b> Depleted Ocl		/MI DA 1	<b>5</b> 1\	U Otner (Exp	plain in Remarks)	
	ark Surface (A12)	CC (ATT)	Iron-Mangan				T) <sup>3</sup> Indicator	rs of hydrophytic veget	tation and
	rairie Redox (A16) (	MLRA 150A)						d hydrology must be pr	
	lucky Mineral (S1)		Delta Ochric			. ,		disturbed or problema	
	Bleyed Matrix (S4)	, ,	Reduced Ver			60A, 150B)		·	
☐ Sandy R	ledox (S5)		Piedmont Flo	odplain S	ioils (F19)	(MLRA 14	49A)		
	Matrix (S6)		Anomalous E	Bright Loai	my Soils (	F20) <b>(MLR</b>	RA 149A, 153C, 15	33D)	
	rface (S7) (LRR P,						ľ		
Restrictive I	_ayer (if observed)	):							
Type:			_						
Depth (inc	ches):						Hydric Soil Pre	esent? Yes	No
Remarks:									
No indic	ators are me	et.							

Project/Site: Golden Triangle Phase 2	City/County: Lowndes County Sampling Date: 2021-0	3-11				
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-02					
Investigator(s): J. Brown, O. Haney Section, Township, Range: S26 T18N R16E						
Landform (hillslope, terrace, etc.): Basin	Local relief (concave, convex, none): Concave Slope (%): 2	2				
Subregion (LRR or MLRA): P 135A Lat: 33.3						
Soil Map Unit Name: Catalpa silty clay	NWI classification: NA					
Are climatic / hydrologic conditions on the site typical for this time of y						
	ly disturbed? Are "Normal Circumstances" present? YesNo					
Are Vegetation, Soil, or Hydrology naturally p	oroblematic? (If needed, explain any answers in Remarks.)					
SUMMARY OF FINDINGS - Attach site map showing	g sampling point locations, transects, important features	s, etc.				
Hydrophytic Vegetation Present? Yes No						
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	is the campion Area					
Wetland Hydrology Present? Yes   ✓ No						
Remarks:	<u>-</u>					
Sample plot in W-PFM-02 The APT shows	ed normal conditions at the time leading up to a	nd				
	sa normal conditions at the time leading up to a	iiiu				
during the field survey.						
HYDROLOGY						
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two requ	uired)				
Primary Indicators (minimum of one is required; check all that apply	Surface Soil Cracks (B6)					
Surface Water (A1) Aquatic Fauna (B	Sparsely Vegetated Concave Surface	(B8)				
High Water Table (A2)  Marl Deposits (B <sup>2</sup>	15) (LRR U)					
Saturation (A3) Hydrogen Sulfide	<del></del>					
	pheres along Living Roots (C3)					
Sediment Deposits (B2)						
	uction in Tilled Soils (C6)	(9)				
Algal Mat or Crust (B4)	<u> </u>					
☐ Iron Deposits (B5) ☐ Other (Explain in						
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)					
Water-Stained Leaves (B9)	☐ Sphagnum moss (D8) (LRR T, U)					
Field Observations:  Surface Water Present?  Yes No Depth (inche	20): 1					
Water Table Present? Yes V No Depth (inche						
Saturation Present? Yes V No Depth (inche						
(includes capillary fringe)						
Describe Recorded Data (stream gauge, monitoring well, aerial pho	tos, previous inspections), if available:					
Remarks:						
Wetland hydrology indicators are met.						

<b>EGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PEM-02
Total Objections (Distriction 30 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1.		Species?		Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species <u>75</u> x 1 = <u>75</u>
50% of total cover:		= Total Cov		FACW species <u>0</u> x 2 = <u>0</u>
Sapling/Shrub Stratum (Plot size: 15 ft r )	20 /6 01	total cover	•	FAC species <u>40</u> x 3 = <u>120</u>
1				FACU species <u>0</u>
2.				UPL species <u>0</u> x 5 = <u>0</u>
3.				Column Totals: 115 (A) 195 (B)
4				Prevalence Index = B/A = 1.7
5.				Hydrophytic Vegetation Indicators:
6.				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				☑ 3 - Prevalence Index is ≤3.0¹
	=	= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Salix nigra	60		OBL	be present, unless disturbed or problematic.
2. Ambrosia trifida	30		FAC	Definitions of Four Vegetation Strata:
3. Eleocharis palustris	- <u>15</u> 10		OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Cyperus rotundus			FAC	more in diameter at breast height (DBH), regardless of height.
5				
6				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				
8			-	<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
			-	
10 11				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12.				noight.
	115% :	= Total Co	ver	
50% of total cover: 58	20% of	total cover	: 23	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
		= Total Co		Vegetation Present?  Yes No
50% of total cover:		total cover	":	
Remarks: (If observed, list morphological adaptations bel	ow).			
Dominance test is met.				

SOIL Sampling Point: W-PEM-02

Profile Desc	ription: (Describe	to the depti	n needed to docum	nent the i	ndicator	or confir	m the absence of i	ndicators.)
Depth	Matrix	0/		x Feature		1 2	Tantona	Developing
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	Texture	Remarks
0 - 18	10YR 3/2	60	10YR 3/6	40	<u>C</u>	M	Clay Loam	
							- <u></u>	
-								
							·	
		<del></del>						
	-						<del></del>	
							·	
	oncentration, D=De					ains.		=Pore Lining, M=Matrix.
l <u> </u>	Indicators: (Appli	cable to all L						Problematic Hydric Soils <sup>3</sup> :
Histosol			Polyvalue Be				. –	(A9) (LRR O)
Histic Ep	oipedon (A2)		Thin Dark Su Loamy Muck					k (A10) (LRR S) Vertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)		Loamy Gleye			(0)		Floodplain Soils (F19) (LRR P, S, T)
	Layers (A5)		Depleted Mat		/			s Bright Loamy Soils (F20)
	Bodies (A6) (LRR I	P, T, U)	Redox Dark		<del>-</del> 6)		(MLRA 1	
5 cm Mu	ıcky Mineral (A7) <b>(L</b>	RR P, T, U)	Depleted Dar	k Surface	(F7)		Red Paren	nt Material (TF2)
	esence (A8) (LRR		Redox Depre		8)			ow Dark Surface (TF12)
	ick (A9) (LRR P, T)		☐ Marl (F10) <b>(L</b>			-40	U Other (Exp	olain in Remarks)
_	d Below Dark Surfa ark Surface (A12)	ce (A11)	Depleted Och				T) <sup>3</sup> Indicator	rs of hydrophytic vegetation and
	rairie Redox (A16) (	MLRA 150A				•	•	d hydrology must be present,
	lucky Mineral (S1)		Delta Ochric					disturbed or problematic.
	Bleyed Matrix (S4)	, ,	Reduced Ver					·
	ledox (S5)		Piedmont Flo					
	Matrix (S6)		Anomalous B	Bright Loa	my Soils	(F20) <b>(MLI</b>	RA 149A, 153C, 15	3D)
	rface (S7) (LRR P,							
	_ayer (if observed	):						
Type:	ahaa):						Hydric Soil Pre	esent? Yes V No No
Remarks:	ches):						nyunc 3011 Fre	sent? resNo
	-:::::::::::::::::::::::::::::::::::::	FC :						
Hyaric s	oil indicator	F6 IS M	et.					

Project/Site: Golden Triangle Phase 2	City/County: Low	ndes County	Sampling Date: 2021-03-11
Applicant/Owner: TVA			Sampling Point: W-PEM-02_UPL
		s26 T18N R16E	· · · · · · · · · · · · · · · · · · ·
Landform (hillslope, terrace, etc.): Upland, Flat	Local relief (conca	ve, convex, none): None	Slope (%): 0
Subregion (LRR or MLRA): P 135A La	ıt: 33.396092	Long: -88.598834	Datum: NAD 83
Soil Map Unit Name: Catalpa silty clay			tion: NA
Are climatic / hydrologic conditions on the site typical for this	time of year? YesN	lo (If no, explain in R	emarks.)
Are Vegetation, Soil, or Hydrology si			
Are Vegetation, Soil, or Hydrology na		(If needed, explain any answe	
SUMMARY OF FINDINGS – Attach site map s			
		·	•
Hydrophytic Vegetation Present? Yes No Hydric Soil Present? Yes No Wetland Hydrology Present? Yes No Hydrology Present?	Is the Sam	•	
Wetland Hydrology Present? Yes No	within a W	etland? Yes	No
Remarks:	<u> </u>		
Upland sample plot adjacent to W-PE	- -M-02. The area v	vas a maintained pa	asture area. The
APT showed normal conditions at the		•	
AFT Showed normal conditions at the	s time leading up t	to and during the h	eiu sui vey.
HYDROLOGY			
Wetland Hydrology Indicators:		Secondary Indica	tors (minimum of two required)
	act apply)	_	
Primary Indicators (minimum of one is required; check all the		Surface Soil	
	Fauna (B13)		getated Concave Surface (B8)
	posits (B15) (LRR U)	☐ Drainage Pat	
	n Sulfide Odor (C1)	Moss Trim Li	
	Rhizospheres along Living F		Water Table (C2)
	e of Reduced Iron (C4)	Crayfish Buri	,
	ron Reduction in Tilled Soils ( ck Surface (C7)		sible on Aerial Imagery (C9)
	xplain in Remarks)	Shallow Aqui	Position (D2)
Inundation Visible on Aerial Imagery (B7)	kpiaiii iii Neiliaiks)	FAC-Neutral	
Water-Stained Leaves (B9)		=	noss (D8) <b>(LRR T, U)</b>
Field Observations:		Opilagilalii ii	1033 (D0) (ERR 1, 0)
Surface Water Present? Yes No Dep	th (inches)		
Water Table Present? Yes No Dep			
Saturation Present? Yes No V Dep		Wetland Hydrology Presen	t? Yes No
(includes capillary fringe)	, , ,		
Describe Recorded Data (stream gauge, monitoring well, a	eriai pnotos, previous inspec	tions), if available:	
Remarks:			
No indicators are met.			

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r	% Cover	Species?	Status	Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 1 (B)
4				
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B)
6				That Ald OBE, I AOW, OF I AO.
7.				Prevalence Index worksheet:
8.				Total % Cover of: Multiply by:
0		Total Cov	·or	OBL species <u>0</u> x 1 = <u>0</u>
50% of total cover:				FACW species <u>0</u> x 2 = <u>0</u>
	20% 01	total cover		FAC species <u>75</u> x 3 = <u>225</u>
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 25
1				UPL species $0$ $x = 0$
2				100 005
3				Column Totals: 100 (A) 325 (B)
4				Prevalence Index = B/A = 3.25
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7.				2 - Dominance Test is >50%
8.				
<u> </u>		Total Cov	/er	☐ 3 - Prevalence Index is ≤3.0¹
50% of total cover:				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
· · · · · · · · · · · · · · · · · · ·	20% 01	lotal cover		
Herb Stratum (Plot size: 5 ft r )  1 Schedonorus arundinaceus	70		FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must
··	· ——			be present, unless disturbed or problematic.
2. Schizachyrium scoparium	15		FACU	Definitions of Four Vegetation Strata:
3. Allium vineale	5		FACU	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Ligustrum sinense	5		FAC	more in diameter at breast height (DBH), regardless of
5. Rosa multiflora	5		FACU	height.
6	-			Sapling/Shrub – Woody plants, excluding vines, less
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
9				of size, and woody plants less than 3.20 it tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	100% =			
50% of total cover: 50	20% of	total cover	<u> 20 </u>	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3.				
4.				
5				
5				Hydrophytic Vegetation
500/ 51 1 1		Total Cov		Present? Yes No
50% of total cover:		total cover		
Remarks: (If observed, list morphological adaptations believed)	ow).			
Dominance test is met.				

Sampling Point: W-PEM-02\_UPL

SOIL Sampling Point: W-PEM-02\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	indicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 18	10YR 3/3	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Co	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masked	Sand Gr	ains	<sup>2</sup> l ocation: Pl	=Pore Lining, M=Matri	<u> </u>
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RR S. T. I		k (A9) <b>(LRR O)</b>	
_	oipedon (A2)		Thin Dark Su				. –	k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside N	MLRA 150A,B)
Hydroge	n Sulfide (A4)		Loamy Gleye	ed Matrix (	(F2)		Piedmont	Floodplain Soils (F19)	(LRR P, S, T)
	l Layers (A5)		Depleted Ma	trix (F3)			Anomalou	s Bright Loamy Soils (	F20)
	Bodies (A6) (LRR I		Redox Dark		•		(MLRA	,	
_	cky Mineral (A7) (L		Depleted Dai					nt Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			low Dark Surface (TF1	2)
	ick (A9) <b>(LRR P, T)</b> d Below Dark Surfa		Marl (F10) <b>(L</b> Depleted Ocl		/MI DA 1	<b>5</b> 1\	U Otner (Exp	plain in Remarks)	
	ark Surface (A12)	CC (ATT)	Iron-Mangan				T) <sup>3</sup> Indicator	rs of hydrophytic veget	tation and
	rairie Redox (A16) (	MLRA 150A)						d hydrology must be pr	
	lucky Mineral (S1)		Delta Ochric			. ,		disturbed or problema	
	Bleyed Matrix (S4)	, ,	Reduced Ver			60A, 150B)		·	
☐ Sandy R	ledox (S5)		Piedmont Flo	odplain S	ioils (F19)	(MLRA 14	49A)		
	Matrix (S6)		Anomalous E	Bright Loai	my Soils (	F20) <b>(MLR</b>	RA 149A, 153C, 15	33D)	
	rface (S7) (LRR P,						ľ		
Restrictive I	_ayer (if observed)	):							
Type:			_						
Depth (inc	ches):						Hydric Soil Pre	esent? Yes	No
Remarks:									
No indic	ators are me	et.							

Project/Site: Golden Triangle	Phase 2	Citv/C	ounty: Lowndes C	ounty	Sampling Date: 2021-03-10
Applicant/Owner: TVA					Sampling Point: W-PEM-03
Investigator(s): J. Brown, O. H	laney	Section	on Townshin Range		Company Cont.
Landform (hillslope, terrace, etc.)	· Basin	Local	rollof (concavo, conv	ov nono): Concave	Slope (%): 2
Subregion (LRR or MLRA): P 13		Lat: 33.385592			Datum: NAD 83
		Lat:	-		
Soil Map Unit Name: Catalpa s				NWI classifica	
Are climatic / hydrologic condition					_
Are Vegetation, Soil	, or Hydrology	significantly distur	bed? Are "Nor	mal Circumstances" p	resent? YesNo
Are Vegetation, Soil	, or Hydrology	naturally problema	atic? (If neede	d, explain any answe	rs in Remarks.)
SUMMARY OF FINDINGS	5 – Attach site m	ap showing sam	pling point loca	tions, transects	, important features, etc.
Hydrophytic Vegetation Present	r? Yes ✔	_ No			
Hydric Soil Present?	Yes 🗸	No	Is the Sampled Are		No
Wetland Hydrology Present?		No	within a Wetland?	res	NO
Remarks:					
Sample plot in W-PE	M-03. The AF	PT showed no	rmal condition	ons at the time	e leading up to and
during the field surv					<b>5</b> 1
daring the held salv	Cy.				
LIVEROLOGY					
HYDROLOGY				Casandan Indiaa	Acre (reining) of the manning di
Wetland Hydrology Indicators		( all that apply)		_	tors (minimum of two required)
Primary Indicators (minimum of Surface Water (A1)				Surface Soil	
<del>                                    </del>		iatic Fauna (B13) 1 Deposits (B15) <b>(LRF</b>	D 11\		getated Concave Surface (B8)
High Water Table (A2) Saturation (A3)		r Deposits (B15) <b>(EK)</b> Irogen Sulfide Odor ((		☐ Drainage Pat ☐ Moss Trim Li	
Water Marks (B1)		dized Rhizospheres a			Water Table (C2)
Sediment Deposits (B2)		sence of Reduced Iro		Crayfish Burr	
Drift Deposits (B3)		cent Iron Reduction in		=,	sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)		n Muck Surface (C7)	()	Geomorphic	
Iron Deposits (B5)		er (Explain in Remark	s)	Shallow Aqui	, ,
Inundation Visible on Aerial	<del></del>	` '	,	FAC-Neutral	` '
Water-Stained Leaves (B9)	)			=	noss (D8) <b>(LRR T, U)</b>
Field Observations:					
Surface Water Present?	Yes No	Depth (inches): 1			
Water Table Present?	Yes No	Depth (inches): 0			
	Yes No	Depth (inches): 0	Wetlan	d Hydrology Presen	t? Yes <u>/</u> No
(includes capillary fringe)  Describe Recorded Data (strear	m gauge, monitoring w	vell, aerial photos, pre	vious inspections), if	available:	
,			. ,		
Remarks:					
Wetland hydrology i	ndicators are	met.			

<b>EGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PEM-03
7. O. 1. (D. 1.) 20 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1.		Species?		Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 115 x 1 = 115
		= Total Cov		FACW species 0 x 2 = 0
50% of total cover: Sapling/Shrub Stratum (Plot size: 15 ft r )	20% 01	total cover		FAC species <u>5</u> x 3 = <u>15</u>
				FACU species $0   x4 = 0$
1				UPL species <u>0</u> x 5 = <u>0</u>
3				Column Totals: <u>120</u> (A) <u>130</u> (B)
4				Prevalence Index = B/A = 1.08
5.				Hydrophytic Vegetation Indicators:
6.				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8.				2 - Bornmande restris > 50 %  ✓ 3 - Prevalence Index is ≤3.0¹
	=	= Total Cov	ver	Problematic Hydrophytic Vegetation¹ (Explain)
50% of total cover:	20% of	total cover	:	<u> </u>
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Salix nigra	50		OBL	be present, unless disturbed or problematic.
2. Carex lurida	30		OBL	Definitions of Four Vegetation Strata:
3. Leersia oryzoides	20	-	OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Alternanthera philoxeroides	10		OBL	more in diameter at breast height (DBH), regardless of
5. Packera glabella	- <del>5</del>		OBL	height.
6. Xanthium strumarium			FAC	Sapling/Shrub – Woody plants, excluding vines, less
7		-	-	than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
		= Total Cov		
50% of total cover: 60	20% of	total cover	: 24	
Woody Vine Stratum (Plot size: 30 ft r )  1.				
2.				
3				
4				
5				Hydrophytic
	=	= Total Cov	ver	Vegetation Present?  Yes No
50% of total cover:	20% of	total cover	:	Present? Yes No
Remarks: (If observed, list morphological adaptations bel	ow).			
Rapid test is met.				

SOIL Sampling Point: W-PEM-03

Profile Desc	ription: (Describe	to the depti	n needed to docun	nent the i	indicator	or confirm	the absence	of indicators.)
Depth	Matrix			x Feature		. 2	<b>-</b> .	
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0 - 18	10YR 3/1	60	10YR 3/6	40	<u>C</u>	PL / M	Clay Loam	
-								
				-				
	-							
				. <u></u>				
<sup>1</sup> Type: C=Co	oncentration, D=De	pletion, RM=I	Reduced Matrix, MS	S=Masked	d Sand G	rains.	<sup>2</sup> Location:	PL=Pore Lining, M=Matrix.
Hydric Soil	Indicators: (Appli	cable to all L	RRs, unless other	wise not	ed.)		Indicators	for Problematic Hydric Soils <sup>3</sup> :
Histosol	(A1)		Polyvalue Be	low Surfa	ce (S8) <b>(</b> I	LRR S, T, L	J) 🛄 1 cm M	uck (A9) (LRR O)
Histic Ep	oipedon (A2)		Thin Dark Su					uck (A10) <b>(LRR S)</b>
Black Hi			Loamy Mucky			R O)		ed Vertic (F18) (outside MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye		(F2)			ont Floodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	. T III	Depleted Mat		-0)			lous Bright Loamy Soils (F20)
	Bodies (A6) (LRR Incky Mineral (A7) (L		Redox Dark S Depleted Dar	•			,	A 153B)  Irent Material (TF2)
_	esence (A8) <b>(LRR</b> I		Redox Depre					nallow Dark Surface (TF12)
	ick (A9) (LRR P, T)		☐ Marl (F10) <b>(L</b>		0)			Explain in Remarks)
	d Below Dark Surfa		Depleted Och		(MLRA 1	51)	_ `	,
	ark Surface (A12)		Iron-Mangane	ese Mass	es (F12)	(LRR O, P,	T) <sup>3</sup> Indica	ators of hydrophytic vegetation and
	rairie Redox (A16) (							and hydrology must be present,
	flucky Mineral (S1)	(LRR O, S)	Delta Ochric					ss disturbed or problematic.
	Gleyed Matrix (S4)		Reduced Ver					
	ledox (S5) Matrix (S6)		Piedmont Flo				эд) А 149A, 153C,	153D)
	rface (S7) <b>(LRR P</b> ,	S. T. U)	Anomalous B	iligili Luai	illy Solis i	(1 20) <b>(WLK</b>	A 149A, 133C,	1330)
	_ayer (if observed)							
Type:								
, ,  —	ches):						Hydric Soil I	Present? Yes V No No
Remarks:			<del></del>				, , , , , ,	
	oil indicator	c 11 and	NEG ara mat					
nyunc s	oil indicator	5 A4 and	a ro are met	•				

Project/Site: Golden Triangle Phase 2	City/County: Lowndes Cour	nty	Sampling Date: 2021-03-10
Applicant/Owner: TVA	· · · · · · · · · · · · · · · · · · ·	State: Mississippi	Sampling Point: W-PEM-03_UPL
	Section, Township, Range: S3		
	Local relief (concave, convex, r		Slone (%): 0
Subregion (LRR or MLRA): P 135A Lat: 33.38	•		Datum: NAD 83
Soil Map Unit Name: Catalpa silty clay	Long	NWI classifica	
Are climatic / hydrologic conditions on the site typical for this time of ye			
Are Vegetation, Soil, or Hydrology significantly			
Are Vegetation, Soil, or Hydrology naturally pr	blematic? (If needed, e.	xplain any answer	rs in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing	sampling point locatio	ns, transects	, important features, etc.
Hydrophytic Vegetation Present? Yes No			
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area		
Wetland Hydrology Present? Yes No _✓	within a Wetland?	Yes	No
Remarks:			
Upland sample plot adjacent to W-PEM-03	The APT showed r	normal cond	ditions at the time
	1110711 1 011011041	iormar com	antionio de tino timo
leading up to and during the field survey.			
HYDROLOGY			
Wetland Hydrology Indicators:			tors (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)		Surface Soil (	Cracks (B6)
Surface Water (A1)			etated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B1)		Drainage Pat	
Saturation (A3)  Hydrogen Sulfide (  Ovidinal Philosoph	` '	Moss Trim Li	` '
☐ Water Marks (B1)       ☐ Oxidized Rhizosph         ☐ Sediment Deposits (B2)       ☐ Presence of Reduction	eres along Living Roots (C3)	Crayfish Burr	Nater Table (C2)
	ion in Tilled Soils (C6)	= 1	sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)  Algal Mat or Crust (B4)  Thin Muck Surface		Geomorphic	
☐ Iron Deposits (B5) ☐ Other (Explain in F	` '	Shallow Aqui	` ,
Inundation Visible on Aerial Imagery (B7)	,	FAC-Neutral	Test (D5)
☐ Water-Stained Leaves (B9)		Sphagnum m	oss (D8) (LRR T, U)
Field Observations:			
Surface Water Present? Yes No Depth (inches			
Water Table Present? Yes No Depth (inches			
Saturation Present? Yes No Depth (inches (includes capillary fringe)	: Wetland H	ydrology Presen	t? Yes No
Describe Recorded Data (stream gauge, monitoring well, aerial phot	s, previous inspections), if avai	ilable:	
Remarks:			
No indicators are met.			

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r	% Cover			Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 3 (B)
4				Descrit of Descinant Conscion
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 33.3 (A/B)
6				(VB)
7				Prevalence Index worksheet:
8.				Total % Cover of: Multiply by:
		= Total Co	ver	OBL species <u>0</u> x 1 = <u>0</u>
50% of total cover:				FACW species 40 x 2 = 80
4F ft	20 /0 01	total cover		FAC species 15 x 3 = 45
				FACU species <u>50</u> x 4 = <u>200</u>
1				UPL species 30 x 5 = 150
2				Column Totals: 135 (A) 475 (B)
3				(1)
4				Prevalence Index = B/A = 3.52
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
	=	= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	<u> </u>
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Cynodon dactylon	50	~	FACU	be present, unless disturbed or problematic.
2. Ranunculus abortivus	40		FACW	Definitions of Four Vegetation Strata:
3. Setaria faberi	30		UPL	John Money Co. 1 Car. 1 Cagottation Carata.
4. Schedonorus arundinaceus	15		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
			1710	more in diameter at breast height (DBH), regardless of height.
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	135% :	= Total Co	ver	
50% of total cover: 68	20% of	total cover	<sub>:</sub> 27	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4			-	
5				Hydrophytic
		Total Co	ver	Vegetation Present? Yes No
50% of total cover:	20% of	total cover	:	riesent: res No
Remarks: (If observed, list morphological adaptations believed)	ow).			
No test is met.				
140 test is met.				

Sampling Point: W-PEM-03\_UPL

SOIL Sampling Point: W-PEM-03\_UPL

Profile Desc	ription: (Describe	to the depth	n needed to docun	nent the i	ndicator	or confirm	the absence of inc	dicators.)
Depth	Matrix			x Feature			_	_
(inches)	Color (moist)	<u> </u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0 - 20	10YR 3/2	95	10YR 3/6	5	С	М	Clay Loam	
	-				-			
-								
_	-							
·	-			· ——				
								•
	oncentration, D=Dep					ains.		Pore Lining, M=Matrix.
Hydric Soil	Indicators: (Applic	able to all L						roblematic Hydric Soils³:
Histosol			Polyvalue Be				. –	
	pipedon (A2)		Thin Dark Su					(A10) (LRR S)
Black Hi			Loamy Mucky			R O)		ertic (F18) <b>(outside MLRA 150A,B)</b> oodplain Soils (F19) <b>(LRR P, S, T)</b>
_	n Sulfide (A4) I Layers (A5)		Loamy Gleye Depleted Mat		F2)			Bright Loamy Soils (F20)
	Bodies (A6) <b>(LRR F</b>	P. T. U)	Redox Dark		6)		(MLRA 15	
I = 1	cky Mineral (A7) (L		Depleted Dar	•	•			Material (TF2)
_	esence (A8) (LRR l		Redox Depre					w Dark Surface (TF12)
1 cm Mu	ck (A9) (LRR P, T)		Marl (F10) <b>(L</b>				U Other (Expla	ain in Remarks)
	Below Dark Surfac	e (A11)	Depleted Och					
=	ark Surface (A12)	MI DA 450A\	Iron-Mangan				•	of hydrophytic vegetation and
	airie Redox (A16) <b>(</b> lucky Mineral (S1) <b>(</b>		Umbric Surfa Delta Ochric			, 0)		hydrology must be present, isturbed or problematic.
	Bleyed Matrix (S4)	LKK 0, 3)	Reduced Ver			OA 150B)		sturbed or problematic.
	ledox (S5)		Piedmont Flo					
_	Matrix (S6)						A 149A, 153C, 153	D)
	rface (S7) (LRR P,							
Restrictive I	_ayer (if observed)	:						
Type:								_
Depth (inc	ches):		<u> </u>				Hydric Soil Prese	ent? Yes No
Remarks:								
Hydric s	oil indicator	F6 is me	et.					

Project/Site: Golden Triangle Phase 2	City/Co	ounty: Lowndes Cou	nty	Sampling Date: 2021-03-10
Applicant/Owner: TVA				Sampling Point: W-PEM-04
	Section			oumpling Forms
Landform (hillslope, terrace, etc.): Depression	Local			Slope (%): 2
	Lat: 33.384726			Slope (%)2 Datum: NAD 83
	Lat: _00.004720			
Soil Map Unit Name: Catalpa silty clay			NWI classificat	
Are climatic / hydrologic conditions on the site typical				
Are Vegetation, Soil, or Hydrology _	significantly disturb	ed? Are "Normal	Circumstances" pr	resent? YesNo
Are Vegetation, Soil, or Hydrology _	naturally problemate	tic? (If needed, e	explain any answers	s in Remarks.)
SUMMARY OF FINDINGS – Attach site	map showing sam	pling point location	ons, transects,	important features, etc.
Hydrophytic Vegetation Present? Yes	No	In the Commission Asses		
Hydric Soil Present? Yes	/ No	Is the Sampled Area within a Wetland?	Voc. V	No
	No	within a Wetland?	res	NO
Remarks:				
Sample plot in W-PEM-04. The	APT showed no	rmal conditions	s at the time	e leading up to and
during the field survey.				<b>5</b> 1
adming the held darvey.				
LIVERGLOCK				
HYDROLOGY				
Wetland Hydrology Indicators:	and all that analys		_	tors (minimum of two required)
Primary Indicators (minimum of one is required; ch			Surface Soil C	
1 <del>                                     </del>	Aquatic Fauna (B13) Marl Deposits (B15) <b>(LRR</b>	III)		etated Concave Surface (B8)
	Mari Deposits (B15) <b>(ERR</b> Hydrogen Sulfide Odor (C		Drainage Patt  Moss Trim Lir	
	Oxidized Rhizospheres al			Vater Table (C2)
	Presence of Reduced Iron		Crayfish Burro	
	Recent Iron Reduction in <sup>1</sup>		=,	sible on Aerial Imagery (C9)
	Thin Muck Surface (C7)	(	Geomorphic F	
	Other (Explain in Remarks	s)	Shallow Aquit	` '
Inundation Visible on Aerial Imagery (B7)		,	FAC-Neutral	` '
Water-Stained Leaves (B9)			_	oss (D8) (LRR T, U)
Field Observations:				
Surface Water Present? Yes No	Depth (inches): .5			
Water Table Present? Yes <u>✓</u> No	Depth (inches): 0			
	Depth (inches): 0	Wetland H	Hydrology Present	t? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitorin	ig well, aerial photos, prev	ious inspections), if ava	ilable:	
		, ,		
Remarks:				
Wetland hydrology indicators a	re met.			

<b>VEGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PEM-04
7 20 4 20 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r )  1	% Cover			Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)
2				Total Number of Dominant Species Across All Strata: 3 (B)
4				Percent of Dominant Species
6				111017110 032,171011, 011710.
7				Prevalence Index worksheet:
8				Total % Cover of:         Multiply by:           OBL species         45           x 1 = 45
	=			OBL species $\frac{45}{0}$ $x = \frac{45}{0}$ FACW species $\frac{6}{0}$ $x = \frac{45}{0}$
50% of total cover:	20% of	total cover	-	FAC species 30 x 2 = 90
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 0 x 4 = 0
1				UPL species 0 x 5 = 0
2.				Column Totals: 75 (A) 135 (B)
3				Prevalence Index = B/A = <u>1.8</u>
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				☑ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
	=			☐ Problematic Hydrophytic Vegetation¹ (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Xanthium strumarium	30		FAC	be present, unless disturbed or problematic.
2. Leersia oryzoides	20		OBL	Definitions of Four Vegetation Strata:
3. Carex lurida	15 10		OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Alternanthera philoxeroides		-	OBL	more in diameter at breast height (DBH), regardless of height.
5				
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8		•		Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				<b>Woody vine</b> – All woody vines greater than 3.28 ft in
11 12.		( <u> </u>		height.
12.	75% =	Total Co	vor	
50% of total cover: 38				
Woody Vine Stratum (Plot size: 30 ft r )	20 /0 01	total covel		
1				
2				
3				
4.				
5				Hydrophytic
			ver	Hydrophytic Vegetation
50% of total cover:				Present? Yes No
Remarks: (If observed, list morphological adaptations bel				1
		on wa	tor	
Dominance test is met. 25% coveraç	ie na ob	en wa	ıeı.	

SOIL Sampling Point: W-PEM-04

Depth	ription: (Describe Matrix	to the dep	th needed to docu Redo	ment the ox Feature			the absence	of indicators.)
(inches)	Color (moist)	%	Color (moist)	%	_Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
0 - 18	10YR 3/1	70	10YR 3/6	30	С	PL / M	Clay Loam	
-								
-								
1- 0.0						<del></del>	21 (1	Di B. Litt M.M.L.
			Reduced Matrix, M			rains.		PL=Pore Lining, M=Matrix.  for Problematic Hydric Soils <sup>3</sup> :
Histosol		ouble to un	Polyvalue B		•	LRR S. T. L	_	luck (A9) <b>(LRR O)</b>
_	oipedon (A2)		Thin Dark S					luck (A10) <b>(LRR S)</b>
Black Hi			Loamy Mucl	-		R O)		ed Vertic (F18) (outside MLRA 150A,B)
	n Sulfide (A4)		Loamy Gley		(F2)			ont Floodplain Soils (F19) (LRR P, S, T)
	l Layers (A5) Bodies (A6) <b>(LRR I</b>	P T II)	Depleted Ma  Redox Dark		-6)			lous Bright Loamy Soils (F20)
	cky Mineral (A7) <b>(L</b>		=		•		1 1 '	arent Material (TF2)
	esence (A8) (LRR		Redox Depr	essions (F	8)			hallow Dark Surface (TF12)
=	ck (A9) (LRR P, T)		☐ Marl (F10) <b>(</b> I	-			U Other (	Explain in Remarks)
	d Below Dark Surfa ark Surface (A12)	ce (A11)	Depleted Oc				T) <sup>3</sup> Indica	ators of hydrophytic vegetation and
I <b>=</b>	rairie Redox (A16) (	MLRA 150A	=		, ,	•		and hydrology must be present,
	lucky Mineral (S1)		Delta Ochric					ess disturbed or problematic.
_	Sleyed Matrix (S4)		Reduced Ve					
	edox (S5) Matrix (S6)		Piedmont Fl					153D)
	rface (S7) <b>(LRR P</b> ,	S. T. U)	<u> </u>	brigrit Lua	illy Solis	(F20) <b>(WILK</b>	A 149A, 153C,	1930)
	_ayer (if observed)							
Type:								
Depth (in	ches):		<u></u>				Hydric Soil	Present? Yes V No No
Remarks:								
Hydric s	oil indicator	s A4 an	d F6 are me	t.				

Project/Site: Golden Triangle Phase 2	ity/County: Lowndes County	Sampling Date: 2021-03-10
Applicant/Owner: TVA	State: Mississippi	Sampling Point: W-PEM-04_UPL
	ection, Township, Range: S34 T18N R16E	
	ocal relief (concave, convex, none): None	Slone (%)· 0
Subregion (LRR or MLRA): P 135A Lat: 33.38		Datum: NAD 83
Soil Map Unit Name: Catalpa silty clay	NWI classifica	
Are climatic / hydrologic conditions on the site typical for this time of ye		
Are Vegetation, Soil, or Hydrology significantly		
Are Vegetation, Soil, or Hydrology naturally pro	lematic? (If needed, explain any answe	rs in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing	sampling point locations, transects	, important features, etc.
Hydrophytic Vegetation Present? Yes No		
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area	
Wetland Hydrology Present? Yes No	within a Wetland? Yes	No
Remarks:		
Upland sample plot adjacent to W-PEM-04	The APT showed normal con	ditions at the time
	The 7th 1 showed hermal con	artionio at the time
leading up to and during the field survey.		
HYDROLOGY		
Wetland Hydrology Indicators:		ators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil	Cracks (B6)
Surface Water (A1) Aquatic Fauna (B1		getated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15)		
Saturation (A3)  Hydrogen Sulfide C	· · ·	` '
		Water Table (C2)
1 <del></del>	— ` <i>'</i>	isible on Aerial Imagery (C9)
Algal Mat or Crust (B4)  Thin Muck Surface		Position (D2)
Iron Deposits (B5) Other (Explain in R	· — ·	,
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral	Test (D5)
☐ Water-Stained Leaves (B9)	Sphagnum n	noss (D8) (LRR T, U)
Field Observations:		
Surface Water Present? Yes No Depth (inches		
Water Table Present? Yes No Depth (inches		
Saturation Present? Yes No Depth (inches (includes capillary fringe)	Wetland Hydrology Preser	nt? Yes No
Describe Recorded Data (stream gauge, monitoring well, aerial photo	, previous inspections), if available:	
Remarks:		
No indicators are met.		

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r )	% Cover			Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 3 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 33.3 (A/B)
6				
7				Prevalence Index worksheet:
8.				Total % Cover of: Multiply by:
		= Total Co	ver	OBL species 0 x 1 = 0
50% of total cover:				FACW species 40 x 2 = 80
45 ft	20 /0 01	total cove		FAC species 15 x 3 = 45
				FACU species 50 x 4 = 200
1				UPL species 30 x 5 = 150
2				Column Totals: 135 (A) 475 (B)
3				(5)
4				Prevalence Index = B/A = 3.52
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
		= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	r:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Cynodon dactylon	50	~	FACU	be present, unless disturbed or problematic.
2. Ranunculus abortivus	40	~	FACW	Definitions of Four Vegetation Strata:
3. Setaria faberi	30		UPL	
4. Schedonorus arundinaceus	15		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
				more in diameter at breast height (DBH), regardless of height.
	<del></del>			
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	135% :	= Total Co	ver	
50% of total cover: 68	20% of	total cover	r: 27	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3				
4				
5				Hydrophytic
		= Total Co		Vegetation Present?  Yes No
50% of total cover:		total cover	r:	100 <u> </u>
Remarks: (If observed, list morphological adaptations below	ow).			
No test is met.				
140 test is met.				

Sampling Point: W-PEM-04\_UPL

SOIL Sampling Point: W-PEM-04\_UPL

Profile Desc	ription: (Describe	to the depth	n needed to docun	nent the i	ndicator	or confirm	the absence of inc	dicators.)
Depth	Matrix			x Feature			_	_
(inches)	Color (moist)	<u> </u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0 - 20	10YR 3/2	95	10YR 3/6	5	С	М	Clay Loam	
	-				-			
-								
_	-							
·	-			· ——				
								•
	oncentration, D=Dep					ains.		Pore Lining, M=Matrix.
Hydric Soil I	Indicators: (Applic	able to all L						roblematic Hydric Soils³:
Histosol			Polyvalue Be				. –	
	pipedon (A2)		Thin Dark Su					(A10) (LRR S)
Black Hi			Loamy Mucky			R O)		ertic (F18) <b>(outside MLRA 150A,B)</b> oodplain Soils (F19) <b>(LRR P, S, T)</b>
_	n Sulfide (A4) I Layers (A5)		Loamy Gleye Depleted Mat		F2)			Bright Loamy Soils (F20)
	Bodies (A6) <b>(LRR F</b>	P. T. U)	Redox Dark		6)		(MLRA 15	
I = 1	cky Mineral (A7) (L		Depleted Dar	•	•			Material (TF2)
_	esence (A8) (LRR l		Redox Depre					w Dark Surface (TF12)
1 cm Mu	ck (A9) (LRR P, T)		Marl (F10) <b>(L</b>				U Other (Expla	ain in Remarks)
	Below Dark Surfac	e (A11)	Depleted Och					
=	ark Surface (A12)	MI DA 450A\	Iron-Mangan				•	of hydrophytic vegetation and
	airie Redox (A16) <b>(</b> lucky Mineral (S1) <b>(</b>		Umbric Surfa Delta Ochric			, 0)		hydrology must be present, isturbed or problematic.
	Bleyed Matrix (S4)	LKK 0, 3)	Reduced Ver			OA 150B)		sturbed or problematic.
	ledox (S5)		Piedmont Flo					
_	Matrix (S6)						A 149A, 153C, 153	D)
	rface (S7) (LRR P,							
Restrictive I	_ayer (if observed)	:						
Type:								_
Depth (inc	ches):		<u> </u>				Hydric Soil Prese	ent? Yes No
Remarks:								
Hydric s	oil indicator	F6 is me	et.					

Project/Site: Golden Triangle Phase 2	ity/County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-05
• • • • • • • • • • • • • • • • • • • •	ection, Township, Range: S34 T18N R16E
	ocal relief (concave, convex, none): Concave Slope (%): 2
Subregion (LRR or MLRA): P 135A Lat: 33.389	
Soil Map Unit Name: Catalpa silty clay	NWI classification: NA
Are climatic / hydrologic conditions on the site typical for this time of year	
	isturbed? Are "Normal Circumstances" present? YesNo
Are Vegetation, Soil, or Hydrology naturally prob	lematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing s	sampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area
Wetland Hydrology Present? Yes V No	within a Wetland? Yes No
Remarks:	
Sample plot in W-PFM-05. The APT showed	normal conditions at the time leading up to and
	normal conditions at the time leading up to and
during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	
High Water Table (A2)  High Water Table (A2)  Marl Deposits (B15)	
Saturation (A3) Hydrogen Sulfide Od	- ` <i>'</i>
	res along Living Roots (C3)
Sediment Deposits (B2)	
☐ Drift Deposits (B3) ☐ Recent Iron Reduction	
Algal Mat or Crust (B4)	
☐ Iron Deposits (B5) ☐ Other (Explain in Rer	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	☐ Sphagnum moss (D8) (LRR T, U)
Field Observations:  Surface Water Present?  Yes No Depth (inches):	1
Water Table Present? Yes V No Depth (inches):	
Saturation Present? Yes V No Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos	, previous inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	

<b>EGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PEM-05
Tree Stratum (Plot size: 30 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r )  1		Species?		Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)
2				Total Number of Dominant
3				Species Across All Strata: 3 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 30 x 1 = 30
		= Total Cov		FACW species 40 x 2 = 80
50% of total cover:	20% of	total cover	·	FAC species 30 x 3 = 90
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species $0   x4 = 0$
1				UPL species $0   x 5 = 0$
2				Column Totals: 100 (A) 200 (B)
3				
4				Prevalence Index = $B/A = 2.0$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7		-		2 - Dominance Test is >50%
8		- Total Car		3 - Prevalence Index is ≤3.0¹
50% of total cover:		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )	20% 01	lotal cover	•	
1 Andropogon glomeratus	40	~	FACW	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Ambrosia trifida	25		FAC	Definitions of Four Vegetation Strata:
3. Ludwigia alternifolia	20		OBL	Deminions of Four Vegetation Strata.
4. Carex lurida	5		OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
5. Packera glabella	5		OBL	more in diameter at breast height (DBH), regardless of height.
6. Rumex crispus	5		FAC	Carling/Charle Washington and Adding vines less
7				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.	<del>-</del>			
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10	<del>-</del>			
10 11				Woody vine – All woody vines greater than 3.28 ft in height.
12.				noight.
·-·		= Total Cov	ver	
50% of total cover: 50				
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4.				
5.				Hudranbutia
· ·		= Total Cov	ver	Hydrophytic Vegetation
50% of total cover:				Present? Yes No
Remarks: (If observed, list morphological adaptations bel				
	o			
Dominance test is met.				

SOIL Sampling Point: W-PEM-05

Profile Desc	cription: (Describe	to the depth	n needed to docur	nent the i	ndicator	or confir	m the absence of i	indicators.)
Depth	Matrix			x Feature		. 2	·	
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0 - 18	10YR 3/1	<u>70                                    </u>	10YR 3/6	30	<u>C</u>	M	Clay Loam	
				· ·			<u> </u>	
-								
_							·	
	-			· <del></del>	-			
							<del></del>	
							- <u> </u>	
						_		
<sup>1</sup> Type: C=C	oncentration, D=De	pletion, RM=F	Reduced Matrix, MS	S=Masked	Sand G	rains.	<sup>2</sup> Location: PL:	=Pore Lining, M=Matrix.
Hydric Soil	Indicators: (Appli	cable to all L	RRs, unless other	wise not	ed.)		Indicators for	Problematic Hydric Soils <sup>3</sup> :
Histosol	(A1)		Polyvalue Be	low Surfa	ce (S8) <b>(</b> I	LRR S, T,	U) 1 cm Muck	k (A9) <b>(LRR O)</b>
	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>
	stic (A3)		Loamy Muck	-		R O)		Vertic (F18) (outside MLRA 150A,B)
	en Sulfide (A4)		Loamy Gleye		F2)			Floodplain Soils (F19) (LRR P, S, T)
	d Layers (A5) Bodies (A6) <b>(LRR</b> l	D T II\	Depleted Ma Redox Dark		-6)		(MLRA 1	s Bright Loamy Soils (F20)
	ucky Mineral (A7) <b>(L</b>		Depleted Dai		,		,	nt Material (TF2)
_	resence (A8) (LRR		Redox Depre					low Dark Surface (TF12)
	ıck (A9) <b>(LRR P, T)</b>		Marl (F10) <b>(L</b>		-,		— 1	olain in Remarks)
Deplete	d Below Dark Surfa	ce (A11)	Depleted Ocl	nric (F11)	(MLRA 1	51)		
	ark Surface (A12)		Iron-Mangan					rs of hydrophytic vegetation and
_	rairie Redox (A16)		_					d hydrology must be present,
_	Mucky Mineral (S1) Bleyed Matrix (S4)	(LRR 0, S)	Delta Ochric					disturbed or problematic.
_	Redox (S5)		Reduced Ver					
	Matrix (S6)						<del>-137.)</del> RA 149A, 153С, 15	(3D)
_	rface (S7) (LRR P,	S, T, U)		9	,	/ (	, , , , ,	- ,
Restrictive	Layer (if observed	):						
Type:			<u></u>					
Depth (in	ches):						Hydric Soil Pre	esent? Yes <u>'</u> No
Remarks:								
Hydric s	soil indicator	F6 is me	et.					
		. • .•						

Project/Site: Golden Triangle Phase 2 Cit	y/County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-05_UPL
Investigator(s): J. Brown, O. Haney Section, Township, Range: S34 T18N R16E	
Landform (hillslope, terrace, etc.): Upland, Flat  Local relief (concave, convex, none): None  Slope (%): 0	
Subregion (LRR or MLRA): P 135A Lat: 33.3891	
Soil Map Unit Name: Catalpa silty clay NWI classification: NA	
Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)	
Are Vegetation, Soil, or Hydrology significantly disturbed?	
Are Vegetation, Soil, or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)	
SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.	
Hydrophytic Vegetation Present?  Hydric Soil Present?  Wetland Hydrology Present?  Remarks:	Is the Sampled Area within a Wetland?  Yes No
Upland sample plot adjacent to W-PEM-05. The APT showed normal conditions at the time	
leading up to and during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)  Surface Soil Cracks (B6)	
Surface Water (A1) Aquatic Fauna (B13) Sparsely Vegetated Concave Surface (B8)	
High Water Table (A2)  Marl Deposits (B15) (LRR U)  Drainage Patterns (B10)	
Saturation (A3) Hydrogen Sulfide Odor (C1) Moss Trim Lines (B16)	
☐ Water Marks (B1) ☐ Oxidized Rhizospheres along Living Roots (C3) ☐ Dry-Season Water Table (C2)	
Sediment Deposits (B2)  Presence of Reduced Iron (C4)  Crayfish Burrows (C8)	
☐ Drift Deposits (B3) ☐ Recent Iron Reduction in Tilled Soils (C6) ☐ Saturation Visible on Aerial Imagery (C9)	
Algal Mat or Crust (B4)  Thin Muck Surface (C7)  Geomorphic Position (D2)  Other (Crustein in Remarks)	
☐ Iron Deposits (B5) ☐ Other (Explain in Remarks) ☐ Shallow Aquitard (D3)	
☐ Inundation Visible on Aerial Imagery (B7) ☐ FAC-Neutral Test (D5) ☐ Sphagnum moss (D8) (LRR T, U)	
Surface Water Present? Yes No Depth (inches): _	
Water Table Present? Yes No _ ✓ Depth (inches): _	
Saturation Present? Yes No Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:	
Remarks:	
Wetland hydrology indicators are met.	

<b>VEGETATION</b>	(Four Strata)	) – Use	scientific	names of	of plants.

EGETATION (Four Stra	ta) – Use scientific na	mes of pl	ants.		Sampling Point: W-PEM-05_UPL
- 0	· · · · · · · · · · · · · · · · · · ·		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft			Species?		Number of Dominant Species
1					That Are OBL, FACW, or FAC: $\underline{1}$ (A)
2 3					Total Number of Dominant Species Across All Strata: 1 (B)
3					Species Across All Strata: 1 (B)
4 5					Percent of Dominant Species That Are ORL FACW or FAC: 100 (A/R)
6					That Are OBL, FACW, or FAC: 100 (A/B)
7					Prevalence Index worksheet:
8					Total % Cover of: Multiply by:
<u> </u>			= Total Cov	/er	OBL species $0   x 1 = 0$
	50% of total cover:				FACW species $60   x 2 = 120$
Sapling/Shrub Stratum (Plot size					FAC species $0 \times 3 = 0$
1					FACU species $0 \times 4 = 0$
2.					UPL species 10 x 5 = 50
3.					Column Totals: <u>70</u> (A) <u>170</u> (B)
4					Prevalence Index = B/A = 2.43
5					Hydrophytic Vegetation Indicators:
6					1 - Rapid Test for Hydrophytic Vegetation
7					2 - Dominance Test is >50%
8					☑ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
		=	= Total Cov	/er	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
	50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft	<u>r</u> )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Packera glabella		60		FACW	be present, unless disturbed or problematic.
2. Lamium amplexicaule		10		UPL	Definitions of Four Vegetation Strata:
3					Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4					more in diameter at breast height (DBH), regardless of height.
					neight.
6					Sapling/Shrub – Woody plants, excluding vines, less
7					than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8					Herb – All herbaceous (non-woody) plants, regardless
9					of size, and woody plants less than 3.28 ft tall.
					<b>Woody vine</b> – All woody vines greater than 3.28 ft in
11 12					height.
12.		70% =	Total Cov		
	50% of total cover: 35				
Woody Vine Stratum (Plot size		20 /0 01	total cover	· <u></u> -	
1					
2					
3					
4.					
5.					Hydrophytic
			= Total Cov	/er	Hydrophytic Vegetation
	50% of total cover:				Present? Yes No
Remarks: (If observed, list mo					<u> </u>
			oorn fi	ad The	are was open soil and dood sorn
•	ample plot was to	aken III	COIII II	:iu. I 116	ere was open soil and dead corn
stalks.					

SOIL Sampling Point: W-PEM-05\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	ndicator	or confirn	n the absence o	f indicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100		<u> </u>			Loam		
									_
-									
					-				
				· ——					
	-								
-									
¹Type: C=Co	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> l ocation: P	PL=Pore Lining, M=Matr	rix
	Indicators: (Appli					anio.		or Problematic Hydric	
Histosol			Polyvalue Be			RR S. T. I		ick (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					ick (A10) (LRR S)	
Black Hi			Loamy Muck					d Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye					nt Floodplain Soils (F19	
Stratified	l Layers (A5)		Depleted Ma	trix (F3)			Anomalo	ous Bright Loamy Soils	(F20)
	Bodies (A6) (LRR I		Redox Dark	Surface (F	<del>-</del> 6)		,	A 153B)	
_	cky Mineral (A7) (L		Depleted Dai					ent Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			allow Dark Surface (TF	12)
	ck (A9) (LRR P, T)		Marl (F10) <b>(L</b>		/MI DA 4	E4)	U Other (E	xplain in Remarks)	
	d Below Dark Surfa ark Surface (A12)	ce (ATT)	Depleted Ocl				T) <sup>3</sup> Indicat	tors of hydrophytic vege	atation and
	rairie Redox (A16) (	MLRA 150A)						nd hydrology must be p	
	lucky Mineral (S1)		Delta Ochric			, •,		s disturbed or problema	
	leyed Matrix (S4)	(,,	Reduced Ver			0A, 150B)			
	edox (S5)		Piedmont Flo						
Stripped	Matrix (S6)		Anomalous E	Bright Loar	my Soils (	F20) <b>(MLF</b>	RA 149A, 153C, 1	153D)	
	rface (S7) (LRR P,								
Restrictive I	_ayer (if observed	):							
Type:			<u>—</u>						
Depth (inc	ches):						Hydric Soil P	resent? Yes	No
Remarks:									
No indic	ators are m	et.							
Tto maio		<b>.</b>							

Project/Site: Golden Triangle Phase 2 City/	/County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-06
	tion, Township, Range: S33 T18N R16E
	al relief (concave, convex, none): Concave Slope (%): 3
	02 Long: -88.624276 Datum: NAD 83
Soil Map Unit Name: Okolona silty clay, 0 to 1 percent slopes	NWI classification: NA
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distu	
Are Vegetation, Soil, or Hydrology naturally problem	natic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing sa	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	
Hydric Soil Present? Yes V No	Is the Sampled Area within a Wetland? Yes V No
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	
Sample plot in W-PEM-06. The APT showed n	ormal conditions at the time leading up to and
during the field survey.	
during the held survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  High Water Table (A2)  Marl Deposits (B15) (LF	
Trydrogen Suinde Odor	· · · · · · · · · · · · · · · · · · ·
	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced In	
Drift Deposits (B3)  Recent Iron Reduction i	
Algal Mat or Crust (B4)  Thin Muck Surface (C7)  Other (Evelsis in Rema	,
☐ Iron Deposits (B5) ☐ Other (Explain in Rema	
Inundation Visible on Aerial Imagery (B7)  Water-Stained Leaves (B9)	FAC-Neutral Test (D5)
Field Observations:	☐ Sphagnum moss (D8) (LRR T, U)
Surface Water Present? Yes No Depth (inches): 2	
Water Table Present? Yes   ✓ No Depth (inches): 0	
Saturation Present?  Yes  No Depth (inches): 0	Wetland Hydrology Present? Yes No
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	revious inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	
Wetland hydrology indicators are met.	

- 01 ( D) ( ) 20 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1.	% Cover			Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 80 x 1 = 80
		= Total Cov		FACW species 0 x 2 = 0
50% of total cover:	20% of	total cover	·	FAC species 5 x 3 = 15
Sapling/Shrub Stratum (Plot size: 30 ft r )				FACU species 0 x 4 = 0
1				UPL species 0 x 5 = 0
2				Column Totals: 85 (A) 95 (B)
3				
4				Prevalence Index = B/A = 1.1
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8		- Total Co		3 - Prevalence Index is ≤3.0¹
50% of total cover:		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 30 ft r )	20% 01	total cover	•	
1 Carex Iurida	40	~	OBL	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Leersia oryzoides	25		OBL	Definitions of Four Vegetation Strata:
3. Alternanthera philoxeroides	15		OBL	Deminions of Four Vegetation Strata.
Xanthium strumarium	5		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
··				more in diameter at breast height (DBH), regardless of height.
5 6.				Carling (Observed) - We asked a large a supply of the province of the supply of the su
6 7				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10.				
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12.				noight.
		= Total Cov	ver	
50% of total cover: 43				
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4				
5.				Hydrophytic
	=	= Total Cov	ver	Vegetation
50% of total cover:	20% of	total cover	:	Present? Yes No
Remarks: (If observed, list morphological adaptations be	elow).			
	,			
Rapid test is met.				

SOIL Sampling Point: W-PEM-06

Profile Desc	ription: (Describe	to the dept	n needed to docum	nent the	indicator	or confir	m the absence of i	indicators.)
Depth	Matrix			x Feature		. 2	·	
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type'	Loc 2	<u>Texture</u>	Remarks
0 - 20	10YR 3/1	80	10YR 3/6	20	С	М	Clay Loam	
-								
	-						·	
·							·	
							<u> </u>	
				. <u></u>				
-								
¹Type: C=C	ncentration D=De	nletion PM=	Reduced Matrix, MS	S=Macked	d Sand G	raine	<sup>2</sup> Location: PL	=Pore Lining, M=Matrix.
			RRs, unless other			airis.		Problematic Hydric Soils <sup>3</sup> :
Histosol			Polyvalue Be			RRST		k (A9) <b>(LRR O)</b>
_	pipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>
Black Hi			Loamy Mucky					Vertic (F18) (outside MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye			. 0,		Floodplain Soils (F19) (LRR P, S, T)
	Layers (A5)		Depleted Mat		/			is Bright Loamy Soils (F20)
	Bodies (A6) (LRR I	P, T, U)	Redox Dark		<del>-</del> 6)		(MLRA	
	cky Mineral (A7) (L		Depleted Dar					nt Material (TF2)
_	esence (A8) (LRR		Redox Depre					low Dark Surface (TF12)
1 cm Mu	ck (A9) (LRR P, T)		Marl (F10) (L	RR U)			Other (Exp	plain in Remarks)
Depleted	d Below Dark Surfa	ce (A11)	Depleted Och	nric (F11)	(MLRA 1	51)		
	ark Surface (A12)		Iron-Mangan	ese Mass	es (F12)	(LRR O, P		rs of hydrophytic vegetation and
	rairie Redox (A16) (		_					d hydrology must be present,
	lucky Mineral (S1)	(LRR O, S)	Delta Ochric					disturbed or problematic.
	Sleyed Matrix (S4)		Reduced Ver					
	edox (S5)		Piedmont Flo					·
	Matrix (S6)	O T II)	Anomalous B	right Loa	my Soils	(F20) <b>(ML</b>	RA 149A, 153C, 15	33D)
	rface (S7) (LRR P,							
	_ayer (if observed)	):						
Type:			<u></u>					
	ches):						Hydric Soil Pre	esent? Yes V No No
Remarks:								
Hydric s	oil indicator	s A4 and	d F6 are met	•				

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-06_UPL
''	tion, Township, Range: S33 T18N R16E
Landform (hillslope, terrace, etc.): Upland, Flat Loca	
Subregion (LRR or MLRA): P 135A Lat: 33.38736	
Soil Map Unit Name: Okolona silty clay, 0 to 1 percent slopes	
	NWI classification: NA
Are climatic / hydrologic conditions on the site typical for this time of year?	.4
Are Vegetation, Soil, or Hydrology significantly distu	ırbed? Are "Normal Circumstances" present? YesNo
Are Vegetation, Soil, or Hydrology naturally problem	natic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing said	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	
Hydric Soil Present? Yes No	Is the Sampled Area within a Wetland? Yes No✓
Hydric Soil Present?  Wetland Hydrology Present?  Yes No V	within a Wetland? Yes No
Remarks:	
Upland sample plot adjacent to W-PEM-06. The	ne APT showed normal conditions at the time
leading up to and during the field survey.	
leading up to and during the held survey.	
HADBOLOGA	
HYDROLOGY	Connectors (minimum of two required)
Wetland Hydrology Indicators:  Primary Indicators (minimum of one is required; check all that apply)	Secondary Indicators (minimum of two required)
Surface Water (A1)  Aquatic Fauna (B13)	Surface Soil Cracks (B6)
High Water Table (A2)  High Water Table (A2)  Advance Faulta (B15)  Marl Deposits (B15) (LF	Sparsely Vegetated Concave Surface (B8)  Drainage Patterns (B10)
Saturation (A3)  Hydrogen Sulfide Odor	
	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced Ir	
Drift Deposits (B3)	n Tilled Soils (C6) Saturation Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	
☐ Iron Deposits (B5) ☐ Other (Explain in Remains)	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	☐ Sphagnum moss (D8) (LRR T, U)
Field Observations:  Surface Water Present?  Yes No Depth (inches):	
Water Table Present?  Yes No _ Depth (inches):	
Saturation Present? Yes No Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	evious inspections), if available:
Remarks:	
No indicators are met.	

20.4		Dominant		Dominance Test worksheet:	
Tree Stratum (Plot size: 30 ft r )		Species?		Number of Dominant Species	
1				That Are OBL, FACW, or FAC: 0 (A)	
2				Total Number of Dominant	
3				Species Across All Strata: 2 (B)	
4				Percent of Dominant Species	
5				That Are OBL, FACW, or FAC: 0 (A/E	3)
6				Prevalence Index worksheet:	
7					
8				Total % Cover of: Multiply by:	
	=	= Total Co	ver	OBL species $0 \times 1 = 0$	
50% of total cover:	20% of	total cover	·:	FACW species $\frac{0}{20}$ x 2 = $\frac{0}{60}$	
Sapling/Shrub Stratum (Plot size: 30 ft r )				FAC species $20$ $\times 3 = 60$	
1				FACU species 50 x 4 = 200	
2.				UPL species 40 x 5 = 200	
3.				Column Totals: <u>110</u> (A) <u>460</u> (B	)
4.				Prevalence Index = B/A = 4.2	
5				Hydrophytic Vegetation Indicators:	
6				1 - Rapid Test for Hydrophytic Vegetation	
7				2 - Dominance Test is >50%	
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>	
		= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)	
50% of total cover:	20% of	total cover	:		
Herb Stratum (Plot size: 30 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must	
1. Cynodon dactylon	50		FACU	be present, unless disturbed or problematic.	
2. Setaria faberi	40		UPL	Definitions of Four Vegetation Strata:	
3. Schedonorus arundinaceus	20		FAC	The Meady plants avaluding vines 2 in (7.0 am)	
4.				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) of more in diameter at breast height (DBH), regardless of	
5.				height.	-
6.				Sapling/Shrub – Woody plants, excluding vines, less	
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.	
8.					
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardles of size, and woody plants less than 3.28 ft tall.	3
10.					
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in	
12.				height.	
12	110% =	- Total Co			
50% of total cover: 55					
	20% 01	total cover			
Woody Vine Stratum (Plot size: 30 ft r )					
1					
2					
3					
4					
5				Hydrophytic	
		= Total Co		Vegetation Present? Yes No	
50% of total cover:		total cover	:	100 100	
Remarks: (If observed, list morphological adaptations below	ow).				
No test is met.					

Sampling Point: W-PEM-06\_UPL

SOIL Sampling Point: W-PEM-06\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Cd	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> Location: PL:	=Pore Lining, M=Matr	riy
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RRSTI		k (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-		-,		Floodplain Soils (F19	
	Layers (A5)		Depleted Ma		,			s Bright Loamy Soils	
Organic	Bodies (A6) (LRR I	P, T, U)	Redox Dark	Surface (F	<del>-</del> 6)		(MLRA 1	153B)	
5 cm Mu	cky Mineral (A7) (L	.RR P, T, U)	Depleted Dar	rk Surface	e (F7)		Red Paren	nt Material (TF2)	
	esence (A8) (LRR		Redox Depre	essions (F	8)		<u></u> Very Shall	ow Dark Surface (TF	12)
	ck (A9) (LRR P, T)						U Other (Exp	olain in Remarks)	
	Below Dark Surfa	ce (A11)	Depleted Ocl				3		
	ark Surface (A12)		Iron-Mangan					rs of hydrophytic vege	
	rairie Redox (A16) (					, U)		d hydrology must be p	
	lucky Mineral (S1)	(LKK 0, 5)	Delta Ochric			OA 150D)		disturbed or problema	atic.
	edox (S5)		Reduced Ver						
	Matrix (S6)						+9A) RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,	S. T. U)	Anomalous L	night Loai	illy Solis (	1 20) (IVILIV	A 149A, 133C, 13	30)	
	_ayer (if observed								
Type:		•							
, , <u> </u>	ches):						Hydric Soil Pre	esent? Yes	No 🗸
Remarks:							Tryuno com r re		
No indic	ators are m	et.							

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-07
	tion, Township, Range: S33 T18N R16E
Landform (hillslope, terrace, etc.): Fringe Local	ol rollof (concavo, convoy, nono): None Slone (%): 0
Subregion (LRR or MLRA): P 135A Lat: 33.38418	
Soil Map Unit Name: Brooksville silty clay, 1 to 3 percent slopes	
	NWI classification: NA
Are climatic / hydrologic conditions on the site typical for this time of year?	· · · · · · · · · · · · · · · · · · ·
Are Vegetation, Soil, or Hydrology significantly distu	ırbed? Are "Normal Circumstances" present? YesNo
Are Vegetation, Soil, or Hydrology naturally problem	natic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing sa	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	In the Complet Area
Hydric Soil Present? Yes V No	Is the Sampled Area within a Wetland? Yes  No
Wetland Hydrology Present? Yes No	within a wettand? Tes No No
Remarks:	
Sample plot in W-PEM-07. The APT showed n	ormal conditions at the time leading up to and
during the field survey.	5 1
daring the note out vey.	
HADBOLOCA	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)  Surface Water (A1)  Aquatic Fauna (B13)	Surface Soil Cracks (B6)
Surface Water (A1)  High Water Table (A2)  Aquatic Fauna (B13)  Marl Deposits (B15) (LF	Sparsely Vegetated Concave Surface (B8)  Drainage Patterns (B10)
Saturation (A3)  Hydrogen Sulfide Odor	
	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced In	
Drift Deposits (B3)  Recent Iron Reduction i	
Algal Mat or Crust (B4) Thin Muck Surface (C7)	_
Iron Deposits (B5) Other (Explain in Rema	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes No Depth (inches): 2_	
Water Table Present? Yes _ V No _ Depth (inches): 0	
Saturation Present? Yes V No Depth (inches): 0	Wetland Hydrology Present? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	evious inspections), if available:
	, ,
Remarks:	
Wetland hydrology indicators are met.	
,	

Tree Stratum (Plot size: 30 ft r)  1 2 3 4 5 6 7.	<u>% Cover</u>		Status	Dominance Test worksheet:  Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)
1				
3				
4				Total Number of Dominant
5 6				Species Across All Strata: 3 (B)
6				Percent of Dominant Species
6 7				That Are OBL, FACW, or FAC: 100 (A/B)
7				Prevalence Index worksheet:
				Total % Cover of: Multiply by:
8				OBL species 42 x 1 = 42
		= Total Cov		FACW species $0$ $x = 0$
50% of total cover:	20% of	total cover	·	FAC species 30 x 3 = 90
Sapling/Shrub Stratum (Plot size: 15 ft r )  1 Salix nigra	5	~	OBL	FACU species $0   x4 = 0$
** <del></del>				UPL species 0 x 5 = 0
2				Column Totals: 72 (A) 132 (B)
3				
4				Prevalence Index = B/A = 1.83
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7		-		2 - Dominance Test is >50%
8		Total Cov		3 - Prevalence Index is ≤3.0¹
50% of total cover: 3				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )	20 /6 01	total cover	•	1
1 Packera glabella	35	~	OBL	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Cyperus esculentus	30		FAC	Definitions of Four Vegetation Strata:
3 Salix nigra			OBL	Definitions of Four Vegetation Strata.
4				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
5				more in diameter at breast height (DBH), regardless of height.
^				Continue (Charaka Manada and and and and and and and and an
6 7				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10.				
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12.				inoight.
	070/	= Total Cov	er	
50% of total cover: 34				
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4.				
5.				Hydrophytic
		= Total Cov	er	Vegetation
50% of total cover:	20% of	total cover	<u> </u>	Present? Yes No
Remarks: (If observed, list morphological adaptations b	elow).			
Dominance test is met.	,			
Dominance test is met.				

SOIL Sampling Point: W-PEM-07

		needed to document the indicator or	confirm the absence	e of indicators.)
Depth (inches)	Matrix Color (moist) %	Redox Features  Color (moist) % Type <sup>1</sup> L	oc <sup>2</sup> Texture	Remarks
-				
				·
				·
-				
_				
1Type: C=C	Properties D-Depleties PM-	Reduced Matrix, MS=Masked Sand Grain	21 coation	: PL=Pore Lining, M=Matrix.
	·	RRs, unless otherwise noted.)		s for Problematic Hydric Soils <sup>3</sup> :
Histosol	, , ,	Polyvalue Below Surface (S8) (LRF		Muck (A9) <b>(LRR O)</b>
l <del>''</del>	pipedon (A2)	Thin Dark Surface (S9) (LRR S, T,		Muck (A10) (LRR S)
Black Hi		Loamy Mucky Mineral (F1) (LRR O		ced Vertic (F18) (outside MLRA 150A,B)
I <b>=</b>	n Sulfide (A4)	Loamy Gleyed Matrix (F2)		nont Floodplain Soils (F19) (LRR P, S, T)
Stratified	Layers (A5)	Depleted Matrix (F3)	L Anom	alous Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark Surface (F6)		.RA 153B)
	cky Mineral (A7) (LRR P, T, U)	Depleted Dark Surface (F7)		Parent Material (TF2)
	esence (A8) (LRR U)	Redox Depressions (F8)		Shallow Dark Surface (TF12)
	ck (A9) <b>(LRR P, T)</b>	Mari (F10) (LRR U)		(Explain in Remarks)
	Below Dark Surface (A11)	Depleted Ochric (F11) (MLRA 151)		cators of hydrophytic vegetation and
	ark Surface (A12) rairie Redox (A16) <b>(MLRA 150A</b> )	Iron-Manganese Masses (F12) (LR Umbric Surface (F13) (LRR P, T, U		etland hydrology must be present,
	lucky Mineral (S1) (LRR O, S)	Delta Ochric (F17) (MLRA 151)		less disturbed or problematic.
_	eleyed Matrix (S4)	Reduced Vertic (F18) (MLRA 150A		icos dicialista di prosioniano.
_	edox (S5)	Piedmont Floodplain Soils (F19) (N		
	Matrix (S6)	Anomalous Bright Loamy Soils (F2)		C, 153D)
Dark Su	face (S7) (LRR P, S, T, U)			
Restrictive I	_ayer (if observed):			
Type:		<u> </u>		,
Depth (inc	ches):	<u></u>	Hydric Soi	I Present? Yes No
Remarks:				
Sample	plot was inundated a	and no soil sample was ta	ken. Indicator	A4 is met.
	,	, , , , , , , , , , , , , , , , , , ,		

Project/Site: Golden Triangle Phase 2 City	//County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-07_UPL
••	ction, Township, Range: S33 T18N R16E
	ral relief (concave, convex, none): None Slope (%): 0
Subregion (LRR or MLRA): P 135A Lat: 33.3957	
Soil Map Unit Name: Brooksville silty clay, 1 to 3 percent slopes	
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly dist	
Are Vegetation, Soil, or Hydrology naturally proble	matic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing sa	ampling point locations, transects, important features, etc.
Hydrophytia Vagetation Propent?	
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	
Unland sample plot adjacent to W-PFM-07 T	he APT showed normal conditions at the time
1	The Air I showed normal conditions at the time
leading up to and during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15) (L	
Saturation (A3) Hydrogen Sulfide Odor	
	s along Living Roots (C3) Dry-Season Water Table (C2)
☐ Sediment Deposits (B2) ☐ Presence of Reduced ☐ Drift Deposits (B3) ☐ Recent Iron Reduction	
Algal Mat or Crust (B4)  Thin Muck Surface (C7)	
Iron Deposits (B5)  Other (Explain in Remains)	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes No Depth (inches):	
Water Table Present? Yes No Depth (inches):	
Saturation Present? Yes No Depth (inches):	Wetland Hydrology Present? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos, p	revious inspections), if available:
Remarks:	
No indicators are met.	

VEGETATION (Four Strata) - Use scientific names of plants. Sampling Point: W-PEM-07\_UPL Absolute Dominant Indicator **Dominance Test worksheet:** Tree Stratum (Plot size: 30 ft r % Cover Species? Status **Number of Dominant Species** That Are OBL, FACW, or FAC: **Total Number of Dominant** 1 \_\_\_\_ (B) Species Across All Strata: Percent of Dominant Species That Are OBL, FACW, or FAC: 0 (A/B) 6. \_\_\_\_\_\_ Prevalence Index worksheet: Total % Cover of: Multiply by: 0 x 1 = 0 OBL species = Total Cover FACW species 0 x 2 = 050% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_ 0 <u>x 3 = 0</u> FAC species Sapling/Shrub Stratum (Plot size: 15 ft r ) FACU species 0 x4 = 010 x = 50**UPL** species Column Totals: 10 \_\_\_\_ (A) <u>50</u> (B) Prevalence Index = B/A = 5.0**Hydrophytic Vegetation Indicators:** 1 - Rapid Test for Hydrophytic Vegetation 2 - Dominance Test is >50% 3 - Prevalence Index is ≤3.0<sup>1</sup> Problematic Hydrophytic Vegetation<sup>1</sup> (Explain) 50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_ Herb Stratum (Plot size: 5 ft r ) <sup>1</sup>Indicators of hydric soil and wetland hydrology must 1 Lamium amplexicaule be present, unless disturbed or problematic. 2. **Definitions of Four Vegetation Strata:** Tree - Woody plants, excluding vines, 3 in. (7.6 cm) or 4. \_\_\_\_\_\_ \_\_\_ \_\_\_\_ \_\_\_\_\_\_\_ more in diameter at breast height (DBH), regardless of height. Sapling/Shrub - Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall. **Herb** – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall. Woody vine - All woody vines greater than 3.28 ft in height. 10% = Total Cover 50% of total cover: 5 20% of total cover: 2 Woody Vine Stratum (Plot size: 30 ft r Hydrophytic \_\_\_\_\_ = Total Cover Vegetation Yes No Present? 50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_ Remarks: (If observed, list morphological adaptations below). No test is met. Sample plot was taken in corn field. There was open soil and dead corn stalks.

SOIL Sampling Point: W-PEM-07\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	ndicator	or confirn	n the absence o	f indicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100		<u> </u>			Loam		
									_
-									
					-				
				· ——					
	-								
-									
¹Type: C=Co	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> l ocation: P	PL=Pore Lining, M=Matr	rix
	Indicators: (Appli					anio.		or Problematic Hydric	
Histosol			Polyvalue Be			RR S. T. I		ick (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					ick (A10) (LRR S)	
Black Hi			Loamy Muck					d Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye					nt Floodplain Soils (F19	
Stratified	l Layers (A5)		Depleted Ma	trix (F3)			Anomalo	ous Bright Loamy Soils	(F20)
	Bodies (A6) (LRR I		Redox Dark	Surface (F	<del>-</del> 6)		,	A 153B)	
_	cky Mineral (A7) (L		Depleted Dai					ent Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			allow Dark Surface (TF	12)
	ck (A9) (LRR P, T)		Marl (F10) <b>(L</b>		/MI DA 4	E4)	U Other (E	xplain in Remarks)	
	d Below Dark Surfa ark Surface (A12)	ce (ATT)	Depleted Ocl				T) <sup>3</sup> Indicat	tors of hydrophytic vege	atation and
	rairie Redox (A16) (	MLRA 150A)						nd hydrology must be p	
	lucky Mineral (S1)		Delta Ochric			, •,		s disturbed or problema	
	leyed Matrix (S4)	(,,	Reduced Ver			0A, 150B)			
	edox (S5)		Piedmont Flo						
Stripped	Matrix (S6)		Anomalous E	Bright Loar	my Soils (	F20) <b>(MLF</b>	RA 149A, 153C, 1	153D)	
	rface (S7) (LRR P,								
Restrictive I	_ayer (if observed	):							
Type:			<u>—</u>						
Depth (inc	ches):						Hydric Soil P	resent? Yes	No
Remarks:									
No indic	ators are m	et.							
Tto maio		<b>.</b>							

Project/Site: Golden Triangle Phase 2	City/County: Lowi	ndes County	Sampling Date: 2021-03-11
Applicant/Owner: TVA	only/county:		Sampling Point: W-PEM-08
	Section, Township,		Camping Cont.
Landform (hillslope, terrace, etc.): Depression	Local relief (concav	o convoy nono). Concave	Slone (%): 5
	<u> </u>	· -	Slope (%) Datum: NAD 83
	Lat: Oc.oboood		
Soil Map Unit Name: Catalpa silty clay		NWI classifica	
Are climatic / hydrologic conditions on the site typical for the			
Are Vegetation, Soil, or Hydrology	significantly disturbed?	Are "Normal Circumstances" p	resent? YesNo
Are Vegetation, Soil, or Hydrology	naturally problematic? (	If needed, explain any answer	rs in Remarks.)
SUMMARY OF FINDINGS – Attach site map	showing sampling poin	nt locations, transects	, important features, etc.
Hydrophytic Vegetation Present? Yes I	No la the Com	alad Awaa	
Hydric Soil Present? Yes	No Is the Samp		No
Wetland Hydrology Present? Yes I		etiality res	NO
Remarks:	<u> </u>		
Sample plot in W-PEM-08. The APT	showed normal co	nditions at the time	e leading up to and
during the field survey.			<b>5</b> 1
during the held survey.			
HADBOLOCA			
HYDROLOGY		Casandan Indiaa	to no (noinino uno of true no suine d)
Wetland Hydrology Indicators:  Primary Indicators (minimum of one is required; check all	I that apply)		tors (minimum of two required)
	c Fauna (B13)	Surface Soil (	getated Concave Surface (B8)
	eposits (B15) <b>(LRR U)</b>	Drainage Pat	
	gen Sulfide Odor (C1)	Moss Trim Li	
	ed Rhizospheres along Living R		Water Table (C2)
	nce of Reduced Iron (C4)	Crayfish Burr	
☑ Drift Deposits (B3) ☐ Recent	t Iron Reduction in Tilled Soils (	C6) Saturation Vis	sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	luck Surface (C7)	✓ Geomorphic I	Position (D2)
☐ Iron Deposits (B5) ☐ Other (	(Explain in Remarks)	Shallow Aqui	` '
Inundation Visible on Aerial Imagery (B7)		FAC-Neutral	` '
☐ Water-Stained Leaves (B9)	Ţ	<u></u> Sphagnum m	noss (D8) <b>(LRR T, U)</b>
Field Observations:			
Surface Water Present? Yes No De			
Water Table Present? Yes No De		Wedler dilbedeels on Bosses	10 V V N-
Saturation Present? Yes Ves No De (includes capillary fringe)	eptn (inches): U	Wetland Hydrology Presen	t? Yes No
Describe Recorded Data (stream gauge, monitoring well,	aerial photos, previous inspect	ions), if available:	
Remarks:			
Wetland hydrology indicators are m	ıet		
Wettaria flyarology maleators are in			

<b>'EGETATION (Four Strata)</b> – Use scientific na	mes of pl	ants.		Sampling Point: W-PEM-08
Total Objections (Dietoine 30 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1.		Species?		Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8.				OBL species <u>5</u> x 1 = <u>5</u>
50% of total cover:		= Total Cov		FACW species <u>50</u> x 2 = <u>100</u>
Sapling/Shrub Stratum (Plot size: 15 ft r )	20 /6 01	total cover		FAC species <u>45</u> x 3 = <u>135</u>
1				FACU species <u>5</u> x 4 = <u>20</u>
2.				UPL species <u>0</u> x 5 = <u>0</u>
3.				Column Totals: 105 (A) 260 (B)
4				Prevalence Index = B/A = 2.48
5.				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				☑ 3 - Prevalence Index is ≤3.0¹
	=	= Total Cov	/er	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Andropogon glomeratus	50		FACW	be present, unless disturbed or problematic.
2. Ambrosia trifida	40		FAC	Definitions of Four Vegetation Strata:
3. Plantago virginica	5 5		FACU	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Typha latifolia 5. Xanthium strumarium	5		OBL	more in diameter at breast height (DBH), regardless of height.
			FAC	
6				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				
8 9.	·			<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10.				Woody vine All woody vines greater than 3.28 ft in
11.				<b>Woody vine</b> – All woody vines greater than 3.28 ft in height.
12				
	105% =	= Total Cov	/er	
50% of total cover: 53	20% of	total cover	21	
Woody Vine Stratum (Plot size: 30 ft r )  1.				
2.				
3.				
4				
5.				Hydrophytic
		= Total Cov	/er	Vegetation
50% of total cover:	20% of	total cover	:	Present? Yes No
Remarks: (If observed, list morphological adaptations below	ow).			
Dominance test is met.				

SOIL Sampling Point: W-PEM-08

Profile Des	cription: (Describe to the depti	n needed to docu	ment the	indicator of	or confirm	the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	%	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
-							
			-				
l — —							_
			_				
-							
			-				
l ———							
			_				
_							
17		Darley and Matrice M	0 Ml			21 1:	David Units of M. Madata
	oncentration, D=Depletion, RM=I Indicators: (Applicable to all L				ains.		Pore Lining, M=Matrix.  Problematic Hydric Soils <sup>3</sup> :
l <u> </u>							•
Histosol	'	Polyvalue Be				. —	
I <b>=</b>	pipedon (A2)	Thin Dark Su					(A10) <b>(LRR S)</b>
_	istic (A3)	Loamy Muck	-		O)		ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)	Loamy Gleye		(F2)			oodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	Depleted Ma					Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark	,	,		(MLRA 15	•
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da		. ,			Material (TF2)
	resence (A8) (LRR U)	Redox Depre		<del>-</del> 8)			w Dark Surface (TF12)
_	uck (A9) <b>(LRR P, T)</b>	Marl (F10) (L				U Other (Expla	ain in Remarks)
	d Below Dark Surface (A11)	Depleted Oc					
	ark Surface (A12)	Iron-Mangan				•	of hydrophytic vegetation and
_	rairie Redox (A16) (MLRA 150A)	_			, U)		nydrology must be present,
	Mucky Mineral (S1) (LRR O, S)	Delta Ochric					sturbed or problematic.
_	Gleyed Matrix (S4)	Reduced Ve					
	Redox (S5)	Piedmont Flo					
	d Matrix (S6)	Anomalous I	Bright Loa	amy Soils (F	-20) <b>(MLR</b>	A 149A, 153C, 153I	D)
	ırface (S7) <b>(LRR P, S, T, U)</b>						
Restrictive	Layer (if observed):						
Type:							
Depth (in	ches):	<u></u>				Hydric Soil Pres	ent? Yes <u>'</u> No
Remarks:							
Coil con	anla waa nat takan d	ua ta inund	ation	Lludria	مماا مم	soumed due	to provolopoo of
Son San	nple was not taken d	ue to muna	ation.	пушіс	Son as	sumed due	to prevalence of
wetland	l hydrology and hydr	ophytic vec	etatio	n.			
	, , .	-	,				

Project/Site: Golden Triangle Phase 2	City/C	ounty: Lowndes County	Sampling Date: 2021-03-11
Applicant/Owner: TVA	0.9,0		ppi Sampling Point: W-PEM-08_2
Investigator(s): J. Brown, O. Haney	Section	n, Township, Range: S34 T18N R16	
Landform (hillslope, terrace, etc.): Depress	sion Local	relief (concave, convex, none): Conc	
	Lat: 33.383191	Long: -88.614966	
Soil Map Unit Name: Catalpa silty clay		NWI class	
Are climatic / hydrologic conditions on the si			
Are Vegetation, Soil, or Hyd	rology significantly disturb	ped? Are "Normal Circumstance	s" present? YesNo
Are Vegetation, Soil, or Hyd	rology naturally problema	tic? (If needed, explain any an	swers in Remarks.)
SUMMARY OF FINDINGS - Attac	ch site map showing sam	pling point locations, transe	cts, important features, etc.
Hydrophytic Vegetation Present?	Yes No	In the Commission Area	
Hydric Soil Present?	Yes No	Is the Sampled Area within a Wetland? Yes	✓ No
	Yes No	within a wetland?	NO
Remarks:			
Sample plot in W-PEM-08.	. The APT showed no	rmal conditions at the ti	me leading up to and
during the field survey.			•
admig the held editely.			
HYDROLOGY			
Wetland Hydrology Indicators:		Secondary In	dicators (minimum of two required)
Primary Indicators (minimum of one is requ	uired; check all that apply)		Soil Cracks (B6)
Surface Water (A1)	Aquatic Fauna (B13)		Vegetated Concave Surface (B8)
High Water Table (A2)	Marl Deposits (B15) (LRR		Patterns (B10)
Saturation (A3)	Hydrogen Sulfide Odor (C		m Lines (B16)
Water Marks (B1)	Oxidized Rhizospheres al	ong Living Roots (C3) 🔲 Dry-Seas	on Water Table (C2)
Sediment Deposits (B2)	Presence of Reduced Iron	n (C4)	Burrows (C8)
Drift Deposits (B3)	Recent Iron Reduction in		n Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	Thin Muck Surface (C7)	<del></del> ·	hic Position (D2)
☐ Iron Deposits (B5)	Other (Explain in Remark	· =	Aquitard (D3)
Inundation Visible on Aerial Imagery ( Water-Stained Leaves (B9)	Б7)	<b>—</b>	tral Test (D5) m moss (D8) <b>(LRR T, U)</b>
Field Observations:		Эрпауни	(LKK 1, 0)
	No Depth (inches): 1		
-	No Depth (inches): 0		
	No Depth (inches): 0	Wetland Hydrology Pre	sent? Yes V No
(includes capillary fringe)			
Describe Recorded Data (stream gauge, n	nonitoring well, aeriai priotos, prev	vious irispections), ii available.	
Remarks:			
Wetland hydrology indicat	ors are met.		
Trouble Hydrology Hidrout			

/EGETATION (Four Strata) – Use scientific na	-			Sampling Point: W-PEM-08_2
Tree Stratum (Plot size: 30 ft r		Dominant Species?		Dominance Test worksheet:
1				Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant Species Across All Strata: 2 (B)
3				
5.				Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 0 x 1 = 0
50% of total cover:		= Total Cover		FACW species <u>25</u> x 2 = <u>50</u>
Sapling/Shrub Stratum (Plot size: 15 ft r )	20 /0 01	total cover		FAC species <u>75</u> x 3 = <u>225</u>
1				FACU species <u>0</u> x 4 = <u>0</u>
2.				UPL species <u>0</u> x 5 = <u>0</u>
3.				Column Totals: <u>100</u> (A) <u>275</u> (B)
4.				Prevalence Index = $B/A = 2.75$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				
8				② 3 - Prevalence Index is ≤3.0 <sup>1</sup>
		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )  1. Ambrosia trifida	40	V	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2 Andropogon glomeratus	25		FACW	Definitions of Four Vegetation Strata:
3. Rumex crispus	15		FAC	Deminions of Four Vegetation Strata.
4. Setaria pumila	15		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
5. Cyperus esculentus	5		FAC	more in diameter at breast height (DBH), regardless of height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				<b>Herb</b> – All herbaceous (non-woody) plants, regardless
0				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
T00/ (1/1/ E0		= Total Cov		
50% of total cover: 50 Woody Vine Stratum (Plot size: 30 ft r )	20% of	total cover	: 20	
1				
2				
4.				
5.				Hydrophytic
		= Total Cov	/er	Vegetation
50% of total cover:	20% of	total cover	:	Present? Yes No
Remarks: (If observed, list morphological adaptations bel	ow).			
Dominance test is met.				

SOIL Sampling Point: W-PEM-08\_2

Profile Desc	cription: (Describe	to the depth	needed to docum	nent the i	ndicator	or confirm	the absence of inc	dicators.)
Depth	Matrix			x Features		. 2		_
(inches)	Color (moist)	<u> </u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
-								
				· ———				
	-							
				<u> </u>				
<sup>1</sup> Type: C=C	oncentration, D=Dep	oletion, RM=R	Reduced Matrix, MS	S=Masked	Sand Gr	ains.	<sup>2</sup> Location: PL=F	Pore Lining, M=Matrix.
Hydric Soil	Indicators: (Applic	able to all Li	RRs, unless other	rwise note	ed.)		Indicators for P	roblematic Hydric Soils <sup>3</sup> :
Histosol	(A1)		Polyvalue Be	low Surfac	ce (S8) <b>(I</b>	.RR S, T, U	) 🔲 1 cm Muck (	(A9) <b>(LRR O)</b>
Histic E	pipedon (A2)		Thin Dark Su					(A10) <b>(LRR S)</b>
_	istic (A3)		Loamy Muck	-		R O)		ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)		Loamy Gleye	,	F2)			oodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	T 11\	Depleted Mar		·e)			Bright Loamy Soils (F20)
_	Bodies (A6) (LRR Fucky Mineral (A7) (L		Depleted Dar	,	,		(MLRA 15	Material (TF2)
_	resence (A8) (LRR L		Redox Depre					w Dark Surface (TF12)
_	uck (A9) <b>(LRR P, T)</b>	,	Marl (F10) <b>(L</b>		-,			ain in Remarks)
_	d Below Dark Surfac	e (A11)	Depleted Och		(MLRA 1	51)		,
Thick D	ark Surface (A12)		Iron-Mangan	ese Masse	es (F12) (	LRR O, P,	<b>T)</b> <sup>3</sup> Indicators	of hydrophytic vegetation and
_	rairie Redox (A16) (		_			', U)		hydrology must be present,
_	Mucky Mineral (S1) (	LRR O, S)	Delta Ochric			4505)	unless di	sturbed or problematic.
_	Gleyed Matrix (S4)		Reduced Ver				24)	
_	Redox (S5) d Matrix (S6)						9A) A 149A, 153C, 153I	ח
	rface (S7) <b>(LRR P, S</b>	S, T, U)		ngni Loui	ny cons (	1 20) (III <b>L</b> IV)	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2,
	Layer (if observed)							
Type:								
Depth (in	ches):		<u></u>				Hydric Soil Pres	ent? Yes <u> </u>
Remarks:	•							
Soil san	nnle was not	taken di	ie to injinda	ation F	Hydric	soil as	sumed due t	to prevalence of
	-				-	Jon as		to prevalence of
wetiand	l hydrology a	na nyar	opnytic veg	etatioi	n.			

Project/Site: Golden Triangle Phase 2	City/Cou	<sub>inty:</sub> Lowndes Cou	nty	Sampling Date: 2021-03-11
Applicant/Owner: TVA				Sampling Point: W-PEM-08_UPL
	Section,			
Landform (hillslope, terrace, etc.): Upland, Flat	Local rel	lief (concave, convex,	none): None	Slope (%): 0
Subregion (LRR or MLRA): P 135A	Lat: 33.383592	Long:	-88.614510	Datum: NAD 83
Soil Map Unit Name: Catalpa silty clay			NWI classifica	tion: NA
Are climatic / hydrologic conditions on the site typical for the	nis time of year? Yes	No	(If no, explain in Re	emarks.)
Are Vegetation, Soil, or Hydrology	significantly disturbed	d? Are "Norma	l Circumstances" p	resent? YesNo
Are Vegetation, Soil, or Hydrology	naturally problematic	c? (If needed,	explain any answei	rs in Remarks.)
SUMMARY OF FINDINGS – Attach site map				
Hydrophytic Vegetation Present? Yes	No. V			
Hydrophytic Vegetation Present? Yes Hydric Soil Present? Yes	No Is	s the Sampled Area		4
Wetland Hydrology Present? Yes	No 🔽	vithin a Wetland?	Yes	No
Remarks:				
Upland sample plot adjacent to W-I	PEM-08. The	APT showed	normal cond	ditions at the time
leading up to and during the field si				
leading up to and during the held so	ui vey.			
HYDROLOGY				
Wetland Hydrology Indicators:			Secondary Indica	tors (minimum of two required)
Primary Indicators (minimum of one is required; check al	ll that apply)		Surface Soil	
	ic Fauna (B13)			getated Concave Surface (B8)
	Deposits (B15) <b>(LRR U</b>	I)	Drainage Pat	
	gen Sulfide Odor (C1)		Moss Trim Li	
	ed Rhizospheres alon			Water Table (C2)
	nce of Reduced Iron (		Crayfish Burr	
1 <del></del>	nt Iron Reduction in Til		= 1	sible on Aerial Imagery (C9)
	luck Surface (C7)	, ,	Geomorphic	
	(Explain in Remarks)		Shallow Aqui	
Inundation Visible on Aerial Imagery (B7)			FAC-Neutral	Test (D5)
Water-Stained Leaves (B9)			Sphagnum m	noss (D8) (LRR T, U)
Field Observations:				
Surface Water Present? Yes No D	epth (inches):			
Water Table Present? Yes No D	epth (inches):			
Saturation Present? Yes No D	epth (inches):	Wetland I	Hydrology Presen	t? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well	, aerial photos, previo	ous inspections), if ava	ailable:	
Remarks:				
No indicators are met.				

<b>/EGETATION (Four Strata)</b> – Use scientific na	mes of pl	ants.		Sampling Point: W-PEM-08_UPL
70 ft r		Dominant		Dominance Test worksheet:
<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> ) 1.	% Cover			Number of Dominant Species That Are OBL, FACW, or FAC: 0 (A)
2				Total Number of Dominant
3				Species Across All Strata: 1 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 0 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 0 x 1 = 0
500/ 51 1 1	=			FACW species $0   x 2 = 0$
50% of total cover: Sapling/Shrub Stratum (Plot size: 15 ft r )	20% of	total cover	<u> </u>	FAC species 20 x 3 = 60
				FACU species 90 x 4 = 360
1				UPL species $0   x 5 = 0$
2				Column Totals: 110 (A) 420 (B)
3				
4				Prevalence Index = B/A = 3.82
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7			-	☐ 2 - Dominance Test is >50%
8		Total Cov	/er	☐ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
50% of total cover:				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )	20 /0 01	total cover	•	1
1. Trifolium repens	75	~	FACU	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Cynodon dactylon	15		FACU	Definitions of Four Vegetation Strata:
3. Setaria pumila	15		FAC	
4. Rumex crispus	5		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
5.				height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				Herb – All herbaceous (non-woody) plants, regardless
9.				of size, and woody plants less than 3.28 ft tall.
10.				Woody vine – All woody vines greater than 3.28 ft in
11.				height.
12				
	110% =	= Total Cov	ver .	
50% of total cover: 55	20% of	total cover	: 22	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
	=	Total Cov	ver .	Vegetation Present? Yes No ✓
50% of total cover:	20% of	total cover	:	Fresent: resNo
Remarks: (If observed, list morphological adaptations belo	ow).			
No test is met.				

SOIL Sampling Point: W-PEM-08\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Cd	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> Location: PL:	=Pore Lining, M=Matr	riy
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RRSTI		k (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-		-,		Floodplain Soils (F19	
	Layers (A5)		Depleted Ma		,			s Bright Loamy Soils	
Organic	Bodies (A6) (LRR I	P, T, U)	Redox Dark	Surface (F	<del>-</del> 6)		(MLRA 1	153B)	
5 cm Mu	cky Mineral (A7) (L	.RR P, T, U)	Depleted Dar	rk Surface	e (F7)		Red Paren	nt Material (TF2)	
	esence (A8) (LRR		Redox Depre	essions (F	8)		<u></u> Very Shall	ow Dark Surface (TF	12)
	ck (A9) (LRR P, T)						U Other (Exp	olain in Remarks)	
	Below Dark Surfa	ce (A11)	Depleted Ocl				3		
	ark Surface (A12)		Iron-Mangan					rs of hydrophytic vege	
	rairie Redox (A16) (					, U)		d hydrology must be p	
	lucky Mineral (S1)	(LKK 0, 5)	Delta Ochric			OA 150D)		disturbed or problema	atic.
	edox (S5)		Reduced Ver						
	Matrix (S6)						+9A) RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,	S. T. U)	Anomalous L	night Loai	illy Solis (	1 20) (IVILIV	A 149A, 133C, 13	30)	
	_ayer (if observed								
Type:		•							
, , <u> </u>	ches):						Hydric Soil Pre	esent? Yes	No 🗸
Remarks:							Tryuno com Tro		
No indic	ators are m	et.							

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-09
• • • • • • • • • • • • • • • • • • • •	ion, Township, Range: S33 T18N R16E
Landform (hillslope, terrace, etc.): Depression Loca	
Subregion (LRR or MLRA): P 135A Lat: 33.38941	
Soil Map Unit Name: Brooksville silty clay, 1 to 3 percent slopes	NWI classification: NA
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distu	
Are Vegetation, Soil, or Hydrology naturally problem	
	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	
Hydric Soil Present? Yes V No	Is the Sampled Area
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	
Sample plot in W-PEM-09. The APT showed n	ormal conditions at the time leading up to and
during the field survey.	orman contained at the time reading up to and
during the field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Secondary Indicators (Infill Indicators (Infill Indicators)  Surface Soil Cracks (B6)
Surface Water (A1)  Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  High Water Table (A2)  Marl Deposits (B15) (LR	
Saturation (A3)  Hydrogen Sulfide Odor	
Water Marks (B1)  Oxidized Rhizospheres	` '
Sediment Deposits (B2) Presence of Reduced In	
Drift Deposits (B3)	_ ` ` · · · · · · · · <del>_</del> · · · · · · · · · · · · · · · · · · ·
Algal Mat or Crust (B4)	☑ Geomorphic Position (D2)
Iron Deposits (B5)	rks) Shallow Aquitard (D3)
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes No Depth (inches):	
Water Table Present? Yes No Depth (inches):	
Saturation Present? Yes No Depth (inches): (includes capillary fringe)	Wetland Hydrology Present? Yes No
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	evious inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	

<b>'EGETATION (Four Strata)</b> – Use scientific na	mes of pl	ants.		Sampling Point: W-PEM-09
Total Objections (Dietoine 30 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r ) 1.		Species?	Status	Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4 5				Percent of Dominant Species That Are ORL FACW or FAC: 100 (A/R)
6.				That file OBE, I flow, SI I flow
7				Prevalence Index worksheet:
8				Total % Cover of:         Multiply by:           OBL species         10         x 1 = 10
500/ of Astal assum		Total Cov		FACW species $\frac{15}{x^2}$ $x = \frac{30}{x^2}$
50% of total cover: Sapling/Shrub Stratum (Plot size: 15 ft r )	20% of	total cover		FAC species 60 x 3 = 180
1. Fraxinus pennsylvanica	10	~	FACW	FACU species <u>0</u>
2.				UPL species $0 \times 5 = 0$
3.				Column Totals: <u>85</u> (A) <u>220</u> (B)
4				Prevalence Index = $B/A = 2.59$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7	·			2 - Dominance Test is >50%
8	10% -	Total Cov		✓ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
50% of total cover: 5				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Ambrosia trifida	60		FAC	be present, unless disturbed or problematic.
2. Persicaria hydropiperoides	10		OBL	Definitions of Four Vegetation Strata:
3. Fraxinus pennsylvanica	5		FACW	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4				more in diameter at breast height (DBH), regardless of height.
5 6.				
6 7				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				Herb – All herbaceous (non-woody) plants, regardless
9.				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12	750/			
50% of total cover: 38		= Total Cov		
Woody Vine Stratum (Plot size: 30 ft r )	20% 01	total cover		
1				
2.				
3				
4				
5				Hydrophytic
		= Total Cov		Vegetation Present?  Yes No
50% of total cover:		total cover	·	
Remarks: (If observed, list morphological adaptations belo	OW).			
Dominance test is met.				

SOIL Sampling Point: W-PEM-09

Profile Des	cription: (Describe to the dept	h needed to docui	ment the	indicator o	or confirm	the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
-							
_							'
			_				_
	·		_				
-							
							_
l — -	·						
<sup>1</sup> Type: C=C	oncentration, D=Depletion, RM=	Reduced Matrix, M	S=Maske	ed Sand Gra	ains.	<sup>2</sup> Location: PL=F	Pore Lining, M=Matrix.
	Indicators: (Applicable to all I						roblematic Hydric Soils <sup>3</sup> :
Histosol	(A1)	Polyvalue Be	elow Surfa	ace (S8) <b>(L</b> l	RR S. T. L	J)	(A9) (LRR O)
I <del></del>	pipedon (A2)	Thin Dark Su					(A10) (LRR S)
1 <del>1 - 1</del>	istic (A3)	Loamy Muck					ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)	Loamy Gleye	-	. , .	٠,		oodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	Depleted Ma		(1 2)			Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark		(F6)		(MLRA 15	. , ,
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da	,	,		,	Material (TF2)
	resence (A8) (LRR U)	Redox Depre					w Dark Surface (TF12)
	uck (A9) (LRR P, T)	Marl (F10) (L		٠,			ain in Remarks)
_	d Below Dark Surface (A11)	Depleted Oc		) (MI RA 15	i1)	Outer (Expire	an in remarks)
	ark Surface (A12)	Iron-Mangan				T) <sup>3</sup> Indicators	of hydrophytic vegetation and
	rairie Redox (A16) <b>(MLRA 150A</b>					•	nydrology must be present,
_	Mucky Mineral (S1) (LRR O, S)	Delta Ochric			• • •		sturbed or problematic.
	Gleyed Matrix (S4)	Reduced Ve			0A 150B)		otalized of problematic.
_	Redox (S5)	Piedmont Flo					
	d Matrix (S6)					A 149A, 153C, 153I	מס
	urface (S7) (LRR P, S, T, U)	/ tilolilalodo I	ongni Loc	anny conc (i	20) <b>(2</b> .(		-,
	Layer (if observed):					1	
Type:							
, , <u> </u>	-1 >					Uhadala Oali Baas	
	ches):					Hydric Soil Pres	ent? Yes No
Remarks:							
No soil	sample taken due to	proximity to	o unde	ergroun	ıd natu	ıral qas pipel	lines. Hydric soil is
		-		_			
assume	d due to prevalence	or wetland	nyaro	logy in	aicato	rs and nydro	pnytic vegetation.

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-11
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PEM-09_UPL
Investigator(s): J. Brown, O. Haney Sect	tion, Township, Range:
	al relief (concave, convex, none): Concave Slope (%):
, ,	25 Long: -88.6267049 Datum: NAD 83
Soil Map Unit Name:	NWI classification:
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distu	
Are Vegetation, Soil, or Hydrology naturally problem	
	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	
Hydric Soil Present? Yes No	Is the Sampled Area
Hydric Soil Present?  Wetland Hydrology Present?  Remarks:	within a Wetland? Yes No
The APT showed normal conditions at the time	e leading up to and during the field survey.
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1)  High Water Table (A2)  Aquatic Fauna (B13)  Marl Deposits (B15) (LR	Sparsely Vegetated Concave Surface (B8)  Drainage Patterns (B10)
Saturation (A3)  Hydrogen Sulfide Odor	
	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced In	
☐ Drift Deposits (B3) ☐ Recent Iron Reduction in	n Tilled Soils (C6) Saturation Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)  Thin Muck Surface (C7)	
☐ Iron Deposits (B5) ☐ Other (Explain in Remai	rks)
Inundation Visible on Aerial Imagery (B7)  Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	<u> </u>
Surface Water Present? Yes No Depth (inches):	
Water Table Present? Yes No Depth (inches):	
Saturation Present? Yes No Depth (inches):	Wetland Hydrology Present? Yes No _✔
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	evious inspections), if available:
Remarks:	

Dominance Test worksheet:   Number of Dominant Species   That Are OBL, FACW, or FAC:   0
Total Number of Dominant Species
Total Number of Dominant Species Across All Strata:    Percent of Dominant Species   That Are OBL, FACW, or FAC: NaN
Percent of Dominant Species   That Are OBL, FACW, or FAC:   NaN   (A/B)
Prevalence Index worksheet:   Total % Cover of:
Prevalence Index worksheet:   Total % Cover of:
Total % Cover of:
OBL species
FACW species 0
FACW species $\frac{0}{0}$ $x = \frac{0}{0}$ FAC species $\frac{0}{0}$ $x = \frac{0}{0}$ FACU species $\frac{0}{0}$ $x = \frac{0}{0}$ UPL species $\frac{0}{0}$ $x = \frac{0}{0}$ Column Totals: $\frac{0}{0}$ (A) $\frac{0}{0}$ (B)  Prevalence Index = B/A = $\frac{nan}{0}$ Hydrophytic Vegetation Indicators: $\frac{0}{0}$ 1 - Rapid Test for Hydrophytic Vegetation
FAC species
UPL species $0$ $x = 0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$ $0$
Column Totals: 0 (A) 0 (B)  Prevalence Index = B/A = nan  Hydrophytic Vegetation Indicators:  1 - Rapid Test for Hydrophytic Vegetation
Prevalence Index = B/A = nan  Hydrophytic Vegetation Indicators:  1 - Rapid Test for Hydrophytic Vegetation
Prevalence Index = B/A = nan  Hydrophytic Vegetation Indicators:  1 - Rapid Test for Hydrophytic Vegetation
Hydrophytic Vegetation Indicators:  1 - Rapid Test for Hydrophytic Vegetation
1 - Rapid Test for Hydrophytic Vegetation
D 2 Deminance Test is \$ 500/
—— ——— <u> </u>
$\square$ 3 - Prevalence Index is $\leq 3.0^1$
l Cover Problematic Hydrophytic Vegetation¹ (Explain)
cover:
<sup>1</sup> Indicators of hydric soil and wetland hydrology must
be present, unless disturbed or problematic.
Definitions of Four Vegetation Strata:
Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
more in diameter at breast height (DBH), regardless of height.
Senting/Shuth Woody plants evaluating visce less
Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
Woody vine - All woody vines greater than 3.28 ft in
Woody vine – All woody vines greater than 3.28 ft in height.
l Cover
cover:
<del></del>
Hydrophytic
Cover   Vegetation   Present?   Yes No
cover:
a c

SOIL Sampling Point: W-PEM-09\_UPL

ype: C=Concentration, D=Depletion, RM=Reduced Matrix, MS=Masked Sand Grains.	Depth _	Matrix	Redox Features	<del>-</del> 1 · 2		
Histosol (A1)	(inches)	Color (moist) %	Color (moist) %	Type <sup>1</sup> Loc <sup>2</sup>	Texture	Remarks
Histosol (A1)						
Histosol (A1)	-					
Histosol (A1)						
Histosol (A1)	<del></del> -					
Histosol (A1)						
Histosol (A1)	-					
Histosol (A1)						
Histosol (A1)	<del></del> -					
Histosol (A1)						
Histosol (A1)					<sup>2</sup> Location: PL=	Pore Lining, M=Matrix.
Histic Epipedon (A2) Black Histic (A3) Hydrogen Sulfide (A4) Stratified Layers (A5) Coamy Gleyed Matrix (F2) Depleted Matrix (F3) Depleted Matrix (F3) Depleted Dark Surface (F7) Depleted Dark Surface (F7) Depleted Below Dark Surface (A1) Depleted Dark Surface (F13) (LRR P, T, U) Depleted Below Dark Surface (A1) Depleted Dark Surface (F13) (LRR P, T, U) Depleted Dark Surface (F14) (MLRA 151) Depleted Below Dark Surface (A12) Depleted Depleted Ochric (F17) (MLRA 151) Depleted Below Dark Surface (A12) Depleted Ochric (F13) (LRR P, T, U) Sandy Mucky Mineral (S1) (LRR O, S) Delta Ochric (F17) (MLRA 151) Sandy Gleyed Matrix (S4) Depleted Vertic (F18) (MLRA 150A, 150B) Sandy Redox (S5) Dark Surface (S7) (LRR P, S, T, U) Destrictive Layer (if observed): Type: Depth (inches):  Thin Dark Surface (S9) (LRR S, T, U) Depleted Matrix (F10) (LRR S) Deadwarts (F13) (LRR P, T, U) Depleted Matrix (F18) (MLRA 150A) Dark Surface (F18) (MLRA 150A, 150B) Dark Surface (S7) (LRR P, S, T, U)  Betrictive Layer (if observed): Type: Depth (inches):  Type: Depth (inches): Dark Surface (F10) (IRR S, T, U) Depleted Matrix (F2) Depth (inches): Dark Surface (F10) (IRR P, S, T, U) Depleted Depleted Matrix (F2) Depth (inches): Dark Surface (F10) (IRR P, S, T, U) Depleted Deplet	ydric Soil Inc	dicators: (Applicable to all	RRs, unless otherwise note	d.)	Indicators for	Problematic Hydric Soils <sup>3</sup> :
Black Histic (A3)	Histosol (A	1)	Polyvalue Below Surface	e (S8) <b>(LRR S, T, U</b>	l) <u> </u>	(A9) <b>(LRR O)</b>
Hydrogen Sulfide (A4)  Stratified Layers (A5)  Organic Bodies (A6) (LRR P, T, U)  Scr Mucky Mineral (A7) (LRR P, T, U)  Depleted Dark Surface (F6)  Muck Presence (A8) (LRR U)  1 cm Muck (A9) (LRR P, T)  Depleted Below Dark Surface (A11)  Thick Dark Surface (A12)  Coast Prairie Redox (A16) (MLRA 150A)  Sandy Mucky Mineral (S1) (LRR O, S)  Sandy Redox (S5)  Sandy Redox (S5)  Depleted Dark Surface (F1)  Marl (F10) (LRR U)  Depleted Ochric (F11) (MLRA 151)  Depleted Ochric (F13) (LRR P, T, U)  Depleted Prairie Redox (A16) (MLRA 150A)  Depleted Ochric (F17) (MLRA 151)  Depleted Redox (A16) (MLRA 150A)  Sandy Mucky Mineral (S1) (LRR O, S)  Sandy Redox (S5)  Delta Ochric (F17) (MLRA 150A, 150B)  Piedmont Floodplain Soils (F20) (MLRA 149A)  Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)  Petricive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No	Histic Epip	edon (A2)	Thin Dark Surface (S9)	(LRR S, T, U)	2 cm Muck	(A10) <b>(LRR S)</b>
Stratified Layers (A5)	Black Histi	c (A3)	Loamy Mucky Mineral (F	F1) <b>(LRR O)</b>	Reduced V	ertic (F18) (outside MLRA 150A,I
Organic Bodies (A6) (LRR P, T, U) Redox Dark Surface (F6) (MLRA 153B)  5 cm Mucky Mineral (A7) (LRR P, T, U) Depleted Dark Surface (F7) Redox Depressions (F8)  Muck Presence (A8) (LRR U) Redox Depressions (F8) Very Shallow Dark Surface (TF12)  1 cm Muck (A9) (LRR P, T) Marl (F10) (LRR U) Other (Explain in Remarks)  Depleted Below Dark Surface (A11) Depleted Ochric (F11) (MLRA 151)  Thick Dark Surface (A12) Iron-Manganese Masses (F12) (LRR O, P, T) Indicators of hydrophytic vegetation and Coast Prairie Redox (A16) (MLRA 150A) Umbric Surface (F13) (LRR P, T, U) wetland hydrology must be present, unless disturbed or problematic.  Sandy Mucky Mineral (S1) (LRR O, S) Delta Ochric (F17) (MLRA 151) unless disturbed or problematic.  Sandy Redox (S5) Reduced Vertic (F18) (MLRA 150A, 150B)  Sandy Redox (S5) Piedmont Floodplain Soils (F19) (MLRA 149A)  Stripped Matrix (S6) Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)  Dark Surface (S7) (LRR P, S, T, U)  Setrictive Layer (if observed):  Type:  Depth (inches): Hydric Soil Present? Yes No	Hydrogen	Sulfide (A4)	Loamy Gleyed Matrix (F	2)		
5 cm Mucky Mineral (A7) (LRR P, T, U)			Depleted Matrix (F3)		Anomalous	s Bright Loamy Soils (F20)
Muck Presence (A8) (LRR U) Redox Depressions (F8) Very Shallow Dark Surface (TF12)  1 cm Muck (A9) (LRR P, T) Marl (F10) (LRR U) Other (Explain in Remarks)  Depleted Below Dark Surface (A11) Depleted Ochric (F11) (MLRA 151)  Thick Dark Surface (A12) Iron-Manganese Masses (F12) (LRR O, P, T) Iron-Manganese Masses (F12) (LRR O, P, T) Iron-Manganese Masses (F13) (LRR P, T, U) Wetland hydrology must be present, unless disturbed or problematic.  Sandy Mucky Mineral (S1) (LRR O, S) Delta Ochric (F17) (MLRA 151) unless disturbed or problematic.  Sandy Gleyed Matrix (S4) Reduced Vertic (F18) (MLRA 150A, 150B)  Sandy Redox (S5) Piedmont Floodplain Soils (F19) (MLRA 149A)  Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)  Setrictive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No			=	•	1 1 '	•
1 cm Muck (A9) (LRR P, T)						, ,
Depleted Below Dark Surface (A11)  Depleted Ochric (F11) (MLRA 151)  Iron-Manganese Masses (F12) (LRR O, P, T)  Coast Prairie Redox (A16) (MLRA 150A)  Umbric Surface (F13) (LRR P, T, U)  Sandy Mucky Mineral (S1) (LRR O, S)  Sandy Gleyed Matrix (S4)  Sandy Redox (S5)  Stripped Matrix (S6)  Dark Surface (S7) (LRR P, S, T, U)  Setrictive Layer (if observed):  Type:  Depth (inches):  Iron-Manganese Masses (F12) (LRR O, P, T)  3Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Poelta Ochric (F17) (MLRA 150A)  Wetland hydrology must be present, unless disturbed or problematic.  Poelta Ochric (F17) (MLRA 150A)  Wetland hydrology must be present, unless disturbed or problematic.  No Vetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Wetland hydrology must be present, unless disturbed or problematic.  Poelta Ochric (F17) (MLRA 151)  Wetland hydrology must be present, unless disturbed or problematic.  Poelta Ochric (F17) (MLRA 150A)  Wetland hydrology must be present, unless disturbed or problematics.			_	)		· · ·
Thick Dark Surface (A12)	_		= ` '` '		U Other (Exp	lain in Remarks)
Coast Prairie Redox (A16) (MLRA 150A) Umbric Surface (F13) (LRR P, T, U) wetland hydrology must be present, bela Ochric (F17) (MLRA 151) unless disturbed or problematic.  Sandy Gleyed Matrix (S4) Reduced Vertic (F18) (MLRA 150A, 150B)  Sandy Redox (S5) Piedmont Floodplain Soils (F19) (MLRA 149A)  Stripped Matrix (S6) Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)  Park Surface (S7) (LRR P, S, T, U)  Pestrictive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No						
Sandy Mucky Mineral (S1) (LRR O, S)  Delta Ochric (F17) (MLRA 151)  Reduced Vertic (F18) (MLRA 150A, 150B)  Sandy Redox (S5)  Stripped Matrix (S6)  Dark Surface (S7) (LRR P, S, T, U)  Destrictive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No					•	
Sandy Gleyed Matrix (S4) Reduced Vertic (F18) (MLRA 150A, 150B)  Sandy Redox (S5) Piedmont Floodplain Soils (F19) (MLRA 149A)  Stripped Matrix (S6) Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)  Dark Surface (S7) (LRR P, S, T, U)  estrictive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No						
Sandy Redox (S5)   Piedmont Floodplain Soils (F19) (MLRA 149A)   Stripped Matrix (S6)   Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)   Dark Surface (S7) (LRR P, S, T, U)   estrictive Layer (if observed): Type:   Depth (inches):   Hydric Soil Present? Yes No V	_					disturbed or problematic.
Stripped Matrix (S6) Dark Surface (S7) (LRR P, S, T, U)  strictive Layer (if observed):  Type: Depth (inches):  Hydric Soil Present? Yes No	_	-				
Dark Surface (S7) (LRR P, S, T, U)  estrictive Layer (if observed):  Type:  Depth (inches):  Hydric Soil Present? Yes No						20)
Depth (inches): Hydric Soil Present? Yes No	_ ::	, ,	Anomalous Bright Loam	y 50115 (F20) (WILK	A 149A, 153C, 15	30)
Type:  Depth (inches):						
Depth (inches): No						
	• •		<del></del> ,			
marks:	Depth (inch	es):			Hydric Soil Pre	sent? Yes No
	temarks:					

Project/Site: Golden Triangle Pl	hase 2	Citv/C	ounty: Lowndes C	ounty	Sampling Date: 2021-03-11
Applicant/Owner: TVA					Sampling Point: W-PEM-10
Investigator(s): J. Brown, O. Hai	ney	Section Section	on Townshin Range		Company Cont.
Landform (hillslope, terrace, etc.):	Depression	Local	rollof (concavo, conv	ox nono): Concave	Slope (%): 3
Subregion (LRR or MLRA): P 135		Lat: 33.382376	•	· -	Datum: NAD 83
		Lat:		•	
Soil Map Unit Name: Catalpa silt				NWI classifica	
Are climatic / hydrologic conditions					_
Are Vegetation, Soil	_, or Hydrology	significantly distur	bed? Are "Nor	mal Circumstances" p	resent? YesNo
Are Vegetation, Soil	, or Hydrology	naturally problema	atic? (If neede	ed, explain any answe	rs in Remarks.)
SUMMARY OF FINDINGS -	- Attach site ma	ap showing sam	pling point loca	ations, transects	, important features, etc.
Hydrophytic Vegetation Present?	Yes 🗸	No	la tha Cammiad Am	_	
Hydric Soil Present?		No	Is the Sampled Are within a Wetland?		No
Wetland Hydrology Present?		No	within a wettand?	res	NO
Remarks:					
Sample plot in W-PEN	<b>∕</b> I-10. The AP	T showed no	rmal conditio	ns at the time	e leading up to and
during the field surve					<b>5</b> 1
during the held surve	у.				
HYDROLOGY					
Wetland Hydrology Indicators:				Socondary Indica	tors (minimum of two required)
Primary Indicators (minimum of or	ne is required: check	all that apply)		Secondary indica	
Surface Water (A1)		atic Fauna (B13)			getated Concave Surface (B8)
High Water Table (A2)		l Deposits (B15) <b>(LRF</b>	s m	Drainage Pat	
Saturation (A3)		rogen Sulfide Odor (0		Moss Trim Li	
Water Marks (B1)		lized Rhizospheres a			Water Table (C2)
Sediment Deposits (B2)		sence of Reduced Iro		Crayfish Burr	
Drift Deposits (B3)		ent Iron Reduction in			sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)		Muck Surface (C7)		Geomorphic	
Iron Deposits (B5)	☐ Oth€	er (Explain in Remark	s)	Shallow Aqui	tard (D3)
Inundation Visible on Aerial Ir	nagery (B7)			FAC-Neutral	Test (D5)
☐ Water-Stained Leaves (B9)				Sphagnum m	noss (D8) <b>(LRR T, U)</b>
Field Observations:					
	es No				
	es <u> </u>				•
Saturation Present? Ye (includes capillary fringe)	es No	Depth (inches): 0	Wetlar	nd Hydrology Presen	t? Yes / No
Describe Recorded Data (stream	gauge, monitoring w	ell, aerial photos, pre	vious inspections), if	available:	
Remarks:					
Wetland hydrology in	dicators are	met.			

1. Alternanthera philoxeroides 10 V OBL

 $_{3.}$  Xanthium strumarium 10  $\checkmark$  FAC 4. Persicaria hydropiperoides 5 OBL

Sapling/Shrub Stratum (Plot size: 15 ft r )

Herb Stratum (Plot size: 5 ft r )

2. Juncus effusus

Absolute Dominant Indicator

% Cover Species? Status

\_\_\_\_\_ = Total Cover

= Total Cover

10

35% = Total Cover

\_\_\_\_\_= Total Cover

50% of total cover: 18 20% of total cover: 7

50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_

OBL

50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_

50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_\_

Sampling Point: W-PEM-10					
Dominance Test worksheet:  Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)					
Total Number of Dominant Species Across All Strata: 3 (B)					
Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B)					
Prevalence Index worksheet:           Total % Cover of:         Multiply by:           OBL species         25         x 1 = 25           FACW species         0         x 2 = 0           FAC species         10         x 3 = 30           FACU species         0         x 4 = 0           UPL species         0         x 5 = 0           Column Totals:         35         (A)         55         (B)					
Prevalence Index = B/A = 1.57					
Hydrophytic Vegetation Indicators:  1 - Rapid Test for Hydrophytic Vegetation  2 - Dominance Test is >50%  3 - Prevalence Index is ≤3.0¹  Problematic Hydrophytic Vegetation¹ (Explain)  ¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.  Definitions of Four Vegetation Strata:  Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of height.  Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.  Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.  Woody vine – All woody vines greater than 3.28 ft in					
Woody vine – All woody vines greater than 3.28 ft in height.  Hydrophytic Vegetation Present? Yes No					

Remarks: (If observed, list morphological adaptations below).

Woody Vine Stratum (Plot size: 30 ft r )

Dominance test is met. More than 40% coverage of open water.

SOIL Sampling Point: W-PEM-10

Profile Des	cription: (Describe to the depti	n needed to docur	ment the	indicator of	or confirm	the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	%	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
-							
l — —						-	_
			_				
-							
			-				
l ———							
			_				
_							
17		D = -ll	0 Ml			21	Daniel Linkson M. Mateix
	oncentration, D=Depletion, RM=I Indicators: (Applicable to all L				ains.		Pore Lining, M=Matrix.  Problematic Hydric Soils <sup>3</sup> :
l <u> </u>							•
Histosol	'	Polyvalue Be				. —	
I <b>=</b>	pipedon (A2)	Thin Dark Su					(A10) <b>(LRR S)</b>
_	istic (A3)	Loamy Muck	-	. , .	O)		ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)	Loamy Gleye		(F2)			loodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	Depleted Ma					Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark	,	,		(MLRA 15	•
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da		, ,			Material (TF2)
	resence (A8) (LRR U)	Redox Depre		<del>-</del> 8)			w Dark Surface (TF12)
_	uck (A9) <b>(LRR P, T)</b>	Marl (F10) <b>(L</b>				U Other (Expla	ain in Remarks)
	d Below Dark Surface (A11)	Depleted Oc					
	ark Surface (A12)	Iron-Mangan				•	of hydrophytic vegetation and
_	rairie Redox (A16) (MLRA 150A)	_			, U)		hydrology must be present,
_	Mucky Mineral (S1) (LRR O, S)	Delta Ochric					isturbed or problematic.
_	Gleyed Matrix (S4)	Reduced Ve					
	Redox (S5)	Piedmont Flo					
	d Matrix (S6)	Anomalous E	Bright Loa	amy Soils (F	-20) <b>(MLR</b>	A 149A, 153C, 153	D)
	ırface (S7) (LRR P, S, T, U)					1	
Restrictive	Layer (if observed):						
Type:							
Depth (in	ches):	<u></u>				Hydric Soil Pres	ent? Yes <u>'</u> No
Remarks:							
No soil	sampla was takan di	ıo to inunda	tion L	Judro	oil ic a	secumed due	to the providence of
NO SOII :	sample was taken di	ie to inunua	tion. F	Tyuro s	oui is a	issumed due	to the prevalence of
wetland	l hydrology and hydr	ophytic ved	etatio	n.			
	, , .	·	,				

Project/Site: Golden Triangle Phase	2	Citv/C	ounty: Lowndes	County	Sampling Date: 2021-03-09
Applicant/Owner: TVA					Sampling Point: W-PFO-01
Investigator(s): J. Brown, O. Haney		Section	n Townshin Range	e: S26 T18N R16E	
Landform (hillslope, terrace, etc.): Depr					Slone (%): 0
Subregion (LRR or MLRA): P 135A					Datum: NAD 83
Soil Map Unit Name: Vaiden silty cla	v. 2 to 5 perce	_ Lat. ent slopes, erode		NWI classifica	
Are climatic / hydrologic conditions on th					
Are Vegetation, Soil, or h					oresent? YesNo
Are Vegetation, Soil, or h	-lydrology	_ naturally problema	itic? (If need	ed, explain any answe	rs in Remarks.)
SUMMARY OF FINDINGS – At	tach site ma	p showing sam	pling point loc	ations, transects	, important features, etc.
Hydrophytic Vegetation Present?	Yes	No	lo the Compled A		
Hydric Soil Present?	Yes 🗸	No	Is the Sampled A within a Wetland?		No
Wetland Hydrology Present?	Yes	No	within a wetland	163	
Remarks:					
Sample plot in W-PFO-0	1. The APT	showed no	rmal condition	ons at the time	leading up to and
during the field survey.					
,					
HYDROLOGY					
Wetland Hydrology Indicators:				Secondary Indica	tors (minimum of two required)
Primary Indicators (minimum of one is	required; check a	all that apply)		Surface Soil	
Surface Water (A1)	Aqua	tic Fauna (B13)			getated Concave Surface (B8)
High Water Table (A2)		Deposits (B15) (LRF	R U)	Drainage Pa	
Saturation (A3)	☐ Hydro	ogen Sulfide Odor (0	21)	Moss Trim Li	nes (B16)
Water Marks (B1)		zed Rhizospheres a			Water Table (C2)
Sediment Deposits (B2)		ence of Reduced Iro		Crayfish Buri	
Drift Deposits (B3)		nt Iron Reduction in	Tilled Soils (C6)		sible on Aerial Imagery (C9)
Algal Mat or Crust (B4) Iron Deposits (B5)		Muck Surface (C7) r (Explain in Remark	e)	Shallow Aqui	Position (D2)
Inundation Visible on Aerial Image		(Explain in Nemark	3)	FAC-Neutral	` '
☐ Water-Stained Leaves (B9)	., (= . ,			_	noss (D8) <b>(LRR T, U)</b>
Field Observations:					
Surface Water Present? Yes	✓ No [	Depth (inches): 2			
Water Table Present? Yes	✓ No [	Depth (inches): 0			
	<u>✓</u> No [	Depth (inches): 0	Wetla	nd Hydrology Presen	t? Yes <u>/</u> No
(includes capillary fringe)  Describe Recorded Data (stream gauge	e, monitoring we	II, aerial photos, pre	vious inspections), i	f available:	
33	, <b>.</b> .	,	,,		
Remarks:					
Wetland hydrology indicate	ators are r	net.			

,	ames of pl	ants.		Sampling Point: W-PFO-0
For a Ottostoria (Dietoina 30 ft r		Dominant		Dominance Test worksheet:
ree Stratum (Plot size: 30 ft r ) Salix nigra	25	Species? ✓	OBL	Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)
·				That Are OBL, FACW, or FAC: 3 (A)
<u>.                                    </u>				Total Number of Dominant
3				Species Across All Strata: 3 (B)
l				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B
S				Prevalence Index worksheet:
				Total % Cover of: Multiply by:
8				OBL species 40 x 1 = 40
40		= Total Cov		FACW species $0 \times 2 = 0$
50% of total cover: 18	20% of	total cover		FAC species 10 x 3 = 30
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 0 x 4 = 0
Salix nigra	5		OBL	
)				· — — —
3				Column Totals: <u>50</u> (A) <u>70</u> (B)
l				Prevalence Index = B/A = 1.4
5				Hydrophytic Vegetation Indicators:
3				1 - Rapid Test for Hydrophytic Vegetation
<b>7</b>				2 - Dominance Test is >50%
3				☑ 3 - Prevalence Index is ≤3.0¹
	5% :	= Total Cov	er	Problematic Hydrophytic Vegetation¹ (Explain)
50% of total cover: 3	20% of	total cover	1	Troblematic Trydrophytic Vegetation (Explain)
Herb Stratum (Plot size: 5 ft r )				1 adjustence of burdence of least model and burdence of burden
Cyperus esculentus	10	~	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2				Definitions of Four Vegetation Strata:
3.				
				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) o
l				more in diameter at breast height (DBH), regardless of height.
5				
S				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7.				than 6 m. BBT and greater than 6.25 tt (1 m) tail.
3			•	Herb – All herbaceous (non-woody) plants, regardless
)				of size, and woody plants less than 3.28 ft tall.
0				Woody vine – All woody vines greater than 3.28 ft in
1				height.
2				
_		= Total Cov		
50% of total cover: 5	20% of	total cover	2	
Noody Vine Stratum (Plot size: 30 ft r )				
l				
2				
3				
1				
5				Hydrophytic
	:	= Total Cov	er	Vegetation
	20% of	total cover		Present? Yes No
50% of total cover:	20 /0 01		·	

SOIL Sampling Point: W-PF0-01

(inches)	Matrix Color (moist)	%	Redox Features Color (moist) % Type <sup>1</sup> Loc <sup>2</sup>			Texture	Remarks
0 - 24	10YR 4/2	_ <u>%</u>	7.5YR 5/6	_ <del>%</del>		Clay Loam	Remarks
0 24	101K 4/2		7.51K 3/0	_ 25 _ (	<u> </u>	Clay Loain	
-							
_							
		<del>-</del>					
		<del>-</del>				<del></del>	
	oncentration, D=De						L=Pore Lining, M=Matrix.
<u>-</u>	ndicators: (Appli	cable to all I			-		or Problematic Hydric Soils <sup>3</sup> :
Histosol					(S8) (LRR S, T,		ck (A9) <b>(LRR O)</b>
Black Hi	pipedon (A2)			urface (S9) <b>(L</b> ky Mineral (F <i>*</i>			ck (A10) <b>(LRR S)</b>   Vertic (F18) <b>(outside MLRA 150A,I</b>
=	n Sulfide (A4)		=	red Matrix (F2		7 7 7	t Floodplain Soils (F19) (LRR P, S, T
	Layers (A5)		Depleted M		,		us Bright Loamy Soils (F20)
	Bodies (A6) (LRR I	P, T, U)		Surface (F6)		(MLRA	
	cky Mineral (A7) (L		Depleted Da	ark Surface (F	7)		ent Material (TF2)
Muck Pr	esence (A8) (LRR	U)	Redox Depi	ressions (F8)			allow Dark Surface (TF12)
=	ck (A9) (LRR P, T)		Marl (F10) (	•		U Other (E	xplain in Remarks)
=	Below Dark Surfa	ce (A11)	= :	chric (F11) <b>(M</b>	•	3	
	irk Surface (A12)	MI DA 450A			(F12) <b>(LRR O, I</b>		ors of hydrophytic vegetation and
	airie Redox (A16) ( lucky Mineral (S1)		· <del>-</del>	ace (F13) <b>(LF</b>			nd hydrology must be present, s disturbed or problematic.
=	leyed Matrix (S4)	(LKK U, 3)		c (F17) <b>(MLR</b> /	-RA 150A, 150E		s disturbed of problematic.
_	edox (S5)				s (F19) <b>(MLRA</b> 1		
_	Matrix (S6)					, .RA 149A, 153C, 1	53D)
= ::	face (S7) (LRR P,	S, T, U)	_	,	` ',		•
estrictive l	ayer (if observed	):					
Type:							
Depth (inc	ches):					Hydric Soil P	resent? Yes 🚩 No
emarks:							
lydric s	oil indicator	F3 is m	et				
iyanc 3	on maicator	1 5 15 111	Ct.				

Project/Site: Golden Triangle Phase 2 City/	County: Lowndes County Sampling Date: 2021-03-09
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PFO-01_UPL
Investigator(s): J. Brown, O. Haney Sec	tion, Township, Range: S26 T18N R16E
Landform (hillslope, terrace, etc.): Hillslope Local	
Subregion (LRR or MLRA): P 135A Lat: 33.39939	,
Soil Map Unit Name: Vaiden silty clay, 2 to 5 percent slopes, eroo	
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distu	
Are Vegetation, Soil, or Hydrology naturally problem	
	mpling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	.1
Upland sample plot adjacent to W-PFO-01. Th	APT showed normal conditions at the time
	e Ai i showed normal conditions at the time
leading up to and during the field survey.	
HYDDOLOGY	
HYDROLOGY	Connedon Indicators (minimum of two required)
Wetland Hydrology Indicators:  Primary Indicators (minimum of one is required; check all that apply)	Secondary Indicators (minimum of two required)
	Surface Soil Cracks (B6)
Surface Water (A1)  High Water Table (A2)  Aquatic Fauna (B13)  Marl Deposits (B15) (LF	Sparsely Vegetated Concave Surface (B8)  RR U)  Drainage Patterns (B10)
Saturation (A3)  Hydrogen Sulfide Odor	
	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced In	ron (C4)
Drift Deposits (B3)	in Tilled Soils (C6) Saturation Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	,
☐ Iron Deposits (B5) ☐ Other (Explain in Rema	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9) Field Observations:	☐ Sphagnum moss (D8) (LRR T, U)
Surface Water Present? Yes No _ Depth (inches):	
Water Table Present?  Yes No _ Depth (inches):	
Saturation Present?  Yes No _ Depth (inches):	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pr	'evious inspections), if available:
Remarks:	
No indicators are met.	
indicators are met.	

20 #		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species
1. Juniperus virginiana	60		FACU	That Are OBL, FACW, or FAC: $\frac{1}{}$ (A)
Celtis laevigata     Celtis laevigata	5		FACW	Total Number of Dominant Species Across All Strata: 2 (B)
4				
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 50 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 0 x 1 = 0
		Total Cov		FACW species $\frac{5}{5}$ $\times 2 = \frac{10}{10}$
50% of total cover: 33	20% of	total cover	13	FAC species 2 x 3 = 6
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 60 x 4 = 240
1				
2				
3				Column Totals: <u>67</u> (A) <u>256</u> (B)
4				Prevalence Index = B/A = 3.82
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
		Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover:		
Herb Stratum (Plot size: 5 ft r )  1. Rubus argutus	2		FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2				Definitions of Four Vegetation Strata:
3				Tara Mandanta avaludin naina 0 in (7.0 am)
4.				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
5.				height.
6.				Conline/Chrub Woody plants avaluding vines less
7				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10				<b>Woody vine</b> – All woody vines greater than 3.28 ft in
11				height.
12	2% -	Total Cov		
50% of total cover: 1				
	20% 01	total cover		
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
	=	Total Cov	er	Vegetation Present?  Yes No
50% of total cover:	20% of	total cover		riesent: res No
Remarks: (If observed, list morphological adaptations bel	ow).			
No indicators are met.				
No maleators are met.				

Sampling Point: W-PFO-01\_UPL

SOIL Sampling Point: W-PFO-01\_UPL

Profile Desc	ription: (Describe	to the dep	th needed to docun	nent the	indicator	or confirm	n the absence	of indicators.)
Depth	Matrix	0/		x Feature			<b>-</b> .	B .
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0-8	10YR 3/3	100					Silty Clay Loam	Fill present
8 - 14	10YR 3/2	35			_		Silty Clay Loam	mixed matrix. fill present
8 - 14	10YR 6/1	35						
8 - 14	10YR 6/6	30						
14 - 18	10YR 6/1	60	10YR 6/6	30	<u>C</u>	M	Silty Clay Loam	fill present
14 - 18			10YR 3/2	10	<u>C</u>	М		
			Reduced Matrix, MS			ains.		PL=Pore Lining, M=Matrix.  for Problematic Hydric Soils <sup>3</sup> :
Black Hi Hydroge Stratified Organic 5 cm Mu Muck Pr 1 cm Mu Depleted Thick Da	pipedon (A2)	RR P, T, U) J) ce (A11)	Redox Depre Marl (F10) (L Depleted Och Iron-Mangane	rface (SS y Minera of Matrix (F3) Surface (F Surface (F Surface (F R U) Arric (F11 esse Mas	9) (LRR S, 1 (F1) (LRF (F2) (F6) e (F7) F8) ) (MLRA 1 ses (F12) (	T, U) ₹ O) 51) LRR O, P,	2 cm M Reductor Piedma Anoma (MLF) Red Piedma Very S Other of T)	Muck (A9) (LRR O) Muck (A10) (LRR S) Muck (A10) (LRR P, S, T) Muck (A10) (LRR S) Muck (A10)
_	lucky Mineral (S1) <b>(</b>		Delta Ochric			, 0)		ess disturbed or problematic.
=	Gleyed Matrix (S4)		Reduced Ver	tic (F18)	(MLRA 15		1	·
_	Redox (S5)		Piedmont Flo					4500)
_	Matrix (S6) rface (S7) <b>(LRR P, S</b>	S. T. U)		right Loa	amy Solis (	F20) (MLR	RA 149A, 153C	, 153D)
	_ayer (if observed)							
Type:								
Depth (inc	ches):						Hydric Soil	Present? Yes No
Remarks:							l	
No indic	ators are me	et.						

Project/Site: Golden Triangle Phase 2 City	//County: Lowndes County Sampling Date: 20	021-03-09
Applicant/Owner: TVA	State: Mississippi Sampling Point: W	
••	ction, Township, Range: S26 T18N R16E	
	al relief (concave, convex, none): Concave Slope	(%): 1
Subregion (LRR or MLRA): P 135A Lat: 33.39319		
Soil Map Unit Name: Catalpa silty clay	NWI classification: PUBHh	
Are climatic / hydrologic conditions on the site typical for this time of year?		
Are Vegetation, Soil, or Hydrology significantly dist		No
Are Vegetation, Soil, or Hydrology naturally proble		
SUMMARY OF FINDINGS – Attach site map showing sa		itures, etc.
Hydrophytic Vegetation Present? Yes No		
Hydric Soil Present? Yes V No	Is the Sampled Area	
Wetland Hydrology Present? Yes V No No	within a Wetland? Yes No	
Remarks:		
Sample plot in W-PFO-02. The APT showed r	ormal conditions at the time leading up	to and
during the field survey.	ar are amine reading up	
during the held survey.		
HYDROLOGY		
Wetland Hydrology Indicators:	Secondary Indicators (minimum of tw	vo required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)	
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Su	urface (B8)
High Water Table (A2)  Marl Deposits (B15) (L		(= 0)
Saturation (A3) Hydrogen Sulfide Odor		
	s along Living Roots (C3) Dry-Season Water Table (C2)	
Sediment Deposits (B2)	Iron (C4) Crayfish Burrows (C8)	
☐ Drift Deposits (B3) ☐ Recent Iron Reduction	in Tilled Soils (C6) Saturation Visible on Aerial Imag	gery (C9)
Algal Mat or Crust (B4) Thin Muck Surface (C7	() Geomorphic Position (D2)	
☐ Iron Deposits (B5) ☐ Other (Explain in Rema	arks)	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)	
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, L	J)
Field Observations:		
Surface Water Present? Yes No Depth (inches):		
Water Table Present? Yes No Depth (inches): 2		
Saturation Present? Yes No Depth (inches): _0 (includes capillary fringe)	Wetland Hydrology Present? Yes	No
Describe Recorded Data (stream gauge, monitoring well, aerial photos, p	revious inspections), if available:	
Remarks:		
Wetland hydrology indicators are met. Addition	onal indicators included buttressing of the	ree
trunks within the wetland.		

Sampling Point:	W-PFO-02
. •	

Tana Charles (Diet sine) 30 ft r		Dominant		Dominance Test worksheet:	
<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> )  1. Celtis laevigata	% Cover 60	Species? ✓	FACW	Number of Dominant Species That Are OBL_FACW_or FAC: 5	<b>(A)</b>
2. Juniperus virginiana	20	<del></del>	FACU	That Are OBL, FACW, or FAC: 5	(A)
3. Acer negundo	10		FAC	Total Number of Dominant	<b>(D)</b>
	- —			Species Across All Strata: 6	(B)
4				Percent of Dominant Species That Are OBL FACW or FAC 83.3	
5				That Are OBL, FACW, or FAC: 83.3	(A/B)
6				Prevalence Index worksheet:	
8.				Total % Cover of: Multiply by:	_
v		Total Cov	er	OBL species <u>70</u> x 1 = <u>70</u>	
50% of total cover: 45				FACW species <u>80</u> x 2 = <u>160</u>	
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species $25$ $\times 3 = 75$	
1. Celtis laevigata	20	~	FACW	FACU species 20 x 4 = 80	
2. Ulmus americana	10	~	FAC	UPL species $0 \times 5 = 0$	
3. Ligustrum sinense	5		FAC	Column Totals: <u>195</u> (A) <u>385</u>	(B)
4				Prevalence Index = B/A = 1.97	
5				Hydrophytic Vegetation Indicators:	
6				1 - Rapid Test for Hydrophytic Vegetation	
7				☑ 2 - Dominance Test is >50%	
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>	
		Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain	)
_	20% of	total cover:	7		
Herb Stratum (Plot size: 5 ft r )	0.5			<sup>1</sup> Indicators of hydric soil and wetland hydrology mu	ust
1. Plantago cordata	35		OBL	be present, unless disturbed or problematic.	
2. Packera glabella	30		OBL	Definitions of Four Vegetation Strata:	
3. Juncus effusus	5		OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cr	m) or
4				more in diameter at breast height (DBH), regardles height.	ss of
5					
6				Sapling/Shrub – Woody plants, excluding vines, I than 3 in. DBH and greater than 3.28 ft (1 m) tall.	ess
7				than 3 in. DBH and greater than 3.20 it (1 in) tail.	
8				<b>Herb</b> – All herbaceous (non-woody) plants, regard of size, and woody plants less than 3.28 ft tall.	lless
9				of size, and woody plants less than 3.20 it tail.	
10				<b>Woody vine</b> – All woody vines greater than 3.28 ft	t in
12.				height.	
12.	70% =	Total Cov	er		
50% of total cover: 35					
Woody Vine Stratum (Plot size: 30 ft r )					
1					
2.					
3.					
4					
5				Hydrophytic	
	=	= Total Cov	er	Vegetation	
50% of total cover:	20% of	total cover:		Present? Yes No	
Remarks: (If observed, list morphological adaptations believed)	ow).			. L	
Dominance test is met.					
Dominance test is met.					

SOIL Sampling Point: W-PF0-02

	cription: (Describe	to the depth				or confirm	the absence o	f indicators.)
Depth (inches)	Matrix Color (moist)	%	Redox Color (moist)	<u>k Feature</u> %	s Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
0 - 18	10YR 3/1		10YR 3/8	30	C	PL / M	Clay Loam	Kemans
	1011(0)1		10111 0/0			1 = 7 141	Oldy Louin	
-								
						·		
				-		<del></del>		
	oncentration, D=De					rains.		PL=Pore Lining, M=Matrix.
l <u> </u>	Indicators: (Appli	cable to all L			•			or Problematic Hydric Soils <sup>3</sup> :
Histosol	. ,		Polyvalue Be		. , .		. —	ick (A9) <b>(LRR O)</b>
_	oipedon (A2) stic (A3)		Thin Dark Su Loamy Mucky					ıck (A10) <b>(LRR S)</b> d Vertic (F18) <b>(outside MLRA 150A</b>
	en Sulfide (A4)		Loamy Gleye			νο,		nt Floodplain Soils (F19) (LRR P, S,
	d Layers (A5)		Depleted Mat		/			ous Bright Loamy Soils (F20)
	Bodies (A6) (LRR I	P, T, U)	Redox Dark S		6)			A 153B)
	ıcky Mineral (A7) <b>(L</b>		Depleted Dar					ent Material (TF2)
	esence (A8) (LRR		Redox Depre		8)			allow Dark Surface (TF12)
	ick (A9) (LRR P, T)		☐ Marl (F10) <b>(L</b>	•		-4\	U Other (E	xplain in Remarks)
_	d Below Dark Surfa ark Surface (A12)	ce (ATT)	Depleted Och				T) <sup>3</sup> Indica	tors of hydrophytic vegetation and
	rairie Redox (A16) (	MI RA 150A)						nd hydrology must be present,
	lucky Mineral (S1)		Delta Ochric					s disturbed or problematic.
	Gleyed Matrix (S4)	, ,	Reduced Ver					·
☐ Sandy R	Redox (S5)		Piedmont Flo	odplain S	oils (F19	(MLRA 14	19A)	
_	Matrix (S6)		Anomalous B	right Loar	my Soils	(F20) <b>(MLR</b>	A 149A, 153C,	153D)
	rface (S7) (LRR P,							
	Layer (if observed)	):						
Type:								
	ches):						Hydric Soil P	resent? Yes V No
Remarks:								
Hydric s	oil indicator	F6 is me	et.					

Project/Site: Golden Triangle Phase 2 City	//County: Lowndes County Sampling Date: 2021-03-10
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PFO-03
	ction, Township, Range: S34 T18N R16E
	cal relief (concave, convex, none): Concave Slope (%): 2
Subregion (LRR or MLRA): P 135A Lat: 33.3931	
Soil Map Unit Name: Vaiden silty clay, 2 to 5 percent slopes, ero	
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly dis	
Are Vegetation, Soil, or Hydrology naturally proble	matic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing sa	ampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes ✓ No	
Hydrophytic Vegetation Present? Yes V No No	Is the Sampled Area
Wetland Hydrology Present? Yes V No No	within a Wetland? Yes No
Remarks:	
Sample plot in W-PFO-03. The APT showed r	normal conditions at the time leading up to and
	iormal conditions at the time leading up to and
during the field survey	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15) (L	
Saturation (A3) Hydrogen Sulfide Odol	· · · · · · · · · · · · · · · · · · ·
	s along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)	- ` ' ' - ' - ' - ' - ' - ' - ' - ' - '
☐ Drift Deposits (B3) ☐ Recent Iron Reduction	
Algal Mat or Crust (B4)  Thin Muck Surface (C7	
☐ Iron Deposits (B5) ☐ Other (Explain in Rem	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	☐ Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes V No Depth (inches): 5	
Water Table Present? Yes   ✓ No Depth (inches): 0  Saturation Present? Yes ✓ No Depth (inches): 0	
(includes capillary fringe)	
Describe Recorded Data (stream gauge, monitoring well, aerial photos, p	revious inspections), if available:
Remarks:	
Indicators are met.	

20.4		Dominant		Dominance Test worksheet:	
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species	
1. Salix nigra	30		OBL	That Are OBL, FACW, or FAC: 2 (A	١)
2				Total Number of Dominant	
3				Species Across All Strata: 2 (B	3)
4				Percent of Dominant Species	
5				' 100	√B)
6				,	,
7.				Prevalence Index worksheet:	
8.				Total % Cover of: Multiply by:	
o	30% =	Total Cov	or	OBL species <u>40</u> x 1 = <u>40</u>	
50% of total cover: <u>15</u>				FACW species $0   x 2 = 0$	
	20% 01	total cover.		FAC species $0 \times 3 = 0$	
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species $0   x4 = 0$	
1				UPL species 0 x 5 = 0	
2				Column Totals: 40 (A) 40 (	'D'
3				Column rotals (A) (	(Β)
4				Prevalence Index = B/A = 1.0	
5					
6					
7.				2 - Dominance Test is >50%	
8.				✓ 3 - Prevalence Index is ≤3.0 <sup>1</sup>	
<u> </u>		Total Cov	er		
50% of total cover:				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)	
<b>= 4</b> :	20 /0 01	lotal cover.	· ——		
Herb Stratum (Plot size: 5 ft r )  1 Alternanthera philoxeroides	10	~	OBL	<sup>1</sup> Indicators of hydric soil and wetland hydrology mus be present, unless disturbed or problematic.	st
"	- —		ODL	,	
2				Definitions of Four Vegetation Strata:	
3				Tree – Woody plants, excluding vines, 3 in. (7.6 cm)	) or
4				more in diameter at breast height (DBH), regardless	of
5				height.	
6				Sapling/Shrub – Woody plants, excluding vines, les	ss
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.	
8.				Harb All barbaccaus (non woody) planta regardle	
9.				<b>Herb</b> – All herbaceous (non-woody) plants, regardle of size, and woody plants less than 3.28 ft tall.	:55
10.					
				<b>Woody vine</b> – All woody vines greater than 3.28 ft i	n
11				height.	
12	10%				
F		Total Cov			
50% of total cover: 5	20% of	total cover			
Woody Vine Stratum (Plot size: 30 ft r )					
1					
2					
3					
4					
5				Hydrophytic	
		Total Cov		Vegetation	
50% of total cover:				Present? Yes No	
Remarks: (If observed, list morphological adaptations belo					
, , ,	JVV).				
Rapid test is met.					

Sampling Point: W-PFO-03

SOIL Sampling Point: W-PF0-03

Profile Desc	cription: (Describe to the depti	needed to docu	ment the	indicator	or confirm	the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
_							'
·			_	<del></del>		· · · · · · · · · · · · · · · · · · ·	_
			_			<u> </u>	
-							
			_				_
<del></del>				<del></del>			
			_				
<sup>1</sup> Type: C=C	oncentration, D=Depletion, RM=I	Reduced Matrix, M	S=Maske	d Sand Gra	ains.	<sup>2</sup> Location: PL=F	Pore Lining, M=Matrix.
	Indicators: (Applicable to all L						roblematic Hydric Soils <sup>3</sup> :
Histosol	(A1)	Polyvalue Be	elow Surfa	ace (S8) <b>(L</b>	RR S. T. U	J) 🔲 1 cm Muck (	(A9) (LRR O)
1 <del></del>	pipedon (A2)	Thin Dark Su					(A10) (LRR S)
	istic (A3)	Loamy Muck					ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)	Loamy Gleye	-	. , .	. •,		oodplain Soils (F19) (LRR P, S, T)
_	d Layers (A5)	Depleted Ma		()			Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark	. ,	F6)		(MLRA 15	
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da	,	,		,	Material (TF2)
	resence (A8) (LRR U)	Redox Depre		. ,			w Dark Surface (TF12)
	uck (A9) (LRR P, T)	Marl (F10) (L		٠,			ain in Remarks)
_	d Below Dark Surface (A11)	Depleted Oc		(MI RA 1	51)	Other (Expire	an in remarks)
	ark Surface (A12)	Iron-Mangan				T) <sup>3</sup> Indicators	of hydrophytic vegetation and
_	rairie Redox (A16) (MLRA 150A)	=		. , .		•	nydrology must be present,
_	Mucky Mineral (S1) (LRR O, S)	Delta Ochric			, •,		sturbed or problematic.
_	Gleyed Matrix (S4)	Reduced Ve			0A. 150B)		otalized of problematic.
_	Redox (S5)	Piedmont Flo					
	d Matrix (S6)					A 149A, 153C, 153I	מס
	urface (S7) (LRR P, S, T, U)	/ triointalous i	ongni Loc	arry cons (i	20) (IIILIX	A 140A, 1000, 1001	2,
	Layer (if observed):						
Type:							
, , <u> </u>		<del></del>					
	ches):					Hydric Soil Pres	ent? Yes No
Remarks:							
Soil san	nple not taken due to	inundation	. Hydr	ric soils	assur	ned due to p	revalence of
	•		-			•	
nyaropi	nytic vegetation and	wettand ny	arolog	y maic	ators.		

Project/Site: Golden Triangle Phase 2 City/C	County: Lowndes County Sampling Date: 2021-03-08
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PFO-04
• • • • • • • • • • • • • • • • • • • •	on, Township, Range: S26 T18N R16E
	relief (concave, convex, none): Concave Slope (%): 2
Subregion (LRR or MLRA): P 135A Lat: 33.39200	,, , ,
Soil Map Unit Name: Catalpa silty clay	NWI classification: PFO1A
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly disturbed.	
Are Vegetation, Soil, or Hydrology naturally problem	atic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing san	npling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes _ ✓ No	
Hydric Soil Present? Yes V No	Is the Sampled Area
Wetland Hydrology Present? Yes   ✓ No	within a Wetland? Yes No
Remarks:	
Sample plot in W-PFO-04. The APT showed no	ormal conditions at the time leading up to and
	which conditions at the time loading up to the
during the field survey	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1)  Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15) (LR	
Saturation (A3)  Hydrogen Sulfide Odor (	· · · · · · · · · · · · · · · · · · ·
Water Marks (B1)  Oxidized Rhizospheres a	
☐ Sediment Deposits (B2) ☐ Presence of Reduced Inc. ☐ Drift Deposits (B3) ☐ Recent Iron Reduction in	
☐ Drift Deposits (B3) ☐ Recent Iron Reduction ir ☐ Algal Mat or Crust (B4) ☐ Thin Muck Surface (C7)	n Tilled Soils (C6)  Saturation Visible on Aerial Imagery (C9)  Geomorphic Position (D2)
Iron Deposits (B5)  Other (Explain in Remar	
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes _ V No Depth (inches): 1	
Water Table Present? Yes <u>✓</u> No Depth (inches): <u>0</u>	
Saturation Present? Yes _ V No Depth (inches): 0	Wetland Hydrology Present? Yes No
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial photos, pre	evious inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	

<b>EGETATION (Four Strata)</b> – Use scientific na	ames of pl	ants.		Sampling Point: W-PFO-04
70 ft r		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r 1. Salix nigra	20	Species? ✓	OBL	Number of Dominant Species
·· <del>·</del>				That Are OBL, FACW, or FAC: $\underline{2}$ (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6.				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species <u>40</u> x 1 = <u>40</u>
50% (1.1.) 15		= Total Cov		FACW species $0   x 2 = 0$
50% of total cover: <u>15</u>	20% of	total cover	. 0	FAC species $0 \times 3 = 0$
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species $0 \times 4 = 0$
1.				UPL species 0 x 5 = 0
2				Column Totals: 40 (A) 40 (B)
3				(1)
4				Prevalence Index = $B/A = 1.0$
5				Hydrophytic Vegetation Indicators:
6				☑ 1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				✓ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
	:	= Total Cov	er	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Alternanthera philoxeroides	10	<b>~</b>	OBL	be present, unless disturbed or problematic.
2				Definitions of Four Vegetation Strata:
3				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4				more in diameter at breast height (DBH), regardless of
5				height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
٥				of size, and woody plants less than 3.28 ft tall.
10.				Was devices Allowed wines proster than 2.20 ft in
11				Woody vine – All woody vines greater than 3.28 ft in height.
12.				
	10% :	= Total Cov	er	
50% of total cover: 5				
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3.				
4 5				
J		Total Cov		Hydrophytic Vegetation
EON/ of total covers				Present? Yes No
50% of total cover:		total cover		
Remarks: (If observed, list morphological adaptations bel Rapid test is met.	ow).			

SOIL Sampling Point: W-PFO-04

Depth (inches)         Matrix         Redox Features           Color (moist)         %         Type¹         Loc²         Texture         Remarks           0 - 20         10YR 3/1         70         10YR 3/8         30         C         PL / M         Clay Loam	
0 - 20 10YR 3/1 70 10YR 3/8 30 C PL / M Clay Loam	
-	
Transport Composition De Doubling DM-Doubled Matrix MC-Marked Cond Coning 21 continue DI - Doublining M-Matrix	
<sup>1</sup> Type: C=Concentration, D=Depletion, RM=Reduced Matrix, MS=Masked Sand Grains.  PL=Pore Lining, M=Matrix.  Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)  Indicators for Problematic Hydric Soils <sup>3</sup> :	
☐ Histosol (A1) ☐ Polyvalue Below Surface (S8) (LRR S, T, U) ☐ 1 cm Muck (A9) (LRR O)	
Histic Epipedon (A2) Thin Dark Surface (S9) (LRR S, T, U) 2 cm Muck (A10) (LRR S)	
Black Histic (A3)  Loamy Mucky Mineral (F1) (LRR O)  Reduced Vertic (F18) (outside MLRA 150	
☐ Hydrogen Sulfide (A4)       ☐ Loamy Gleyed Matrix (F2)       ☐ Piedmont Floodplain Soils (F19) (LRR P,         ☐ Stratified Layers (A5)       ☐ Depleted Matrix (F3)       ☐ Anomalous Bright Loamy Soils (F20)	i, T)
Image: Stratified Layers (A5)       Image: Depleted Matrix (F3)       Image: Anomalous Bright Loamy Soils (F20)         Image: Organic Bodies (A6) (LRR P, T, U)       Image: Redox Dark Surface (F6)       (MLRA 153B)	
5 cm Mucky Mineral (A7) (LRR P, T, U) Depleted Dark Surface (F7) Red Parent Material (TF2)	
Muck Presence (A8) (LRR U) Redox Depressions (F8) Very Shallow Dark Surface (TF12)	
1 cm Muck (A9) (LRR P, T) Marl (F10) (LRR U) Under (Explain in Remarks)	
Depleted Below Dark Surface (A11)  Depleted Ochric (F11) (MLRA 151)  Thick Dark Surface (A12)  Depleted Ochric (F11) (MLRA 151)  Iron-Manganese Masses (F12) (LRR O, P, T)  January 10 (LRR O, P, T)	
Coast Prairie Redox (A16) (MLRA 150A) Umbric Surface (F13) (LRR P, T, U) wetland hydrology must be present,	
Sandy Mucky Mineral (S1) (LRR O, S) Delta Ochric (F17) (MLRA 151) unless disturbed or problematic.	
Sandy Gleyed Matrix (S4)  Reduced Vertic (F18) (MLRA 150A, 150B)  Reduced Vertic (F10) (MLRA 150A)	
Sandy Redox (S5) Piedmont Floodplain Soils (F19) (MLRA 149A) Stripped Matrix (S6) Anomalous Bright Loamy Soils (F20) (MLRA 149A, 153C, 153D)	
Dark Surface (S7) (LRR P, S, T, U)	
Restrictive Layer (if observed):	
Type:	
Depth (inches): No	
Remarks:	
Hydric soil indicator F6 is met.	

Project/Site: Golden Triangle Phase 2	City/County: Lownde	es County	Sampling Date: 2021-03-08			
Applicant/Owner: TVA						
••	.,					
	Local relief (concave, o		Slope (%): 0			
Subregion (LRR or MLRA): P 135A Lat: 33		ong: <u>-88.5958254</u>				
Soil Map Unit Name: Catalpa silty clay			tion: PFO1A			
Are climatic / hydrologic conditions on the site typical for this time of	of waar? Waa					
Are Vegetation, Soil, or Hydrology signification,						
Are Vegetation, Soil, or Hydrology naturally	/ problematic? (If ne	eeded, explain any answe	rs in Remarks.)			
SUMMARY OF FINDINGS - Attach site map show	ing sampling point l	ocations, transects	, important features, etc.			
Hydrophytic Vogototion Propert?						
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampleo					
Wetland Hydrology Present? Yes No _✓		nd? Yes	No			
Remarks:	<u> </u>					
Upland sample plot adjacent to W-PFO-0	)4 The area was	a maintained na	asture area. The			
		•				
APT showed normal conditions at the tin	ie leading up to a	and during the fi	eid survey.			
HYDROLOGY						
Wetland Hydrology Indicators:		Secondary Indica	tors (minimum of two required)			
Primary Indicators (minimum of one is required; check all that app	oly)	Surface Soil	Cracks (B6)			
Surface Water (A1)	(B13)	Sparsely Veg	getated Concave Surface (B8)			
High Water Table (A2)  Harl Deposits (		☐ Drainage Pat				
☐ Saturation (A3) ☐ Hydrogen Sulfi	, ,	<u></u> Moss Trim Li	` '			
	spheres along Living Roots		Water Table (C2)			
	educed Iron (C4)	Crayfish Burr	,			
	eduction in Tilled Soils (C6)		sible on Aerial Imagery (C9)			
☐ Algal Mat or Crust (B4) ☐ Thin Muck Suri ☐ Iron Deposits (B5) ☐ Other (Explain	` '	Geomorphic  Shallow Agui	, ,			
Inundation Visible on Aerial Imagery (B7)	iii Neillaiks)	☐ Shallow Aquitard (D3) ☐ FAC-Neutral Test (D5)				
Water-Stained Leaves (B9)			noss (D8) <b>(LRR T, U)</b>			
Field Observations:		<u> </u>	,,,,,			
Surface Water Present? Yes No Depth (inc	hes):					
Water Table Present? Yes No _ ✓ Depth (inc	hes):					
Saturation Present? Yes No Depth (inc	hes): We	etland Hydrology Presen	t? Yes No			
(includes capillary fringe)  Describe Recorded Data (stream gauge, monitoring well, aerial p	hotos previous inspections	s) if available:				
Describe recorded Buta (stream gauge, monitoring won, dental p	notos, previous inspectione	o, ii availabie.				
Remarks:						
No indicators are met.						

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r	% Cover			Number of Dominant Species
1				That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Descrit of Descinant Conscion
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 50 (A/B)
6				
7				Prevalence Index worksheet:
8.				Total % Cover of: Multiply by:
		= Total Co	/er	OBL species <u>0</u> x 1 = <u>0</u>
50% of total cover:				FACW species <u>0</u> x 2 = <u>0</u>
20 #	20 /0 01	total cover		FAC species <u>70</u> x 3 = <u>210</u>
				FACU species <u>25</u> x 4 = <u>100</u>
1				UPL species $0   x 5 = 0$
2				Column Totals: 95 (A) 310 (B)
3				(1)
4				Prevalence Index = B/A = 3.3
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				☐ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
	=	= Total Co	ver	Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover:	20% of	total cover	:	
Herb Stratum (Plot size: 30 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Schedonorus arundinaceus	65	<b>✓</b>	FAC	be present, unless disturbed or problematic.
2. Schizachyrium scoparium	20	~	FACU	Definitions of Four Vegetation Strata:
3. Allium vineale	5		FACU	
4. Ligustrum sinense	5		FAC	<b>Tree</b> – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
· · · · · · · · · · · · · · · · · · ·				height.
5				
6				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				than 5 m. bbit and greater than 5.20 ft (1 m) tail.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	95% :	= Total Co	ver	
50% of total cover: 48	20% of	total cover	<sub>:</sub> 19	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4				
5				Hydrophytic Vegetation
		= Total Co		Present? Yes No
50% of total cover:		total cover	<u> </u>	
Remarks: (If observed, list morphological adaptations believed)	ow).			
No test is met.				

Sampling Point: W-PFO-04\_UPL

SOIL Sampling Point: W-PFO-04\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	ndicator	or confirn	n the absence of i	indicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	%	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 20	10YR 3/3	100					Clay Loam		
-									
-									
				<del>.</del>					
	oncentration, D=De					ains.		=Pore Lining, M=Matr	
	ndicators: (Appli	cable to all Li						Problematic Hydric	Soils":
Histosol			Polyvalue Be					k (A9) <b>(LRR O)</b>	
	pipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck	-		R O)		Vertic (F18) (outside	
	n Sulfide (A4)		Loamy Gleye		F2)			Floodplain Soils (F19	
	Layers (A5)	···	Depleted Ma		-0)			s Bright Loamy Soils	(F20)
	Bodies (A6) (LRR I		Redox Dark	•	•		(MLRA	,	
_	cky Mineral (A7) (L		Depleted Day					nt Material (TF2) low Dark Surface (TF	10)
	esence (A8) <b>(LRR</b> ( ck (A9) <b>(LRR P, T)</b>		Redox Depre		0)			plain in Remarks)	12)
	Below Dark Surfa		Depleted Ocl		/MIRA1	51)	U Other (LX)	piairi ir Nemarks)	
	ark Surface (A12)	CC (A11)	Iron-Mangan				T) <sup>3</sup> Indicato	rs of hydrophytic vege	tation and
	rairie Redox (A16) (	MLRA 150A)						d hydrology must be p	
	lucky Mineral (S1)		Delta Ochric			, •,		disturbed or problema	
	leyed Matrix (S4)	-, -,	Reduced Ver			OA, 150B)		, , , , , , , , , , , , , , , , , , ,	
	edox (S5)		Piedmont Flo						
	Matrix (S6)						RA 149A, 153C, 15	3D)	
Dark Su	face (S7) (LRR P,	S, T, U)							
Restrictive I	ayer (if observed)	):							
Type:									
Depth (inc	ches):						Hydric Soil Pre	esent? Yes	No 🗸
Remarks:	,						,		
	otoro oro m.	•							
No indic	ators are me	et.							
1									

Project/Site: Golden Triangle Phase 2	ty/County: Lowndes County Sampling Date: 2021-03-11					
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PFO-05					
Investigator(s): J. Brown, O. Haney Section, Township, Range: S33 T18N R16E						
	Landform (hillslope, terrace, etc.): Depression Local relief (concave, convex, none): Concave Slope (%): 2					
Subregion (LRR or MLRA): P 135A Lat: 33.3826						
Soil Map Unit Name: Catalpa silty clay NWI classification: NA						
Are climatic / hydrologic conditions on the site typical for this time of year'						
Are Vegetation, Soil, or Hydrology significantly dis						
Are Vegetation, Soil, or Hydrology naturally proble						
	ampling point locations, transects, important features, etc.					
Hydrophytic Vegetation Present? Yes _ ✓ No						
Hydric Soil Present? Yes V No	Is the Sampled Area					
Wetland Hydrology Present? Yes _ ✓ No	within a Wetland? Yes No					
Remarks:						
Sample plot in W-PFO-05. The APT showed	normal conditions at the time leading up to and					
during the field survey	Tormal contained at the time reading up to and					
during the held survey						
HYDROLOGY						
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)					
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)					
Surface Water (A1)  Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)					
High Water Table (A2)  Marl Deposits (B15)						
Saturation (A3)  Hydrogen Sulfide Odd						
1 <del></del>	es along Living Roots (C3) Dry-Season Water Table (C2)					
Sediment Deposits (B2)  Presence of Reduced						
☐ Drift Deposits (B3) ☐ Recent Iron Reduction						
Algal Mat or Crust (B4) Thin Muck Surface (C	(7) Geomorphic Position (D2)					
☐ Iron Deposits (B5) ☐ Other (Explain in Rem	narks)					
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)					
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)					
Field Observations:						
Surface Water Present? Yes No Depth (inches): _						
Water Table Present? Yes No Depth (inches): _						
Saturation Present? Yes No Depth (inches): _ (includes capillary fringe)	Wetland Hydrology Present? Yes No					
Describe Recorded Data (stream gauge, monitoring well, aerial photos,	previous inspections), if available:					
Remarks:						
Wetland hydrology indicators are met.						

Sampling Point: W	/-P	FΟ	1-05
-------------------	-----	----	------

00.6	Absolute	Dominant	Indicator	Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species
1. Celtis laevigata	55		FACW	That Are OBL, FACW, or FAC: 5 (A)
2				Total Niverban of Dansin and
3.				Total Number of Dominant Species Across All Strata: 6 (B)
4.				(b)
				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 83.3 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 5 x 1 = 5
	<u>55%</u> =	Total Cov	er	
50% of total cover: <u>28</u>	20% of	total cover:	11	FACW species $\frac{60}{10}$ x 2 = $\frac{120}{20}$
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species 10 x 3 = 30
1. Celtis laevigata	5	~	FACW	FACU species <u>5</u>
				UPL species <u>0</u> x 5 = <u>0</u>
2				Column Totals: <u>80</u> (A) <u>175</u> (B)
3				
4				Prevalence Index = $B/A = 2.19$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8				✓ 3 - Prevalence Index is ≤3.0 <sup>1</sup>
		Total Cov	er	
50% of total cover: 3				☐ Problematic Hydrophytic Vegetation¹ (Explain)
·	20% 01	lotal cover.	· <u>·</u>	
Herb Stratum (Plot size: 5 ft r )	E	,	E40	<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1. Cyperus esculentus			FAC	be present, unless disturbed or problematic.
2. Elymus virginicus	5		FAC	Definitions of Four Vegetation Strata:
3. Packera glabella	5		OBL	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4. Trifolium repens	5	<b>✓</b>	FACU	more in diameter at breast height (DBH), regardless of
5				height.
6.				Couling/Charle Weeds about a cook diagraphic
				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				than 6 in. BBit and greater than 6.20 it (1 in) tail.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12.				
	20% =	Total Cov	er	
50% of total cover: 10				
Woody Vine Stratum (Plot size: 30 ft r )	20 70 01	10101 00101		
1				
2				
3				
4				
5				Hydrophytic
		Total Cov	er	Vegetation
50% of total cover:				Present? Yes No
		total cover.	<u> </u>	
Remarks: (If observed, list morphological adaptations below	OW).			
Dominance test is met.				

SOIL Sampling Point: W-PF0-05

Profile Desc	ription: (Describe	to the dept	h needed to docum	nent the	indicator	or confir	m the absence of i	ndicators.)	
Depth	Matrix		Redox Features			1 2			
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u> 	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks	
0 - 18	10YR 3/2	95	10YR 3/6	5	<u>C</u>	M	Clay Loam		
							<del></del>		
-									
							· -		
		<del></del> -			-		<del></del>		
	-					. ———	·		
							<del></del>		
			Reduced Matrix, MS			rains.		=Pore Lining, M=Matrix.	
l <u> </u>		cable to all I	RRs, unless other					Problematic Hydric Soils <sup>3</sup> :	
Histosol			Polyvalue Be				. —	(A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					(A10) (LRR S)	
Black Hi	en Sulfide (A4)		Loamy Muck	-		(0)		Vertic (F18) <b>(outside MLRA 150A,B)</b> Floodplain Soils (F19) <b>(LRR P, S, T)</b>	
	d Layers (A5)		Depleted Mar		(1 2)			s Bright Loamy Soils (F20)	
	Bodies (A6) (LRR I	P, T, U)	Redox Dark		<del>-</del> 6)		(MLRA 1		
	ıcky Mineral (A7) <b>(L</b>		Depleted Dar		,		Red Paren	nt Material (TF2)	
	esence (A8) (LRR		Redox Depre		8)			ow Dark Surface (TF12)	
	ick (A9) (LRR P, T)		Marl (F10) <b>(L</b>	-			U Other (Exp	olain in Remarks)	
_	d Below Dark Surfa	ce (A11)	Depleted Och				T) <sup>3</sup> Indicator	rs of hydrophytic vegetation and	
	ark Surface (A12) rairie Redox (A16) (	ΜΙ ΒΔ 150Δ						d hydrology must be present,	
	lucky Mineral (S1)		Delta Ochric					disturbed or problematic.	
	Gleyed Matrix (S4)	(	Reduced Ver						
_	Redox (S5)		Piedmont Flo						
	Matrix (S6)		Anomalous E	Bright Loa	my Soils	(F20) <b>(ML</b>	RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,								
	Layer (if observed	):							
Type:			<u></u>						
	ches):						Hydric Soil Pre	esent? Yes No	
Remarks:			_						
Hydric s	oil indicator	F6 is m	et.						

Project/Site: Golden Triangle Phase 2	City/County: Lowndes County Sa	ampling Date: 2021-03-11				
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PF0-05_UPL					
• • • • • • • • • • • • • • • • • • • •						
	Local relief (concave, convex, none): None	Slone (%)· 3				
Subregion (LRR or MLRA): P 135A Lat: 33.3		Datum: NAD 83				
Soil Map Unit Name: Catalpa silty clay	NWI classification					
Are climatic / hydrologic conditions on the site typical for this time of y						
Are Vegetation, Soil, or Hydrology significantl						
Are Vegetation, Soil, or Hydrology naturally p						
SUMMARY OF FINDINGS – Attach site map showin	sampling point locations, transects, in	nportant features, etc.				
Hydrophytic Vegetation Present? Yes No						
Hydric Soil Present? Yes No	Is the Sampled Area within a Wetland? Yes	No. V				
Wetland Hydrology Present? Yes No✓	within a Wetland? Yes	NO				
Remarks:						
Upland adjacent to W-PFO-05. The APT sl	owed normal conditions at the ti	ime leading up to				
and during the field survey		• .				
HYDROLOGY						
Wetland Hydrology Indicators:	Secondary Indicators	s (minimum of two required)				
Primary Indicators (minimum of one is required; check all that apply	Surface Soil Cra					
Surface Water (A1)  Aquatic Fauna (B		ated Concave Surface (B8)				
High Water Table (A2)  High Water Table (A2)  Marl Deposits (B1)						
Saturation (A3)  Hydrogen Sulfide						
	eres along Living Roots (C3) Dry-Season Wat	` '				
Sediment Deposits (B2)						
1 <del></del>		le on Aerial Imagery (C9)				
Algal Mat or Crust (B4)	(C7) Geomorphic Pos	sition (D2)				
☐ Iron Deposits (B5) ☐ Other (Explain in	temarks)	d (D3)				
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Tes	FAC-Neutral Test (D5)				
Water-Stained Leaves (B9)	☐ Sphagnum moss	s (D8) <b>(LRR T, U)</b>				
Field Observations:						
Surface Water Present? Yes No Depth (inche						
Water Table Present? Yes No Depth (inche						
Saturation Present? Yes No Depth (inche (includes capillary fringe)	): Wetland Hydrology Present?	Yes No				
Describe Recorded Data (stream gauge, monitoring well, aerial pho	os, previous inspections), if available:					
Remarks:						
No indicators are met.						

<b>VEGETATION</b>	(Four Strata)	- Use scientific na	mes of plants.

<b>EGETATION (Four Strata)</b> – Use scientific na	mes of pl	ants.		Sampling Point: W-PFO-05_UPL
20.65		Dominant		Dominance Test worksheet:
<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> ) 1. <u>Celtis laevigata</u>	% Cover 35	Species? ✓	<u>Status</u> FACW	Number of Dominant Species That Are OBL, FACW, or FAC: 3 (A)
2				Total Number of Dominant
3				Species Across All Strata: 4 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 75 (A/B)
6				
7				Prevalence Index worksheet:
8				Total % Cover of: Multiply by:
	35% =	Total Cov	er	OBL species $0 \times 1 = 0$
50% of total cover: <u>18</u>	20% of	total cover:	7	FACW species $\frac{35}{70}$ $\times 2 = \frac{70}{210}$
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species $\frac{70}{35}$ $\times 3 = \frac{210}{100}$
1. Juniperus virginiana	15		FACU	FACU species $\frac{25}{2}$ $\times 4 = \frac{100}{2}$
2				UPL species $0 \times 5 = 0$
3				Column Totals: <u>130</u> (A) <u>380</u> (B)
4.				Prevalence Index = B/A = 2.92
5				Hydrophytic Vegetation Indicators:
6.				1 - Rapid Test for Hydrophytic Vegetation
7				2 - Dominance Test is >50%
8.				2 - Dominance rest is >50 %  ✓ 3 - Prevalence Index is ≤3.0¹
· ·	15% =	Total Cov	er	
50% of total cover: 8				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )			·	The disease of brodeic and another discount and another discount
1 Carex sp.*	45	~	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Ligustrum sinense	20		FAC	Definitions of Four Vegetation Strata:
3. Allium canadense	10		FACU	Dominions of Four Vogotation Gradua
4 Smilax rotundifolia	5		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
5.				height.
			-	
6				Sapling/Shrub – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.
7				
8	- ——			Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
9			-	of size, and woody plants less than 3.20 it tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11			-	height.
12				
10		Total Cov		
50% of total cover: 40	20% of	total cover:	10	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
		Total Cov		Vegetation Present? Yes No
50% of total cover:		total cover:	:	100 <u>100</u>
Remarks: (If observed, list morphological adaptations belo		nt ha id	ontifica	d to species It was assumed to be

FAC since most species in the area are FAC or wetter.

SOIL Sampling Point: W-PFO-05\_UPL

Profile Description: (Describe to the depth	needed to docur	nent the indi	cator or confir	m the absence of ir	ndicators.)
Depth Matrix		x Features	1 . 2		
(inches) Color (moist) %	Color (moist)	<u>% T</u>	ype <sup>1</sup> Loc <sup>2</sup>	<u>Texture</u>	Remarks
<u>0 - 16</u> <u>10YR 3/2</u> <u>100</u> _				Silty Clay	
				<u> </u>	
-					
			; <u></u>	·	
		<del></del>			
		<del></del>		<del>-</del>	
				<del></del>	
				<u> </u>	
<sup>1</sup> Type: C=Concentration, D=Depletion, RM=F	Reduced Matrix, MS	S=Masked Sa	nd Grains.	<sup>2</sup> Location: PL=	Pore Lining, M=Matrix.
Hydric Soil Indicators: (Applicable to all L	RRs, unless other	rwise noted.)		Indicators for	Problematic Hydric Soils <sup>3</sup> :
☐ Histosol (A1)	Polyvalue Be	elow Surface (	S8) <b>(LRR S, T,</b>	U) 1 cm Muck	(A9) (LRR O)
Histic Epipedon (A2)		ırface (S9) <b>(LF</b>			(A10) <b>(LRR S)</b>
Black Histic (A3)	_	y Mineral (F1)	(LRR O)		ertic (F18) (outside MLRA 150A,B)
Hydrogen Sulfide (A4)	=	ed Matrix (F2)			Floodplain Soils (F19) (LRR P, S, T)
Stratified Layers (A5) Organic Bodies (A6) (LRR P, T, U)	Depleted Ma Redox Dark			Anomalous	Bright Loamy Soils (F20)
5 cm Mucky Mineral (A7) (LRR P, T, U)	=	rk Surface (F7	7)	,	t Material (TF2)
Muck Presence (A8) (LRR U)	Redox Depre		,		ow Dark Surface (TF12)
1 cm Muck (A9) (LRR P, T)	Marl (F10) <b>(L</b>				lain in Remarks)
Depleted Below Dark Surface (A11)	_	hric (F11) <b>(ML</b>			
Thick Dark Surface (A12)			F12) <b>(LRR O, P</b>		s of hydrophytic vegetation and
Coast Prairie Redox (A16) (MLRA 150A)		ace (F13) <b>(LRI</b>			hydrology must be present,
Sandy Mucky Mineral (S1) (LRR O, S) Sandy Gleyed Matrix (S4)		(F17) <b>(MLRA</b>			disturbed or problematic.
Sandy Gleyed Matrix (34)			R <b>A 150A, 150B</b> (F19) <b>(MLRA 1</b>		
Stripped Matrix (S6)				<del>-137.)</del> RA 149A, 153C, 153	BD)
Dark Surface (S7) (LRR P, S, T, U)	_	5 , .		, , , , , ,	,
Restrictive Layer (if observed):					
Type:					
Depth (inches):				Hydric Soil Pres	sent? Yes No 🕑
Remarks:					
No indicators are met.					
1					

Project/Site: Golden Triangle Phase 2 City.	County: Lowndes County Sampling Date: 2021-03-10				
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PFO-3_UPL				
nvestigator(s): J. Brown, O. Haney Section, Township, Range: S34 T18N R16E					
Landform (hillslope, terrace, etc.): Hillslope Local relief (concave, convex, none): Convex Slope (%): 15					
Subregion (LRR or MLRA): P 135A Lat: 33.38594	,				
Soil Map Unit Name: Vaiden silty clay, 2 to 5 percent slopes, eroded  NWI classification:  NA					
Are climatic / hydrologic conditions on the site typical for this time of year?					
Are Vegetation, Soil, or Hydrology significantly distr					
Are Vegetation, Soil, or Hydrology naturally probler	natic? (If needed, explain any answers in Remarks.)				
SUMMARY OF FINDINGS - Attach site map showing sa	mpling point locations, transects, important features, etc.				
Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?  Remarks:  Upland sample plot adjacent to W-PFO-03. The sample plot adjacent to W-PFO-03.	Is the Sampled Area within a Wetland?  Yes No  No  No				
leading up to and during the field survey	ie Ai i snowed normal conditions at the time				
leading up to and during the field survey					
LIVEROL OCY					
HYDROLOGY Westernd Hydrology Indicators	Secondary Indicators (minimum of two required)				
Wetland Hydrology Indicators:  Primary Indicators (minimum of one is required; check all that apply)	Secondary mulcators (minimum or two required)  Surface Soil Cracks (B6)				
Surface Water (A1)  Surface Water (A1)  Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)				
High Water Table (A2)  Addatic Faulia (B13)  Marl Deposits (B15) (L1					
Saturation (A3)  Hydrogen Sulfide Odor					
1 <del></del>	along Living Roots (C3) Dry-Season Water Table (C2)				
Sediment Deposits (B2)  Presence of Reduced II					
Drift Deposits (B3)	- ` ' ' - ' - ' - ' - ' - ' - ' - ' - '				
Algal Mat or Crust (B4) Thin Muck Surface (C7	Geomorphic Position (D2)				
☐ Iron Deposits (B5) ☐ Other (Explain in Rema	rks) Shallow Aquitard (D3)				
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)				
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)				
Field Observations:					
Surface Water Present? Yes No Depth (inches):					
Water Table Present? Yes No _ ✓ Depth (inches):					
Saturation Present? Yes No Depth (inches): (includes capillary fringe)	Wetland Hydrology Present? Yes No _✓				
Describe Recorded Data (stream gauge, monitoring well, aerial photos, p	revious inspections), if available:				
Remarks:					
No indicators are met.					

<b>VEGETATION</b>	(Four Strata)	– Use	scientific	names of	of plants.

Tree Stratum (Plot size: 30 ft r )

2. Maclura pomifera

3. Cornus sp. \_\_\_\_\_

Sapling/Shrub Stratum (Plot size: 15 ft r )

Herb Stratum (Plot size: 5 ft r )

1 Carex sp.\*

1. Juniperus virginiana

San	npling Point: W-PF	O-3_UPL						
Dominance Test worksheet:								
Number of Dominant Species That Are OBL, FACW, or FAC	2	(A)						
Total Number of Dominant Species Across All Strata:	3	(B)						
Percent of Dominant Species That Are OBL, FACW, or FAC	66.7	(A/B)						
Prevalence Index worksheet								
Total % Cover of:	Multiply by:	_						
OBL species 0	x 1 = 0	_						
FACW species 0	x 2 = 0	_						
FAC species 4	x 3 = 12	_						
FACU species 70	x 4 = 280	_						
UPL species 0	x 5 = 0	_						
	(A) <u>292</u>	_ (B)						
	= 3.95							
Hydrophytic Vegetation India								
1 - Rapid Test for Hydroph	, ,							
2 - Dominance Test is >50								
☐ 3 - Prevalence Index is ≤3 ☐ Problematic Hydrophytic \		in)						
be present, unless disturbed o	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.							
Definitions of Four Vegetation	n Strata:							
Tree – Woody plants, excludin more in diameter at breast heigheight.								
Sapling/Shrub – Woody plant than 3 in. DBH and greater tha								
<b>Herb</b> – All herbaceous (non-woof size, and woody plants less		rdless						
Woody vine – All woody vines height.	greater than 3.28	3 ft in						

Remarks: (If observed, list morphological adaptations below).

Woody Vine Stratum (Plot size: 30 ft r)

Dominance test is met. \*The Carex could not be identified to species. It was assumed to be FAC since most species in the area are FAC or wetter.

\_\_\_\_\_= Total Cover

4% = Total Cover

50% of total cover: 2 20% of total cover: 0.8

50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_

Absolute Dominant Indicator

% Cover Species? Status

5 \_\_\_\_ \_\_\_

75% = Total Cover

50% of total cover: 38 20% of total cover: 15

50% of total cover: \_\_\_\_\_ 20% of total cover: \_\_\_\_

2 Ligustrum sinense 2 FAC

Yes \_\_\_\_ No \_\_\_\_

Hydrophytic

Vegetation

Present?

SOIL Sampling Point: W-PFO-3\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Cd	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> Location: PL:	=Pore Lining, M=Matr	riy
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RRSTI		k (A9) (LRR O)	
	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-		-,		Floodplain Soils (F19	
	Layers (A5)		Depleted Ma		,			s Bright Loamy Soils	
Organic	Bodies (A6) (LRR I	P, T, U)	Redox Dark	Surface (F	<del>-</del> 6)		(MLRA 1	153B)	
5 cm Mu	cky Mineral (A7) (L	.RR P, T, U)	Depleted Dar	rk Surface	e (F7)		Red Paren	nt Material (TF2)	
	esence (A8) (LRR		Redox Depre	essions (F	8)		<u></u> Very Shall	ow Dark Surface (TF	12)
	ck (A9) (LRR P, T)						U Other (Exp	olain in Remarks)	
	Below Dark Surfa	ce (A11)	Depleted Ocl				3		
	ark Surface (A12)		Iron-Mangan					rs of hydrophytic vege	
	rairie Redox (A16) (					, U)		d hydrology must be p	
	lucky Mineral (S1)	(LKK 0, 5)	Delta Ochric			OA 150D)		disturbed or problema	atic.
	edox (S5)		Reduced Ver						
	Matrix (S6)						+9A) RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,	S. T. U)	Anomalous L	night Loai	illy Solis (	1 20) (IVILIV	A 149A, 133C, 13	30)	
	_ayer (if observed								
Type:		•							
, , <u> </u>	ches):						Hydric Soil Pre	esent? Yes	No 🗸
Remarks:							Tryuno com r re		
No indic	ators are m	et.							

Project/Site: Golden Triangle	Phase 2	Citv/C	ounty: Lowndes C	ounty	Sampling Date: 2021-03-11
Applicant/Owner: TVA					Sampling Point: W-PSS-01
Investigator(s): J. Brown, O. H	laney	Section	n, Township, Range:		
Landform (hillslope, terrace, etc.)	· Depression				Slope (%): 5
Subregion (LRR or MLRA): P 13		Lat: 33.382512			Datum: NAD 83
Soil Map Unit Name: Catalpa s		_ Lat			
				NWI classifica	
Are climatic / hydrologic condition					
Are Vegetation, Soil					present? YesNo
Are Vegetation, Soil	, or Hydrology	naturally problema	itic? (If neede	d, explain any answer	rs in Remarks.)
SUMMARY OF FINDINGS	5 – Attach site ma	ap showing sam	pling point loca	tions, transects	, important features, etc.
Hydrophytic Vegetation Present	t? Yes 🗸	No			
Hydric Soil Present?		No	Is the Sampled Are within a Wetland?		No
Wetland Hydrology Present?		No	within a wettand?	1es	NO
Remarks:					
Sample plot in W-PS	SS-01. The AP	T showed noi	mal condition	ns at the time	leading up to and
during the field surv					
	cy.				
HYDROLOGY					
Wetland Hydrology Indicators				Secondary Indica	ators (minimum of two required)
Primary Indicators (minimum of		all that annly)		Surface Soil	
Surface Water (A1)		atic Fauna (B13)			getated Concave Surface (B8)
High Water Table (A2)		Deposits (B15) <b>(LRF</b>	R U)	Drainage Pat	
Saturation (A3)		rogen Sulfide Odor (0		Moss Trim Li	
Water Marks (B1)		-	long Living Roots (C3		Water Table (C2)
Sediment Deposits (B2)		ence of Reduced Iro		Crayfish Burr	
Drift Deposits (B3)		ent Iron Reduction in			sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)		Muck Surface (C7)		Geomorphic	=
Iron Deposits (B5)	Othe	er (Explain in Remark	s)	Shallow Aqui	tard (D3)
Inundation Visible on Aerial	I Imagery (B7)			FAC-Neutral	Test (D5)
☐ Water-Stained Leaves (B9)	)			Sphagnum m	noss (D8) <b>(LRR T, U)</b>
Field Observations:					
	Yes No				
	Yes No				
Saturation Present? (includes capillary fringe)	Yes No	Depth (inches): 0	Wetlan	d Hydrology Presen	nt? Yes No
Describe Recorded Data (stream	m gauge, monitoring we	ell, aerial photos, pre	vious inspections), if	available:	
Remarks:					
Wetland hydrology i	indicators are i	met.			

Sampling Point:	W-PSS-01

7 01 1 (D) 1 20 ft r		Dominant		Dominance Test worksheet:
<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> ) 1.		Species?		Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
4.         5.				Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species <u>5</u> x 1 = <u>5</u>
		Total Cov		FACW species 75 x 2 = 150
50% of total cover:	20% of	total cover		FAC species 0 x 3 = 0
Sapling/Shrub Stratum (Plot size: 15 ft r )	C.E.		FACW	FACU species 0 x 4 = 0
1. Fraxinus pennsylvanica			OBL	UPL species $0 \times 5 = 0$
2. Salix nigra 3.				Column Totals: 80 (A) 155 (B)
4.				Prevalence Index = B/A = 1.94
5				Hydrophytic Vegetation Indicators:
6				
7				
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>
		Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover: 35	20% of	total cover	14	
Herb Stratum (Plot size: 5 ft r )  1. Andropogon glomeratus	10	<b>~</b>	FACW	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2.				Definitions of Four Vegetation Strata:
3.				Tree Message related a suple discouring a Q in (7.0 and) a
4.				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) o more in diameter at breast height (DBH), regardless o
5.				height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7.				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8.				Harle All back account (a an area de ) relate an areadle an
9.				Herb – All herbaceous (non-woody) plants, regardless of size, and woody plants less than 3.28 ft tall.
10.				
11.				Woody vine – All woody vines greater than 3.28 ft in
12.				height.
12.	10% =	Total Cov	er	
50% of total cover: 5	20% of	total cover	2	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2.				
3.				
4.				
5.				Hydronhytio
		Total Cov	er	Hydrophytic Vegetation
50% of total cover:				Present? Yes No
Remarks: (If observed, list morphological adaptations bel		10101 00101	·	
	iow).			
Rapid test is met.				

SOIL Sampling Point: W-PSS-01

Profile Desc	cription: (Describe to the depti	needed to docur	ment the	indicator of	or confirm	n the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
	. <u> </u>						
_							
·			_				_
-							
			_				_
<del></del>	·						
<sup>1</sup> Type: C=C	oncentration, D=Depletion, RM=I	Reduced Matrix. M	S=Maske	d Sand Gra	ains.	<sup>2</sup> Location: PL=F	Pore Lining, M=Matrix.
	Indicators: (Applicable to all L						Problematic Hydric Soils <sup>3</sup> :
Histosol		Polyvalue Be			RRSTI		•
1 <del></del>	pipedon (A2)	Thin Dark Su					(A10) <b>(LRR S)</b>
I <b>=</b>	istic (A3)	Loamy Muck					ertic (F18) (outside MLRA 150A,B)
_	en Sulfide (A4)	Loamy Gleye	-		0,		loodplain Soils (F19) (LRR P, S, T)
_	d Layers (A5)	Depleted Ma		(1 2)			Bright Loamy Soils (F20)
	Bodies (A6) <b>(LRR P, T, U)</b>	Redox Dark		E6)		(MLRA 15	
	ucky Mineral (A7) (LRR P, T, U)	Depleted Da	,	,		,	Material (TF2)
	resence (A8) (LRR U)	Redox Depre					w Dark Surface (TF12)
	uck (A9) (LRR P, T)	Marl (F10) (L		J)			ain in Remarks)
_	d Below Dark Surface (A11)	Depleted Oc		\	51)	U Other (Expir	all III Nelliaiks)
	ark Surface (A12)	Iron-Mangan				T) <sup>3</sup> Indicators	of hydrophytic vegetation and
	rairie Redox (A16) <b>(MLRA 150A</b> )	_				•	hydrology must be present,
_	Mucky Mineral (S1) (LRR O, S)	Delta Ochric			, 0,		isturbed or problematic.
_	Gleyed Matrix (S4)	Reduced Ve			0A 150B)		isturbed or problematic.
_	Redox (S5)	Piedmont Flo					
	Matrix (S6)					RA 149A, 153C, 153I	D)
	rface (S7) (LRR P, S, T, U)	Anomaious I	ongni Lua	airiy Solis (i	20) (WILK	A 149A, 133C, 133	<i>D</i> )
	Layer (if observed):					<u> </u>	
_	Layer (ii observeu).						
Type:							
Depth (in	ches):					Hydric Soil Pres	ent? Yes No
Remarks:							
Soil san	nple was not taken d	ue to inunda	ation.	Hvdrid	e soil i	s assumed d	lue to prevalence of
	•			•			
nyaropi	nytic vegetation and	wetland ny	arolog	Jy.			
I							

Project/Site: Golden Triangle Phase 2	City/County	<sub>y:</sub> Lowndes County		Sampling Date: 2021-03-11
Applicant/Owner: TVA				Sampling Point: W-PSS-01_UPL
• •	Section, To	ownship, Range: S34 T		· • ———
				Slope (%): 1
Subregion (LRR or MLRA): P 135A Lat		Long:88.6		Datum: NAD 83
Soil Map Unit Name: Catalpa silty clay		_	NWI classificati	
Are climatic / hydrologic conditions on the site typical for this t	ima af waar2 .Vaa			
Are Vegetation, Soil, or Hydrology sig				
Are Vegetation, Soil, or Hydrology nat			-	
SUMMARY OF FINDINGS – Attach site map sl	nowing samplin	ng point locations,	transects,	important features, etc.
Hydrophytic Vegetation Present? Yes No	✓ lo th	ne Sampled Area		
Hydric Soil Present? Yes No	✓ with	nin a Wetland?	Vos	No
Wetland Hydrology Present? Yes No	<u> </u>	iii a wellana:	103	
Remarks:				
Upland sample plot adjacent to W-PSS-0	1. The sample	e plot was in a ma	aintained/	mowed area. The
APT showed normal conditions at the tim	e leading up t	to and during the	e field surv	vey.
HYDROLOGY				
Wetland Hydrology Indicators:		Sec	ondary Indicate	ors (minimum of two required)
Primary Indicators (minimum of one is required; check all that	at apply)	_	Surface Soil C	
	auna (B13)			etated Concave Surface (B8)
	osits (B15) <b>(LRR U)</b>		Drainage Patt	
	Sulfide Odor (C1)		Moss Trim Lin	
	Rhizospheres along L			Vater Table (C2)
	of Reduced Iron (C4)		Crayfish Burro	
☐ Drift Deposits (B3) ☐ Recent Iro	n Reduction in Tilled	d Soils (C6)	Saturation Vis	sible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	Surface (C7)		Geomorphic F	osition (D2)
☐ Iron Deposits (B5) ☐ Other (Ex	plain in Remarks)		Shallow Aquita	ard (D3)
Inundation Visible on Aerial Imagery (B7)			FAC-Neutral 7	Гest (D5)
☐ Water-Stained Leaves (B9)			Sphagnum mo	oss (D8) (LRR T, U)
Field Observations:				
Surface Water Present? Yes No Depti				
Water Table Present? Yes No Depti				
Saturation Present? Yes No Depti (includes capillary fringe)	ı (inches):	Wetland Hydro	ology Present	? Yes No
Describe Recorded Data (stream gauge, monitoring well, ae	rial photos, previous	inspections), if available	e:	
Remarks:				
No indicators are met.				

<b>/EGETATION (Four Strata)</b> – Use scientific na	mes of pl	ants.		Sampling Point: W-PSS-01_UPL
		Dominant		Dominance Test worksheet:
<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> ) 1.	% Cover			Number of Dominant Species That Are OBL, FACW, or FAC: 1 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 50 (A/B)
6				Prevalence Index worksheet:
7				Total % Cover of: Multiply by:
8				OBL species 0 x 1 = 0
				FACW species 0 x 2 = 0
50% of total cover:	20% of	total cover	·	FAC species 55 x 3 = 165
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 80 x 4 = 320
1				UPL species $0$ $x = 0$
2				Column Totals: 135 (A) 485 (B)
3				
4				Prevalence Index = $B/A = 3.59$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				☐ 2 - Dominance Test is >50%
o		Total Cov	/er	3 - Prevalence Index is ≤3.0 <sup>1</sup>
50% of total cover:				Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
Herb Stratum (Plot size: 5 ft r )			·	Indicators of hydric coil and watered hydrology movet
1. Cynodon dactylon	60	~	FACU	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2. Setaria pumila	30	~	FAC	Definitions of Four Vegetation Strata:
3. Cyperus esculentus	25		FAC	_
4. Erigeron canadensis	20		FACU	<b>Tree</b> – Woody plants, excluding vines, 3 in. (7.6 cm) or more in diameter at breast height (DBH), regardless of
5.				height.
6.				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
		Total Cov		
50% of total cover: 68	20% of	total cover	<u> 27 </u>	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
		Total Cov		Vegetation Present? Yes No
50% of total cover:		total cover	:	
Remarks: (If observed, list morphological adaptations below	OW).			
No test is met.				

SOIL Sampling Point: W-PSS-01\_UPL

Profile Desc	ription: (Describe	to the depth	needed to docur	nent the i	indicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Feature					
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks	
0 - 18	10YR 3/2	100					Clay Loam		
-									
	-								
·									
	-								
-									
¹Type: C=Cd	oncentration, D=De	nletion RM=R	educed Matrix MS	S=Masker	Sand Gr	ains	<sup>2</sup> Location: PL:	=Pore Lining, M=Matr	riy
	Indicators: (Appli					anio.		Problematic Hydric	
☐ Histosol			Polyvalue Be			RRSTI		k (A9) (LRR O)	
_	oipedon (A2)		Thin Dark Su					k (A10) <b>(LRR S)</b>	
Black Hi			Loamy Muck					Vertic (F18) (outside	MLRA 150A,B)
	n Sulfide (A4)		Loamy Gleye	-		-,		Floodplain Soils (F19	
	Layers (A5)		Depleted Ma		,			s Bright Loamy Soils	
Organic	Bodies (A6) (LRR I	P, T, U)	Redox Dark	Surface (F	<del>-</del> 6)		(MLRA 1	153B)	
5 cm Mu	cky Mineral (A7) (L	.RR P, T, U)	Depleted Dar	rk Surface	e (F7)		Red Paren	nt Material (TF2)	
	esence (A8) (LRR		Redox Depre	essions (F	8)		<u></u> Very Shall	ow Dark Surface (TF	12)
	ck (A9) (LRR P, T)						U Other (Exp	olain in Remarks)	
	Below Dark Surfa	ce (A11)	Depleted Ocl				3		
	ark Surface (A12)		Iron-Mangan					rs of hydrophytic vege	
	rairie Redox (A16) (					, U)		d hydrology must be p	
	lucky Mineral (S1)	(LKK 0, 5)	Delta Ochric			OA 150D)		disturbed or problema	atic.
	edox (S5)		Reduced Ver						
	Matrix (S6)						+9A) RA 149A, 153C, 15	3D)	
	rface (S7) (LRR P,	S. T. U)	Anomalous L	night Loai	illy Solis (	1 20) (IVILIV	A 149A, 133C, 13	30)	
	_ayer (if observed								
Type:		•							
, , <u> </u>	ches):						Hydric Soil Pre	esent? Yes	No 🗸
Remarks:							Tryuno com rite		
No indic	ators are m	et.							

Project/Site: Golden Triangle Phase 2 City	County: Lowndes County Sampling Date: 2021-03-12
Applicant/Owner: TVA	State: Mississippi Sampling Point: W-PSS-02
• • • • • • • • • • • • • • • • • • • •	tion, Township, Range: S33 T18N R16E
	al relief (concave, convex, none): Concave Slope (%): 1
Subregion (LRR or MLRA): P 135A	, , , , , , , , , , , , , , , , , , , ,
Soil Map Unit Name: Catalpa silty clay	NWI classification: PFO1A
Are climatic / hydrologic conditions on the site typical for this time of year?	
Are Vegetation, Soil, or Hydrology significantly distr	
Are Vegetation, Soil, or Hydrology naturally probler	natic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing sa	mpling point locations, transects, important features, etc.
Lhydrophytic Vegetation Present?	
Hydrophytic Vegetation Present? Yes No  Hydric Soil Present? Yes No	Is the Sampled Area
Wetland Hydrology Present? Yes V No No	within a Wetland? Yes No
Remarks:	-
Sample plot in W-PSS-02. No upland sample plot	was taken. Boundary was determined by significant
	, ,
topographical change. The APT showed normal co	multions at the time leading up to and during the
field survey.	
HYDROLOGY	
Wetland Hydrology Indicators:	Secondary Indicators (minimum of two required)
Primary Indicators (minimum of one is required; check all that apply)	Surface Soil Cracks (B6)
Surface Water (A1) Aquatic Fauna (B13)	Sparsely Vegetated Concave Surface (B8)
High Water Table (A2)  Marl Deposits (B15) (LI	RR U) Drainage Patterns (B10)
Saturation (A3) Hydrogen Sulfide Odor	(C1) Moss Trim Lines (B16)
Water Marks (B1) ☐ Oxidized Rhizospheres	along Living Roots (C3) Dry-Season Water Table (C2)
Sediment Deposits (B2)  Presence of Reduced In	ron (C4)
Drift Deposits (B3)	in Tilled Soils (C6) Saturation Visible on Aerial Imagery (C9)
Algal Mat or Crust (B4)	) Geomorphic Position (D2)
Iron Deposits (B5) Other (Explain in Rema	rks) Shallow Aquitard (D3)
Inundation Visible on Aerial Imagery (B7)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	Sphagnum moss (D8) (LRR T, U)
Field Observations:	
Surface Water Present? Yes No Depth (inches): 6	
Water Table Present? Yes No Depth (inches): 0	
Saturation Present? Yes V No Depth (inches): 0 (includes capillary fringe)	Wetland Hydrology Present? Yes No
Describe Recorded Data (stream gauge, monitoring well, aerial photos, pl	revious inspections), if available:
Remarks:	
Wetland hydrology indicators are met.	

20.4		Dominant		Dominance Test worksheet:	
Tree Stratum (Plot size: 30 ft r		Species?		Number of Dominant Species	
1. Fraxinus pennsylvanica	50		FACW	That Are OBL, FACW, or FAC: 2 (A)	
2. Gleditsia aquatica 3.	5		OBL	Total Number of Dominant Species Across All Strata: 2 (B)	
4.					
5				Percent of Dominant Species That Are OBL, FACW, or FAC: 100 (A/B	)
6				Prevalence Index worksheet:	
7				Total % Cover of: Multiply by:	
8				OBL species 5 x 1 = 5	
		Total Cov		FACW species 50	
50% of total cover: 28	20% of	total cover	11		
Sapling/Shrub Stratum (Plot size: 15 ft r )				1 AO 3pcoics X 0 =	
1				FACU species $0 \times 4 = 0$	
2				UPL species $0 \times 5 = 0$	
3				Column Totals: <u>57</u> (A) <u>111</u> (B)	
4				Prevalence Index = B/A = 1.95	
5				Hydrophytic Vegetation Indicators:	
6				1 - Rapid Test for Hydrophytic Vegetation	
7					
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>	
	=			Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)	
50% of total cover:	20% of	total cover	:		
Herb Stratum (Plot size: 5 ft r )  1. Rumex crispus	2	V	FAC	<sup>1</sup> Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.	
2.				Definitions of Four Vegetation Strata:	
3.					
4.				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) o more in diameter at breast height (DBH), regardless of	
				height.	
5					
6				<b>Sapling/Shrub</b> – Woody plants, excluding vines, less than 3 in. DBH and greater than 3.28 ft (1 m) tall.	
7				and the second s	
8				Herb – All herbaceous (non-woody) plants, regardless	i
9				of size, and woody plants less than 3.28 ft tall.	
10				<b>Woody vine</b> – All woody vines greater than 3.28 ft in	
11				height.	
12	20/				
		= Total Cov			
50% of total cover: 1	20% of	total cover			
Woody Vine Stratum (Plot size: 30 ft r )					
1					
2					
3					
4					
5				Hydrophytic	
	=	Total Cov	er	Vegetation	
50% of total cover:	20% of	total cover	·	Present? Yes No	
Remarks: (If observed, list morphological adaptations belo	ow).			. I	
, , , , , , , , , , , , , , , , , , , ,	,				
Dominance test is met.					

Sampling Point: W-PSS-02

SOIL Sampling Point: W-PSS-02

Profile Desc	cription: (Describe to the depti	n needed to docu	ment the	indicator	or confirm	n the absence of in	dicators.)
Depth	Matrix	Redo	x Feature	es			
(inches)	Color (moist) %	Color (moist)	%	Type <sup>1</sup>	Loc <sup>2</sup>	<u>Texture</u>	Remarks
-							
l — —							-
			_				
-							
			-				
l ———						-	
			_				
_							
17	- D.	Darley and Matrice M	0 Ml			21	Daniel Linkson M. Marketo
	oncentration, D=Depletion, RM=I Indicators: (Applicable to all L				ains.		Problematic Hydric Soils <sup>3</sup> :
							•
Histosol		Polyvalue Be					
	pipedon (A2)	Thin Dark Su					(A10) <b>(LRR S)</b>
_	istic (A3)	Loamy Muck	-	. , .	(O)		ertic (F18) (outside MLRA 150A,B)
	en Sulfide (A4)	Loamy Gleye		(F2)			oodplain Soils (F19) (LRR P, S, T)
	d Layers (A5)	Depleted Ma	. ,				Bright Loamy Soils (F20)
	Bodies (A6) (LRR P, T, U)	Redox Dark	•			(MLRA 15	,
_	ucky Mineral (A7) (LRR P, T, U)	Depleted Da					Material (TF2)
	resence (A8) (LRR U)	Redox Depre		=8)			w Dark Surface (TF12)
_	uck (A9) <b>(LRR P, T)</b>	☐ Marl (F10) <b>(L</b>				U Other (Expla	ain in Remarks)
	d Below Dark Surface (A11)	Depleted Oc					
_	ark Surface (A12)	Iron-Mangan		, , ,		•	of hydrophytic vegetation and
	rairie Redox (A16) (MLRA 150A)				, U)		hydrology must be present,
_	Mucky Mineral (S1) (LRR O, S)	Delta Ochric					sturbed or problematic.
_	Gleyed Matrix (S4)	Reduced Ve					
	Redox (S5)	Piedmont Flo					
	d Matrix (S6)	Anomalous I	Bright Loa	amy Soils (	F20) <b>(MLR</b>	RA 149A, 153C, 153I	D)
	ırface (S7) (LRR P, S, T, U)					_	
Restrictive	Layer (if observed):						
Type:							
Depth (in	ches):					Hydric Soil Pres	ent? Yes <u>'</u> No
Remarks:							
No com	nla takan Sail was ir	undatad					
NO Sam	ple taken. Soil was ir	iunuateu.					

Project/Site: Golden Triangle F	Phase 2	City/C	<sub>ounty:</sub> Lowndes	County	Sampling Date: 2021-03-10				
Applicant/Owner: TVA  State: Mississippi Sampling Point: W-PSS/PFO-01									
Investigator(s): J. Brown, O. Haney  Section, Township, Range: S34 T18N R16E									
Landform (hillslope, terrace, etc.):	Depression				Slope (%): 2				
	Subregion (LRR or MLRA): P 135A Lat: 33.383971 Long: -88.603066 Datum: NAD 83								
Soil Map Unit Name: Catalpa sil				NWI classifica					
Are climatic / hydrologic conditions									
Are Vegetation, Soil	, or Hydrology	significantly distur	bed? Are "N	ormal Circumstances" p	resent? YesNo				
Are Vegetation, Soil, or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)									
SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.									
Hydrophytic Vegetation Present?	Yes 🗸	_ No							
Hydric Soil Present?	Yes 🗸	No	Is the Sampled A						
Wetland Hydrology Present?		No	within a Wetland	!? Yes	No				
Remarks:									
Sample plot in W-PS	S/PFO-01 Th	e ΔPT showe	d normal co	anditions at the	time leading up to				
		10 711 1 3110 110	a normal co	mandons at the	time leading up to				
and during the field	survey.								
HYDROLOGY									
Wetland Hydrology Indicators:	;			Secondary Indica	tors (minimum of two required)				
Primary Indicators (minimum of c	one is required; check	( all that apply)		Surface Soil	Cracks (B6)				
Surface Water (A1)		ıatic Fauna (B13)			jetated Concave Surface (B8)				
High Water Table (A2)		1 Deposits (B15) (LRF		☑ Drainage Pat					
Saturation (A3)		Irogen Sulfide Odor (0		Moss Trim Li					
Water Marks (B1)		dized Rhizospheres a			Water Table (C2)				
Sediment Deposits (B2)		sence of Reduced Iro		Crayfish Burr					
Drift Deposits (B3)		cent Iron Reduction in	Tilled Soils (C6)		sible on Aerial Imagery (C9)				
Algal Mat or Crust (B4)		n Muck Surface (C7)		Geomorphic	` '				
Iron Deposits (B5)		er (Explain in Remark	.S)	Shallow Aqui	` '				
Inundation Visible on Aerial Water-Stained Leaves (B9)	imagery (B7)			FAC-Neutral	noss (D8) <b>(LRR T, U)</b>				
Field Observations:				Эрпаупипп	1035 (D0) (LIKIT 1, <b>0)</b>				
	∕es	Depth (inches): 1							
	/es No								
	/es / No		Wetl	land Hydrology Presen	t? Yes No				
(includes capillary fringe)									
Describe Recorded Data (stream	ı gauge, monitoring w	vell, aerial photos, pre	vious inspections),	if available:					
Remarks:									
Wetland hydrology indicators are met.									
Wedana nyarelegy n	idioatoro dre	11100.							

<b>/EGETATION (Four Strata)</b> – Use scientific n	ames of pl	ants.		Sampling Point: W-PSS/PFO-01
00.6		Dominant		Dominance Test worksheet:
Tree Stratum (Plot size: 30 ft r 1. Fraxinus pennsylvanica	% Cover 70	Species? ✓	Status FACW	Number of Dominant Species That Are OBL, FACW, or FAC: 2 (A)
2				Total Number of Dominant
3				Species Across All Strata: 2 (B)
4				Percent of Dominant Species
5				That Are OBL, FACW, or FAC: 100 (A/B)
6				Prevalence Index worksheet:
7				
8				
		= Total Cov		OBL species $0 \times 1 = 0$ FACW species $120 \times 2 = 240$
50% of total cover: 35	20% of	total cover	14	FAC species 0 x 3 = 0
Sapling/Shrub Stratum (Plot size: 15 ft r )				FACU species 0 x 4 = 0
1. Fraxinus pennsylvanica	50		FACW	UPL species 0 x 5 = 0
2				100
3				Column Totals: 120 (A) 240 (B)
4				Prevalence Index = $B/A = 2.0$
5				Hydrophytic Vegetation Indicators:
6				1 - Rapid Test for Hydrophytic Vegetation
7				
8				② 3 - Prevalence Index is ≤3.0 <sup>1</sup>
		= Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)
50% of total cover: 25	20% of	total cover	10	
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology must
1				be present, unless disturbed or problematic.
2				Definitions of Four Vegetation Strata:
3				Tree – Woody plants, excluding vines, 3 in. (7.6 cm) or
4				more in diameter at breast height (DBH), regardless of
5				height.
6				Sapling/Shrub – Woody plants, excluding vines, less
7				than 3 in. DBH and greater than 3.28 ft (1 m) tall.
8				Herb – All herbaceous (non-woody) plants, regardless
9				of size, and woody plants less than 3.28 ft tall.
10				Woody vine – All woody vines greater than 3.28 ft in
11				height.
12				
	=	= Total Cov	er	
50% of total cover:	20% of	total cover	·	
Woody Vine Stratum (Plot size: 30 ft r )				
1				
2				
3				
4				
5				Hydrophytic
	=	= Total Cov	er	Vegetation Present? Yes ✓ No
50% of total cover:	20% of	total cover	·	Present? Yes No
Remarks: (If observed, list morphological adaptations be	elow).			
Rapid test is met.				

SOIL Sampling Point: W-PSS/PFO-01

Profile Desc	ription: (Describe	to the depth	needed to docun	nent the i	ndicator	or confirm	the absence of in	dicators.)
Depth	Matrix			x Feature	S1	. 2		
(inches)	Color (moist)	_ <u>%</u>	Color (moist)	<u>%</u>	Type'	Loc <sup>2</sup>	<u>Texture</u>	Remarks
0 - 20	10YR 2/1	100					Clay Loam	
					-			
-								
_								
				· <del></del>				
				· <del></del>				
-								
	oncentration, D=Dep					ains.		Pore Lining, M=Matrix.
-	ndicators: (Applic	cable to all L	_		•		_	Problematic Hydric Soils <sup>3</sup> :
Histosol (			Polyvalue Be				. —	
Black His	oipedon (A2)		Thin Dark Su Loamy Muck					(A10) <b>(LRR S)</b> ertic (F18) <b>(outside MLRA 150A,B)</b>
I <b>=</b>	n Sulfide (A4)		Loamy Gleye		. , .	. 0,		oodplain Soils (F19) (LRR P, S, T)
	Layers (A5)		Depleted Mat		,			Bright Loamy Soils (F20)
Organic	Bodies (A6) (LRR F	P, T, U)	Redox Dark	Surface (F	6)		(MLRA 15	
	cky Mineral (A7) (L		Depleted Dar					Material (TF2)
_	esence (A8) (LRR l	J)	Redox Depre		8)			w Dark Surface (TF12)
_	ck (A9) <b>(LRR P, T)</b> I Below Dark Surfac	re (Δ11)	Marl (F10) (L Depleted Och		(MIRA 1	51)	Utner (Expla	ain in Remarks)
	ark Surface (A12)	oc (ATT)	Iron-Mangan		•	-	T) <sup>3</sup> Indicators	of hydrophytic vegetation and
_	rairie Redox (A16) (	MLRA 150A)	=				•	hydrology must be present,
Sandy M	lucky Mineral (S1) (	LRR O, S)	Delta Ochric	(F17) <b>(ML</b>	.RA 151)		unless di	sturbed or problematic.
_	leyed Matrix (S4)		Reduced Ver					
	edox (S5)		Piedmont Flo					D)
_	Matrix (S6) face (S7) <b>(LRR P,</b> 3	S T II)	Anomalous B	right Loar	ny Solis (	F20) (WLR	A 149A, 153C, 153I	טן
	ayer (if observed)							
Type:	, (,							
, , <u> </u>	ches):						Hydric Soil Pres	ent? Yes 🗸 No
Remarks:			<del></del>				1.,	
	ne of rodov	are mos	t likely mac	kad di	io to t	ha avtr	omoly dark	and saturated nature
			-				-	
	•			the p	revale	nce of	hydrophytic	vegetation and
wetland	hydrology i	ndicator	s.					
	, ,,							

Project/Site: Golden Triangle	Phase 2	City/C	<sub>ounty:</sub> Lownde	s County	Sampling Date: 2021-03-10				
Applicant/Owner: TVA State: Mississippi Sampling Point: W-PSS/PFO-01_UPL									
Investigator(s): J. Brown, O. Haney Section, Township, Range: S34 T18N R16E									
Landform (hillslope, terrace, etc.					Slope (%): 10				
	Subregion (LRR or MLRA): P 135A Lat: 33.383022 Long: -88.605213 Datum: NAD 83								
Soil Map Unit Name: PFO1A NWI classification: NA									
Are climatic / hydrologic conditio	ns on the site typical for	this time of year? Yo	es No						
Are Vegetation, Soil					present? YesNo				
Are Vegetation, Soil, or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)  SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.									
Hydrophytic Vegetation Preser	nt? Yes 🔽	No							
Hydric Soil Present?	Yes	No V	Is the Sampled		4				
Wetland Hydrology Present?	Yes	No V	within a Wetlar	id? Yes	No				
Remarks:									
Upland sample plot	adjacent to W	-PSS/PFO-01	. The APT	showed normal	conditions at the				
time leading up to a	•								
HYDROLOGY									
Wetland Hydrology Indicator	s:			Secondary Indica	tors (minimum of two required)				
Primary Indicators (minimum o	f one is required; check	all that apply)		Surface Soil	Cracks (B6)				
Surface Water (A1)		atic Fauna (B13)		Sparsely Veç	getated Concave Surface (B8)				
High Water Table (A2)		Deposits (B15) (LRR		☐ Drainage Pat					
Saturation (A3)		rogen Sulfide Odor (C		Moss Trim Li					
Water Marks (B1)		lized Rhizospheres al sence of Reduced Iror			Water Table (C2)				
Sediment Deposits (B2)  Drift Deposits (B3)		ent Iron Reduction in		☐ Crayfish Burr	sible on Aerial Imagery (C9)				
Algal Mat or Crust (B4)		Muck Surface (C7)	Tilled Collo (CO)		Position (D2)				
Iron Deposits (B5)		er (Explain in Remark	s)	Shallow Aquitard (D3)					
Inundation Visible on Aeria				FAC-Neutral Test (D5)					
☐ Water-Stained Leaves (B9	·)			Sphagnum m	noss (D8) (LRR T, U)				
Field Observations:									
Surface Water Present?	Yes No								
Water Table Present?	Yes No								
Saturation Present? (includes capillary fringe)	Yes No	Depth (inches):	We	etland Hydrology Presen	t? Yes No				
Describe Recorded Data (stream	ım gauge, monitoring we	ell, aerial photos, prev	vious inspections	), if available:					
Remarks:									
No indicators are m	et								
	<b>C</b> t.								

Sampling Point:	W-PSS/PFO-01_UP
-----------------	-----------------

<u>Tree Stratum</u> (Plot size: <u>30 ft r</u> )  1. Juniperus virginiana	70 COVE		Status		
1. Sumperus virgimana	40	Species? ✓	FACU	Number of Dominant Species That Are OBL FACW or FAC: 6	<b>^</b> \
2 Maclura pomifera	40	<u> </u>	FACU	That Are OBL, FACW, or FAC: 6	A)
3. Ulmus americana	20		FAC	Total Number of Dominant	
	· ——			Species Across All Strata: 10 (I	B)
4				Percent of Dominant Species	
5					A/B)
6				Prevalence Index worksheet:	
7				Total % Cover of: Multiply by:	
8					
	100% =	= Total Cov	/er		
50% of total cover: 50	20% of	total cover	20	FACW species $0 \times 2 = 0$	
Sapling/Shrub Stratum (Plot size: 15 ft r )				FAC species $\frac{120}{135}$ x 3 = $\frac{360}{540}$	
1. Juniperus virginiana	40		FACU	FACU species 135 x 4 = 540	
2. Ulmus americana	15	~	FAC	UPL species $0 \times 5 = 0$	
3. Cornus florida	10		FACU	Column Totals: 255 (A) 900	(B)
4.				Decomposite description D/A 3.53	
				Prevalence Index = B/A = 3.53	
5				Hydrophytic Vegetation Indicators:	
6				1 - Rapid Test for Hydrophytic Vegetation	
7				2 - Dominance Test is >50%	
8				3 - Prevalence Index is ≤3.0 <sup>1</sup>	
0.0		Total Cov		Problematic Hydrophytic Vegetation <sup>1</sup> (Explain)	
50% of total cover: 33	20% of	total cover	: 13		
Herb Stratum (Plot size: 5 ft r )				<sup>1</sup> Indicators of hydric soil and wetland hydrology mu	st
1. Ligustrum sinense	40		FAC	be present, unless disturbed or problematic.	
2. Carex sp.*	20		FAC	Definitions of Four Vegetation Strata:	
3. Viola sororia	15		FAC	Tree – Woody plants, excluding vines, 3 in. (7.6 cm	a) or
4				more in diameter at breast height (DBH), regardles	
5.				height.	
6.				Sapling/Shrub – Woody plants, excluding vines, le	000
7.					:55
8				Herb – All herbaceous (non-woody) plants, regardled of size, and woody plants less than 3.28 ft tall.	ess
9				of size, and woody plants less than 5.20 it tall.	
10				Woody vine – All woody vines greater than 3.28 ft	in
11				height.	
12					
		Total Cov			
50% of total cover: 38	20% of	total cover	: <u>15</u>		
Woody Vine Stratum (Plot size: 30 ft r )					
1. Smilax rotundifolia	10		FAC		
2. Lonicera japonica	5		FACU		
3.					
4.					
5					
J	15% =	Total Cov	·or	Hydrophytic Vegetation	
500/ 51.1.				Present? Yes No	
50% of total cover: 8		total cover	:		
	w).				
Remarks: (If observed, list morphological adaptations below					

SOIL Sampling Point: W-PSS/PFO-01\_UPL

Profile Desci	iption: (Describe	to the depth	needed to docur	nent the i	ndicator	or confirn	n the absence of i	ndicators.)	
Depth	Matrix			x Features		1 2 2	Tavetura	Damade	_
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remark	<u>s</u>
0 - 18	10YR 2/1	100					Clay Loam		
-									
		<del></del> -							
							<u> </u>		
<sup>1</sup> Type: C=Co	ncentration, D=De	pletion, RM=F	educed Matrix, MS	S=Masked	Sand Gr	ains.	<sup>2</sup> Location: PL	=Pore Lining, M=Ma	atrix.
Hydric Soil II	ndicators: (Appli	cable to all L	RRs, unless other	rwise note	ed.)		Indicators for	Problematic Hydr	ic Soils³:
Histosol (	<b>A1</b> )		Polyvalue Be	elow Surfac	ce (S8) <b>(L</b>	RR S, T, l	<b>ل</b> ) <u> </u>	(A9) <b>(LRR O)</b>	
Histic Ep	pedon (A2)		Thin Dark Su					(A10) <b>(LRR S)</b>	
Black His			Loamy Muck	-		(O)		/ertic (F18) (outsid	
	Sulfide (A4)		Loamy Gleye		F2)			Floodplain Soils (F1	
	Layers (A5)	р т и\	Depleted Mar	. ,	(C)			s Bright Loamy Soil	s (F20)
	Bodies (A6) <b>(LRR</b> ky Mineral (A7) <b>(L</b>		Redox Dark	`	,		(MLRA 1	nt Material (TF2)	
_	sence (A8) <b>(LRR</b>		Redox Depre		, ,			ow Dark Surface (T	F12)
	ck (A9) <b>(LRR P, T)</b>		Marl (F10) (L		3)			plain in Remarks)	,
	Below Dark Surfa		Depleted Ocl		(MLRA 1	51)		,	
Thick Da	k Surface (A12)		Iron-Mangan	ese Masse	es (F12) <b>(</b>	LRR O, P,	T) <sup>3</sup> Indicator	rs of hydrophytic ve	getation and
	airie Redox (A16)					, U)		d hydrology must be	-
_	ucky Mineral (S1)	(LRR O, S)	Delta Ochric					disturbed or probler	matic.
_	eyed Matrix (S4)		Reduced Ver						
Sandy Re			Piedmont Flo					3D)	
_	Matrix (S6) face (S7) <b>(LRR P,</b>	S T II)	Anomaious E	srigni Loar	riy Solis (	-20) (WILK	RA 149A, 153C, 15	30)	
	ayer (if observed								
Type:	, (	,-							
, ,	nes):						Hydric Soil Pre	esent? Yes	No <u> </u>
Remarks:			<u> </u>				Trydric 30ii Fre	.sent: 1es	
No indica	ators are m	et.							
1									



Site: PU	B-001		Rater(s): JB; OH		Date: 03/2021
max 6 pts.	2 subtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Capital	Select one size class and assign  >50 acres (>20.2 ha) (6 pi  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 h  X 0.3 to <3 acres (0.04 to <  0.1 to <0.3 acre (0.04 ha) (0)	s) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around 225 m to <50 m (82 to <164 pe 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double controlled for older forest, prairie, savand), shrubland, young 2nd grosidential, fenced pasture, p	wetland perimeter (7)  ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) heck and average. hah, wildlife area, etc. (7) howth forest (5) hark, conservation tillage, new	(4) er (1)
	13	Metric 3. Hydrolog	ЭУ		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Seleston No.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/Cl 3e. Modifications to natural hydrom None or none apparent (1) Recovered (7)  X Recevering (3)  Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) ct only one and assign scor a.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in.) logic regime. Score one or or	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat Seasonally satura double check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	6	Metric 4. Habitat A	Alteration and D	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (1) Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and averag	Check all disturbances of   ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-001		Rater(s): OH:jb	Date: 03/2021				
21 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Wetlands					
	*If the documented raw score for	*If the documented raw score for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.					
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. t. Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain, Gross morph. adapt. in >5 tre Ecological community with glo Known occurrence state/fedet [*use higher rank where mix Superior/enhanced habitat/us	multiple values apply in row, score row as single feature with highest point value. Provide action (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] re such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5), perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) ac (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) aplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) rith global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] [exclude records which are only "historic"] tat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3): <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
1	Metric 6. Plant Co	mmunities, Interspe	rsion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 Moss/lichen. Other	ale. 0 = Absent or <0.1 ha  [For BR/CM <0.0] 1 = Present and eithe  moderate quality, 2 = Present and eithe is of moderate qu	a (0.25 acre) contiguous acre 4 ha (0.1 acre)] er comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality er comprises a significant part of wetland's vegetation and lality, or comprises a small part and is of high quality prises a significant part or more of wetland's vegetation				
	6b. Horizontal (plan view) intersp Select only one.  High (5)  Moderately high (4) [BR/CM Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CI	iow = Low species dinative species  iom (5)] mod = Native species	Narrative Description of Vegetation Quality  low = Low species diversity &/or dominance of nonnative or disturbance tolerant native species  mod = Native species are dominant component of the vegetation, although nonnative &/or disturbance tolerant native species can also be present, and species diversity moderate to moderately high, but generally				
	Low (1) [BR/CM (2)]  X None (0)	high = A predominand tolerant native	f rare, threatened or endangered species e of native species with nonnative sp &/or disturbance sp absent or virtually absent, and high sp diversity and often the presence of rate, threatened, or endangered species				
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5 Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	e. Mudflat and Open War 0 = Absent <0.1 ha (0.3) 1 = Low 0.1 to <1 ha (0.1 to 0.5 acre)] 2 = Moderate 1 to <4					
	6d. Microtopography.  Score all present using 0 to 3 sci Vegetated hummocks/tusi Coarse woody debris >15 Standing dead >25 cm (10 Amphibian breeding pools	Hypothetical Wetland ale. socks cm (6 in.) 0 in.) dbh	I for Estimating Degree of Interspersion  Low Moderate Moderate High				
		2 = Present in moder amounts of higher	mall amounts or if more common of marginal quality ate amounts, but not of highest quality or in small st quality ate or greater amounts and of highest quality				
22	GRAND 1 (max 100	30- 59 = Category 2	, low wetland function, condition, quality** , good/moderate wetland function, condition, quality** , superior wetland function, condition, quality**				

Site: PUB-002		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (Size) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
max 14 pts. subtotal	MEDIUM. Buffers average	th. Select only one and assi 0 m (164 ft) or more around e 25 m to <50 m (82 to <164	gn score. Do not double check wetland perimeter (7) If t) around wetland perimeter	<. (4)
	VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	average <10 m (<32 ft) aro use. Select one or double co or older forest, prairie, savai s), shrubland, young 2nd gro esidential, fenced pasture, p	heck and average. nnah, wildlife area, etc. (7) bwth forest (5) ark, conservation tillage, new	
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selector > 0.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydrom None or none apparent (1) Recovered (7)  X Recovering (3)  Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) cot only one and assign score) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in ologic regime. Score one or 2)  Check all disturbance	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) auturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
		weir stormwater input	☐ dredging ☐ other cattle grazing	
6	Metric 4. Habitat	Alteration and I	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-002		Rater(s): OH:jb	Rater(s): OH:jb		Date: 03/2021		
21 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	<b>Vetlands</b>					
	*If the documented raw score for	f the documented raw score for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.					
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedel [*use higher rank where mixe Superior/enhanced habitat/us	nere multiple values apply in row, score row as single feature with highest point value. Provide selection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). e (10); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) &/or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] feature such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) lated, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) I acre (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) floodplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) bt. in >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) inity with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] state/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) where mixed rank or qualifier] [exclude records which are only "historic"] habitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) ality) : <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
max 20 pts. subtotal	Metric 6. Plant Co	mmunities, Inters	persion, Mic	rotopography			
max 20 pts. Subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 l	le. 0 = Absent or <	uality, or comprises a signal either comprises a signate quality, or comprises domprises a significant	Juous acre  all part of wetland's vegetation and is gnificant part but is of low quality lificant part of wetland's vegetation a small part and is of high quality part or more of wetland's vegetation	and		
	6b. Horizontal (plan view) intersponded to Select only one.		iption of Vegetation Quies diversity &/or domina	uality ance of nonnative or disturbance tole	erant		
	High (5)  Moderately high (4) [BR/C Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CM Low (1) [BR/CM (2)]  None (0)	M (5)] mod = Native sp mod = Native sp nonnative and speci w/o prese high = A predom tolerant n	ecies ecies are dominant com &/or disturbance tolerar es diversity moderate to ence of rare, threatened e inance of native species ative sp absent or virtual	ponent of the vegetation, although not native species can also be present moderately high, but generally or endangered species with nonnative sp &/or disturbance ly absent, and high sp diversity and	nt,		
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5 Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	Mudflat and Op  0 = Absent <0.0  1 = Low 0.1 to 0.5 a  (0.1 to 0.5 a  2 = Moderate 1	en Water Class Quality   ha (0.25 acres) [For BF <1 ha (0.25 to 2.5 acres)   acre)     to <4 ha (2.5 to 9.9 acre	R/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha [BR/CM 0.2 to <0.2 ha (0.5 to 5 acms)] [BR/CM 0.5 to 5 acms]			
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetland for Estimating Degree of Interspersion  3 scale. s/tussocks s > 15 cm (6 in.) m (10 in.) dbh pools  None Low Low Moderate  Microtopography Cover Scale  0 = Absent 1 = Present in very small amounts or if more common of marginal quality					
		2 = Present in r amounts of	noderate amounts, but n highest quality	ot of highest quality or in small unts and of highest quality			
22	GRAND 1 (max 100	30- 59 = Cate	gory 1, low wetland functions 2, good/moderate wory 3, superior wetland	tion, condition, quality** etland function, condition, quality** function, condition, quality**			

Site: PUB-003		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
·	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4)  X 0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to )	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland B	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pastidential, fenced past	vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) seck and average. hah, wildlife area, etc. (7) wth forest (5) irk, conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	<b>ду</b>		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Seleston 20.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydrom None or none apparent (1 Recovered (7)  X Recovering (3)  Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) cct only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or ccc  2)  Check all disturbances  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura louble check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and E	evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-001		s): OH:jb	Date: 03/2021				
22 subtotal previous page							
max 10 pts. subtotal	ic 5. Special Wetla	nds					
	cumented raw score for Metric 5 is	30 points or higher, the site is automatically co	onsidered a Category 3 wetland.				
documer  A: A: S: V: Is B: C:	ntation for each selection (photos, or og, fen, wet prairie (10); acidophilic vegosoc. forest (wetl. &/or adj. upland) incl. ensitive geologic feature such as spring ernal pool (5); isolated, perched, or slop land wetland >0.1 acre (0.04 ha) in reservaided channel or floodplain/terrace depross morph. adapt. in >5 trees >10 in. (2000) cological community with global rank (Nown occurrence state/federal threaten.*  "use higher rank where mixed rank or querior/enhanced habitat/use: migratory	nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) lain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) (5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] t/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
1 Metri	ic 6. Plant Commu	nities, Interspersion, Mici	rotopography				
Score all A E S F W X O	and vegetation communities. present using 0 to 3 scale. quatic bed mergent hrub prest judflats pen water <20 acres (8 ha) loss/lichen. Other	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contig [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a significant and is of high quality, or comprises a significant and is of high quality	all part of wetland's vegetation and is of quificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality				
6b. Horiz	contal (plan view) interspersion.	Narrative Description of Vegetation Qu	uality				
Select or  H  N  N  L	. , .	low = Low species diversity &/or domina native species  mod = Native species are dominant component nonnative &/or disturbance tolerare and species diversity moderate to w/o presence of rare, threatened of high = A predominance of native species tolerant native sp absent or virtual	ponent of the vegetation, although nt native species can also be present, moderately high, but generally or endangered species with nonnative sp &/or disturbance lly absent, and high sp diversity and often				
Add or d E M S X N	rage of invasive plants. educt points for coverage. xtensive >75% cover (-5) loderate 25-75% cover (-3) parse 5-25% cover (-1) early absent <5% cover (0) bsent (1)	Mudflat and Open Water Class Quality 0 = Absent <0.1 ha (0.25 acres) [For BF 1 = Low 0.1 to <1 ha (0.25 to 2.5 acres) (0.1 to 0.5 acre)] 2 = Moderate 1 to <4 ha (2.5 to 9.9 acres) 3 = High 4 ha (9.9 acres) or more [BR/C	R/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha es) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]				
6d. Micr Score al V C S	otopography. I present using 0 to 3 scale. egetated hummocks/tussocks oarse woody debris >15 cm (6 in.) tanding dead >25 cm (10 in.) dbh mphibian breeding pools	Hypothetical Wetland for Estimating D  None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if the second seco	Moderate Moderate High  more common of marginal quality of of highest quality or in small				
23	GRAND TOTAL (max 100 pts)	0- 29 = Category 1, low wetland funct 30- 59 = Category 2, good/moderate w 60-100 = Category 3, superior wetland	etland function, condition, quality**				

Site: PUB-004		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (Size) op	en water body (excluding aquati-	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  X 0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland B	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around a 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arouse. Select one or double clor older forest, prairie, savars), shrubland, young 2nd grostdential, fenced pasture, pasidential, pasidential	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter ft) around wetland perimeter ft) ft) around wetland perimeter ft)	(4) er (1)
17	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  cot only one and assign scor  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  lologic regime. Score one or (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat () (2)] Seasonally satura double check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and [	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one None or none apparent (S) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and averag	Check all disturbances of   ☑ mowing ☑ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-004		Rater(s): OH:jb	Date: 03/2021				
26 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Wetlands					
	*If the documented raw score for	Metric 5 is 30 points or higher, the site i	s automatically considered a Category 3 wetland.				
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perci Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedet [*use higher rank where mixe Superior/enhanced habitat/use	multiple values apply in row, score row as single feature with highest point value. Provide action (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). (a); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] are such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5), perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) and (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) adplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) and (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) arith global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] are federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) emixed rank or qualifier] [exclude records which are only "historic"] atat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) in care (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
1	Metric 6. Plant Co	mmunities, Intersper	sion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communit Score all present using 0 to 3 scar Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 Mass/lichen. Other Moss/lichen. Other High (5) Moderately high (4) [BR/C	lle. 0 = Absent or <0.1 ha  [For BR/CM <0.04] 1 = Present and either  moderate quality, of 2 = Present and either  is of moderate quality of 3 = Present and comp  and is of high quality of 1 h	(0.25 acre) contiguous acre ha (0.1 acre)] comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and lity, or comprises a small part and is of high quality rises a significant part or more of wetland's vegetation ty				
	Moderate (3)[BR/CM (5)] Moderately low (2) [BR/CN Low (1) [BR/CM (2)] None (0)	M (3)] and species dive w/o presence of high = A predominance tolerant native sp	sturbance tolerant native species can also be present, rsity moderate to moderately high, but generally rare, threatened or endangered species of native species with nonnative sp &/or disturbance beabsent or virtually absent, and high sp diversity and often				
	6c. Coverage of invasive plants. Add or deduct points for coverage  Extensive >75% cover (-5  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  X Nearly absent <5% cover  Absent (1)	Mudflat and Open Wat 0 = Absent < 0.1 ha (0.3) 1 = Low 0.1 to <1 ha (0.1 to 0.5 acre)] (0)	er Class Quality 25 acres) [For BR/CM <0.04 ha (0.1 acre)] 0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha a (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] es) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Ale. Socks cm (6 in.) O in.) dbh  None  Low  Microtopography Cove 0 = Absent 1 = Present in very sm 2 = Present in modera amounts of highes	all amounts or if more common of marginal quality e amounts, but not of highest quality or in small				
27	GRAND 7 (max 100	O-29 = Category 1, 30-59 = Category 2,	low wetland function, condition, quality** good/moderate wetland function, condition, quality** superior wetland function, condition, quality**				

Site: PUB-(	005		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subt		Metric 1. Wetland	Area (Size) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign   >50 acres (>20.2 ha) (6 pt)   25 to <50 acres (10.1 to <   10 to <25 acres (4 to <10.   3 to <10 acres (1.2 to <4 h   X 0.3 to <3 acres (0.1 to <1.   0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	s) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sui	rounding Land	Use
max 14 pts. subt	total	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around a 25 m to <50 m (82 to <164 je 10 m to <25 m (32 ft to < average <10 m (<32 ft) arouse. Select one or double cor older forest, prairie, savar), shrubland, young 2nd grosidential, fenced pasture, p	wetland perimeter (7)  If the around wetland perimeter  Ether around wetland perimeter  wetland perimeter (0)  wetland perimeter (0)  wetland average.  mah, wildlife area, etc. (7)  with forest (5)  ark, conservation tillage, new	(4) er (1)
1		Metric 3. Hydrolog	ЭУ		
max 30 pts. subt	total	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BF X Precipitation (1) [unless B X Seasonal/intermittent surfill Perennial surface water 3c. Maximum water depth. Sele >0.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/CI 3e. Modifications to natural hydromapholic processes (7)  X Recovered (7)  X Recovering (3)  Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) ct only one and assign scort.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in logic regime. Score one or	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally saturat double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) sed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	6	Metric 4. Habitat A	Alteration and I	Development	
max 20 pts. subt	total	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (1) Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score. or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-005		Rater(s): OH:jb	Date: 03/2021			
26 subtotal previous page						
max 10 pts. subtotal	Metric 5. Special \	Wetlands				
	*If the documented raw score for	Metric 5 is 30 points or higher, the site is	automatically considered a Category 3 wetland.			
	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, percl Island wetland > 0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	sultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) 0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) lain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) is trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] truse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1	Metric 6. Plant Co	mmunities, Interspers	ion, Microtopography			
	6a. Wetland vegetation communit Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 h	lle. 0 = Absent or <0.1 ha (0   For BR/CM <0.04 h. 1 = Present and either comoderate quality, or 2 = Present and either comoderate quality.	.25 acre) contiguous acre			
	6b. Horizontal (plan view) interspersion.  Select only one.  High (5)  Moderately high (4) [BR/CM (5)]  Narrative Description of Vegetation Quality low = Low species diversity &/or dominance of nonnative native species  mod = Native species are dominant component of the ve					
	Moderate (3)[BR/CM (5)] Moderately low (2) [BR/CM Low (1) [BR/CM (2)] None (0)	M (3)] and species divers  w/o presence of ra  high = A predominance o  tolerant native spa	urbance tolerant native species can also be present, ity moderate to moderately high, but generally re, threatened or endangered species native species with nonnative sp &/or disturbance absent or virtually absent, and high sp diversity and often presence of rate, threatened, or endangered species			
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5) Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	Mudflat and Open Water  0 = Absent < 0.1 ha (0.25)  1 = Low 0.1 to <1 ha (0.35)  (0.1 to 0.5 acre)  2 = Moderate 1 to <4 ha	·			
	6d. Microtopography.  Score all present using 0 to 3 sca Vegetated hummocks/tuss Coarse woody debris >15 Standing dead >25 cm (10 Amphibian breeding pools	ale. socks cm (6 in.) 0 in.) dbh  None  Low  Microtopography Cover	r Estimating Degree of Interspersion  Low Moderate Moderate High			
		2 = Present in moderate amounts of highest of	amounts or if more common of marginal quality amounts, but not of highest quality or in small uality or greater amounts and of highest quality			
27	GRAND T (max 100	30- 59 = Category 2, go	w wetland function, condition, quality** od/moderate wetland function, condition, quality** perior wetland function, condition, quality**			

Page 2 of 2

Site: PUB-006		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
·	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around version to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <8 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pastidential, fenced pastidenti	vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) seck and average. hah, wildlife area, etc. (7) wth forest (5) irk, conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	<b>ду</b>		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) cct only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or ccc  2)  Check all disturbances  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura louble check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat A	Alteration and E	evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select  Excellent (7)  Very good (6)  Good (5)  Moderately good (4)  Fair (3)  Poor to fair (2)  Poor (1)  4c. Habitat alteration. Score one  None or none apparent (9)  Recovered (6)  Recovering (3)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-006		Rater(s): OH:jb	Date: 03/2021				
22 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special Wetlands						
	*If the documented raw score for	Metric 5 is 30 points or higher, the site is	automatically considered a Category 3 wetland.				
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perci Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedet [*use higher rank where mixe Superior/enhanced habitat/use	nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] e such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) dain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] truse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
1	Metric 6. Plant Co	mmunities, Interspers	sion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communit Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 h	lle. 0 = Absent or <0.1 ha (    For BR/CM <0.04     1 = Present and either     moderate quality, or     2 = Present and either     is of moderate qual	0.25 acre) contiguous acre na (0.1 acre)] comprises a small part of wetland's vegetation and is of r comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and ity, or comprises a small part and is of high quality ses a significant part or more of wetland's vegetation				
	6b. Horizontal (plan view) intersponded in the select only one.  High (5)  Moderately high (4) [BR/C Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/C Moderately low (2) [BR/C Moderately low (2)]  None (0)	rsity &/or dominance of nonnative or disturbance tolerant e dominant component of the vegetation, although sturbance tolerant native species can also be present, rsity moderate to moderately high, but generally are, threatened or endangered species of native species with nonnative sp &/or disturbance					
	6c. Coverage of invasive plants.  Add or deduct points for coverage Extensive >75% cover (-5) Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) Nearly absent <5% cover Absent (1)	but not always, the  Mudflat and Open Wate  0 = Absent <0.1 ha (0.2  1 = Low 0.1 to <1 ha (0.2  (0.1 to 0.5 acre)]  2 = Moderate 1 to <4 ha	absent or virtually absent, and high sp diversity and often e presence of rate, threatened, or endangered species  Pr Class Quality  25 acres) [For BR/CM < 0.04 ha (0.1 acre)]  .25 to 2.5 acres) [BR/CM 0.04 to < 0.2 ha  a (2.5 to 9.9 acres) [BR/CM 0.2 to < 02 ha (0.5 to 5 acre)]  s) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetland f  ale. socks cm (6 in.) 0 in.) dbh  None  Low  Microtopography Cove 0 = Absent 1 = Present in very sma 2 = Present in moderat amounts of highest	To Estimating Degree of Interspersion  Low Moderate Moderate High  T Scale  all amounts or if more common of marginal quality amounts, but not of highest quality or in small				
23	GRAND 7 (max 100	0- 29 = Category 1, I 30- 59 = Category 2, g	ow wetland function, condition, quality** lood/moderate wetland function, condition, quality** uperior wetland function, condition, quality**				

Site: PUB-007		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pastidential, fenced past	wetland perimeter (7) ft) around wetland perimeter (2 ft) around wetland perimeter (3 ft) around wetland perimeter (4 wetland perimeter (0) (5 eck and average. (7) (6 math, wildlife area, etc. (7) (7) (8 wth forest (5) (8 conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) cct only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or ccc  2)  Check all disturbances  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and E	)evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovered (6) Recovered (6) Recovering (3)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-007	Rater(s):	OH:jb	Date: 03/2021			
22 subtotal previous page						
max 10 pts. subtotal Metric 5	. Special Wetland	s				
	nted raw score for Metric 5 is 30	points or higher, the site is automatically co	onsidered a Category 3 wetland.			
documentation  Bog, fen, Assoc. fc Sensitive Vernal pt Island we Braided of Gross me Ecologic Known o  [*use hi	for each selection (photos, check, wet prairie (10); acidophilic veg., motorest (wetl. &/or adj. upland) incl. >0.2 et geologic feature such as spring/see and (5); isolated, perched, or slope wetland >0.1 acre (0.04 ha) in reservoin channel or floodplain/terrace depressorph. adapt. in >5 trees >10 in. (25 crall community with global rank (Naturaccurrence state/federal threatened/eigher rank where mixed rank or qualif/enhanced habitat/use: migratory son	ultiple values apply in row, score row as single feature with highest point value. Provide on (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) ij. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) erched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) .0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) ain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] deral threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) nixed rank or qualifier] [exclude records which are only "historic"] (use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) 1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 Metric 6	. Plant Communit	ies, Interspersion, Mici	rotopography			
Score all prese Aquatic Emerge Shrub Forest Mudflat:	ent	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contig [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a sig moderate quality, or comprises a sig 2 = Present and either comprises a sign is of moderate quality, or comprises 3 = Present and comprises a significant and is of high quality	all part of wetland's vegetation and is of gnificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality			
6b. Horizontal	(plan view) interspersion.	Narrative Description of Vegetation Qu	uality			
Select only one High (5) Modera Modera Modera Modera	e. ) tely high (4) [BR/CM (5)] te (3)[BR/CM (5)] tely low (2) [BR/CM (3)] [BR/CM (2)]	low = Low species diversity &/or dominance of nonnative or disturbance of native species				
Add or deduct Extensi Modera Sparse	of invasive plants. points for coverage. ve >75% cover (-5) te 25-75% cover (-3) 5-25% cover (-1) absent <5% cover (0) (1)	but not always, the presence of ra  Mudflat and Open Water Class Quality 0 = Absent < 0.1 ha (0.25 acres) [For BR 1 = Low 0.1 to <1 ha (0.25 to 2.5 acres) (0.1 to 0.5 acre)] 2 = Moderate 1 to <4 ha (2.5 to 9.9 acre 3 = High 4 ha (9.9 acres) or more [BR/C	R/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha es) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]			
6d. Microtopoo Score all prese Vegetat Coarse	. ,	Hypothetical Wetland for Estimating D  None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if r  2 = Present in moderate amounts, but n amounts of highest quality  3 = Present in moderate or greater amounts	Moderate Moderate High  more common of marginal quality not of highest quality or in small			
23	GRAND TOTAL (max 100 pts)	0- 29 = Category 1, low wetland funct 30- 59 = Category 2, good/moderate w 60-100 = Category 3, superior wetland	etland function, condition, quality**			

Site: PUB-008		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
·	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around version to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <8 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pastidential, fenced pastidenti	vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) seck and average. hah, wildlife area, etc. (7) wth forest (5) irk, conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) cct only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or ccc  2)  Check all disturbances  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura louble check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and E	evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovered (6) Recovered (6) Recovering (3)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-008		Rater(s): OH:jb	s): OH:jb			
22 subtotal previous page						
max 10 pts. subtotal	Metric 5. Special \	Vetlands				
	*If the documented raw score for	Metric 5 is 30 points or higher, the	ne site is automatically co	onsidered a Category 3 wetland.		
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedel [*use higher rank where mixe Superior/enhanced habitat/us	per for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland. The multiple values apply in row, score row as single feature with highest point value. Provide lection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). (10); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] ture such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) ad, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) cre (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) codplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) on >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] te/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) ere mixed rank or qualifier] [exclude records which are only "historic"] bitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) y) : <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
max 20 pts. subtotal	Metric 6. Plant Co	mmunities, Inters	persion, Mic	rotopography		
max 20 pts. Subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 l	le. 0 = Absent or <	uality, or comprises a signal either comprises a signate quality, or comprises domprises a significant	Juous acre  all part of wetland's vegetation are anificant part but is of low quality ificant part of wetland's vegetation as a small part and is of high quality part or more of wetland's vegetation.	on and	
	6b. Horizontal (plan view) intersponded to Select only one.		iption of Vegetation Quies diversity &/or domina		tolerant	
	High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]	High (5)  Moderately high (4) [BR/CM (5)]  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CM (3)]  Low (1) [BR/CM (2)]  Indive species  mod = Native species are dominant component of the vegetation, although nonnative &/or disturbance tolerant native species can also be presumed and species diversity moderate to moderately high, but generally w/o presence of rare, threatened or endangered species				
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5) Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	Mudflat and Op  0 = Absent <0.  1 = Low 0.1 to  (0.1 to 0.5 a)  2 = Moderate 1	en Water Class Quality   ha (0.25 acres) [For BF <1 ha (0.25 to 2.5 acres)   acre)     to <4 ha (2.5 to 9.9 acre	R/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha es) [BR/CM 0.2 to <02 ha (0.5 to		
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10	Microtopography. Hypothetical Wetland for Estimating Degree of Interspersion				
		0 = Absent 1 = Present in v 2 = Present in v amounts of	very small amounts or if in moderate amounts, but in highest quality	more common of marginal qualit ot of highest quality or in small unts and of highest quality	у	
23	GRAND 1 (max 100	30- 59 = Cate	gory 1, low wetland func- gory 2, good/moderate w gory 3, superior wetland	tion, condition, quality** etland function, condition, qualit function, condition, quality**	y**	

Site: PUB-009	Rater(s): JB; OH		Date: 03/2021
2 Metric 1. Wetlar	id Area (Size) opi	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
Select one size class and ass  >50 acres (>20.2 ha) (  25 to <50 acres (10.1  10 to <25 acres (4 to <  3 to <10 acres (1.2 to  X 0.3 to <3 acres (0.1 to	6 pts) to <20.2 ha) (5) [BR/CM (6)] t10.1 ha) (4) [BR/CM (6)] t24 ha) (3) [BR/CM (5)] t1.2 ha) (2) [BR/CM (3)] to <0.1 ha) (1) [BR/CM (2)]	Sources/assumptions for s	size estimate (list):
max 14 pts. subtotal  2a. Calculate average buffer with wide and with wind with the subtotal with the	vidth. Select only one and assign to 164 ft) or more around wage 25 m to <50 m (82 to <164 terage 10 m to <25 m (32 ft to <80 ters average <10 m (<32 ft) around use. Select one or double control of the or older forest, prairie, savanters), shrubland, young 2nd gro	in score. Do not double check wetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) seck and average. nah, wildlife area, etc. (7)	c. (4)
MODERATELY HIGH.	Residential, fenced pasture, pa , open pasture, row cropping, m	ark, conservation tillage, new	fallow field (3)
max 30 pts. subtotal  3a. Sources of water. Score a High pH groundwater X Other groundwater (3) Precipitation (1) [unless Seasonal/intermittent and	Il that apply.  [BR/CM (5)]  S BR/CM primary source (5)]  surface water (3)  ter (lake or stream) (5)  select only one and assign score  in in.) (2) [BR/CM (3)]  c/CM 0.15 to 0.4 m (6 to <16 in.)  rdrologic regime. Score one or or  t (12)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/s Se. Semi- to permane X Regularly inundat Seasonally inundat (2) Seasonally saturations and average. Seasonally source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6 Metric 4. Habita	t Alteration and D	Development	
None or none apparer Recovered (3) X Recovering (2) Recent or no recovery 4b. Habitat development. Sele Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)	(1) ect only one and assign score.  ne or double check and averag t (9)	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-001		Rater(s): OH:jb	Date: 03/2021			
21 subtotal previous page						
max 10 pts. subtotal	Metric 5. Special \	Wetlands				
	*If the documented raw score for	Metric 5 is 30 points or higher, the s	ite is automatically considered a Category 3 wetland.			
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedel [*use higher rank where mixe Superior/enhanced habitat/us	e multiple values apply in row, score row as single feature with highest point value. Provide lection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  0); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] ure such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) d, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) re (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) adplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) resolved to 10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] tel/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) remediate maker and ror qualifier] [exclude records which are only "historic"] oitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) reservoir solves of invasives OR nonvegetated on mined/excavated land (-10)				
max 20 pts. subtotal	Metric 6. Plant Co	mmunities, Interspe	ersion, Microtopography			
max 20 pts. Subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 l	nity Cover Scale ha (0.25 acre) contiguous acre .04 ha (0.1 acre)] her comprises a small part of wetland's vegetation and is of ty, or comprises a significant part but is of low quality her comprises a significant part of wetland's vegetation and quality, or comprises a small part and is of high quality mprises a significant part or more of wetland's vegetation uality				
	6b. Horizontal (plan view) intersponded only one.  High (5)	low = Low species native specie				
	Moderately high (4) [BR/C Moderate (3)[BR/CM (5)] Moderately low (2) [BR/CN Low (1) [BR/CM (2)]  X None (0)	nonnative &/c and species c w/o presence high = A predominar	es are dominant component of the vegetation, although or disturbance tolerant native species can also be present, diversity moderate to moderately high, but generally of rare, threatened or endangered species note of native species with nonnative sp &/or disturbance as p absent or virtually absent, and high sp diversity and often			
	6c. Coverage of invasive plants.  Add or deduct points for coverage Extensive >75% cover (-5  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover Absent (1)	e. Mudflat and Open N 0 = Absent < 0.1 ha 3) 1 = Low 0.1 to <1 h (0.1 to 0.5 acre 2 = Moderate 1 to <	Nater Class Quality (0.25 acres) [For BR/CM <0.04 ha (0.1 acre)] (0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha (0.25 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] (24 ha (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] (35 acres) or more [BR/CM 2 ha (5 acres) or more]			
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10	Hypothetical Wetla	nd for Estimating Degree of Interspersion			
	Amphibian breeding pools					
		1 = Present in very 2 = Present in mod amounts of high	small amounts or if more common of marginal quality erate amounts, but not of highest quality or in small nest quality erate or greater amounts and of highest quality			
31	GRAND 1 (max 100	30- 59 = Category	1, low wetland function, condition, quality** 2, good/moderate wetland function, condition, quality** 3, superior wetland function, condition, quality**			

Site: PUB-010	Rater(s): JB; OH		Date: 03/2021
2 Metric 1. Wetland	Area (SIZE) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  X 0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
MEDIUM. Buffers averag NARROW. Buffers averag VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	th. Select only one and assig 0 m (164 ft) or more around v e 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <8 average <10 m (<32 ft) arou	n score. Do not double check vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter nd wetland perimeter (0) eck and average. nah, wildlife area, etc. (7) wth forest (5) rk, conservation tillage, new	c. (4) er (1)
max 30 pts.  3a. Sources of water. Score all the High pH groundwater (5)    High pH groundwater (3)   Birty     Precipitation (1) [unless Eight     Seasonal/intermittent surfice water     Perennial surface water     Sources of water. Score all the High pH groundwater (5)   Other groundwater (3)   Birty     Perennial surface water     Perennial surface water     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Seasonal/intermittent surfice     Perennial surface water     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Seasonal/intermittent surfice     Perennial surface water     Seasonal/intermittent surfice     Perennial surface water     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Not precipitation (1) [unless Eight     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Not precipitation (1) [unless Eight     Seasonal/intermittent surfice     Perennial surface water     Sources of water. Score all the light     Not precipitation (1) [unless Eight     Not precipitation (1) [unless Eight     Seasonal/intermittent surfice     Seasonal/intermittent surfice     Not precipitation (1) [unless Eight     Not precipitation (1) [unless Eight     Seasonal/intermittent     Perennial surface water     Sources of water     Not precipitation (1) [unless Eight     Not precipitation (1) [un	nat apply.  R/CM (5)]  R/CM primary source (5)] face water (3) (lake or stream) (5) ect only one and assign score  1.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) plogic regime. Score one or d  1.2)  Check all disturbances  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura louble check and average.  observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6 Metric 4. Habitat	Alteration and D	evelopment	
4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one None or none apparent (5) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and average	Check all disturbances of   ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-010		Rater(s): OH:jb	Date: 03/2021			
22 subtotal previous page						
max 10 pts. subtotal	Metric 5. Special \	Wetlands				
	*If the documented raw score for	Metric 5 is 30 points or higher, the site	is automatically considered a Category 3 wetland.			
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedet [*use higher rank where mixe Superior/enhanced habitat/us	multiple values apply in row, score row as single feature with highest point value. Provide action (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] are such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5), perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) and (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) aplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) and this pool in (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) ith global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] [exclude records which are only "historic"] atat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) : <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
max 20 pts. subtotal	Metric 6. Plant Co	mmunities, Intersper	sion, Microtopography			
max 20 pts. Subidial	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 l	ole. 0 = Absent or <0.1 ha    For BR/CM <0.04	(0.25 acre) contiguous acre  ha (0.1 acre)]  r comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality r comprises a significant part of wetland's vegetation and ality, or comprises a small part and is of high quality brises a significant part or more of wetland's vegetation			
	6b. Horizontal (plan view) intersponded to Select only one.		of Vegetation Quality ersity &/or dominance of nonnative or disturbance tolerant			
	High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]	High (5)  Moderately high (4) [BR/CM (5)]  Moderately low (2) [BR/CM (3)]  Low (1) [BR/CM (2)]  Moderately low (2) [BR/CM (3)]  Low (1) [BR/CM (2)]  Moderately low (2) [BR/CM (3)]  Low (3) [BR/CM (3)]  Moderately low (2) [BR/CM (3)]  Low (3) [BR/CM (3)]  Moderately low (2) [BR/CM (3)]  Low (4) [BR/CM (2)]  Moderately low (5) [BR/CM (3)]  Moderately low (6) [BR/CM (3)]  Moderately low (7) [BR/CM (3)]  Moderately low (8) [BR/CM (3)]  Moderately low (9) [BR/CM (3)]  Moderately				
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5 Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	e. Mudflat and Open Wa  ) 0 = Absent <0.1 ha (0  -3) 1 = Low 0.1 to <1 ha	the presence of rate, threatened, or endangered species  ter Class Quality  .25 acres) [For BR/CM <0.04 ha (0.1 acre)]  0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha  ha (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]  es) or more [BR/CM 2 ha (5 acres) or more]			
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetland for Estimating Degree of Interspersion  Low Low Moderate Moderate High  Microtopography Cover Scale				
		0 = Absent 1 = Present in very sr 2 = Present in modera amounts of higher	nall amounts or if more common of marginal quality ate amounts, but not of highest quality or in small			
23	GRAND 1 (max 100	30- 59 = Category 2,	low wetland function, condition, quality** good/moderate wetland function, condition, quality** superior wetland function, condition, quality**			

Site: PUB-011		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	1 subtotal	Metric 1. Wetland	Area (Size)	pen water body (excluding aquati-	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign   >50 acres (>20.2 ha) (6 p   25 to <50 acres (10.1 to <   10 to <25 acres (4 to <10   3 to <10 acres (1.2 to <4     0.3 to <3 acres (0.1 to <1)   X 0.1 to <0.3 acre (0.04 to <0.1 acre (0.04 to <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
may 14 mts	1 subtotal	Metric 2. Upland E	Buffers and Su	rrounding Land	Use
max 14 pts.	Sublotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around e 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to average <10 m (<32 ft) are use. Select one or double or older forest, prairie, sava s), shrubland, young 2nd gresidential, fenced pasture, p	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. unnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
	13	Metric 3. Hydrolog	gy		
max 30 pts.	subtotal	3a. Sources of water. Score all the light physical states of the light phy	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign scc  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in the color of the color	Part of wetland/up Part of riparian or 3d. Duration inundation/s ore. Semi- to permane X Regularly inundat Seasonally inundat Seasonally satura of double check and average.  es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)
	6	Metric 4. Habitat	Alteration and	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select  Excellent (7)  Very good (6)  Good (5)  Moderately good (4)  Fair (3)  Poor to fair (2)  Recovered (6)  Recovered (6)	only one and assign score.  or double check and avera	Check all disturbances of mowing grazing clearcutting selective cutting farming	
		X Recovering (3) Recent or no recovery (1)		toxic pollutants	nutrient enrichment

Site: PUB-011		Rater(s): OH:jb	Date: 03/2021				
21 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Wetlands					
	*If the documented raw score for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.						
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain, Gross morph. adapt. in >5 tre Ecological community with glo Known occurrence state/fedet [*use higher rank where mix Superior/enhanced habitat/us	(photos, checklists, maps, resource s dophilic veg., mossy substrate >10 sq.m, spupland) incl. >0.25 acre (0.1 ha); old growth the as spring/seep, sink, losing/underground hed, or slope wetland (4); headwater wetland (4 ha) in reservoir, river, or perennial water > (ferrace depressions (floodplain pool, slough the ses >10 in. (25 cm) dbh: buttress, multitrunk/ obal rank (NatureServe): G1*(10), G2*(5), Garal threatened/endangered species (10); other drank or qualifier] [exclude records which e: migratory songbird/waterfowl (5); in-reservant of the service	6 ft (2 m) deep (5) n, oxbow, meander scar, etc.) (3) stool, stilted, shallow roots/tip-up, or pneumatophores (3) 3*(3) [*use higher rank where mixed rank or qualifier] er rare species with global rank G1*(10), G2*(5), G3*(3)				
1	Metric 6. Plant Co	mmunities, Interspe	rsion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 l	1	a (0.25 acre) contiguous acre 4 ha (0.1 acre)] er comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality er comprises a significant part of wetland's vegetation and lality, or comprises a small part and is of high quality prises a significant part or more of wetland's vegetation				
	Moss/lichen. Other and is of high quality  6b. Horizontal (plan view) interspersion.  Select only one.  High (5)    Marrative Description of Vegetation Quality						
	Moderately high (4) [BR/C Moderate (3)[BR/CM (5)] Moderately low (2) [BR/CN Low (1) [BR/CM (2)] None (0)	mod = Native species nonnative &/or (3)] and species div w/o presence of high = A predominance	are dominant component of the vegetation, although disturbance tolerant native species can also be present, versity moderate to moderately high, but generally frare, threatened or endangered species e of native species with nonnative sp &/or disturbance sp absent or virtually absent, and high sp diversity and often				
	6c. Coverage of invasive plants. Add or deduct points for coverage	but not always,	the presence of rate, threatened, or endangered species				
	Extensive >75% cover (-5  Moderate 25-75% cover (-5  Sparse 5-25% cover (-1)  X Nearly absent <5% cover  Absent (1)	1 = Low 0.1 to <1 ha (0.1 to 0.5 acre)] 2 = Moderate 1 to <4	0.25 acres) [For BR/CM <0.04 ha (0.1 acre)] (0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha  ha (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] res) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tus:  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	ale. socks cm (6 in.) 0 in.) dbh None Low	A for Estimating Degree of Interspersion  Low Moderate Moderate High				
		2 = Present in moder amounts of higher	mall amounts or if more common of marginal quality ate amounts, but not of highest quality or in small				
22	GRAND 1 (max 100	30- 59 = Category 2	, low wetland function, condition, quality** , good/moderate wetland function, condition, quality** , superior wetland function, condition, quality**				

Site: PUB-012		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 l  X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pastidential, fenced pastidenti	wetland perimeter (7) ft) around wetland perimeter (2 ft) around wetland perimeter (12 ft) around wetland perimeter (13 ft) around wetland perimeter (14 ft) neck and average. nah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	ЭУ		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selector > 0.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydrom None or none apparent (1) Recovered (7)  X Recovering (3)  Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) act only one and assign score act) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in.) blogic regime. Score one or other	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane X Regularly inundat Seasonally inundat Ouble check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and D	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one Recovered (6) X Recovering (3)	only one and assign score. or double check and averag	Check all disturbances of  ☐ mowing ☐ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-012		Rater(s): OH:jb	Date: 03/2021
21 subtotal previous page			
max 10 pts. subtotal	Metric 5. Special \	Wetlands	
	*If the documented raw score for	Metric 5 is 30 points or higher, the site	s automatically considered a Category 3 wetland.
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedel [*use higher rank where mixe Superior/enhanced habitat/us	(photos, checklists, maps, resource sp dophilic veg., mossy substrate >10 sq.m, sph- ipland) incl. >0.25 acre (0.1 ha); old growth (1 th as spring/seep, sink, losing/underground s hed, or slope wetland (4); headwater wetland tha) in reservoir, river, or perennial water >6 ferrace depressions (floodplain pool, slough, es >10 in. (25 cm) dbh: buttress, multirunk/s obal rank (NatureServe): G1*(10), G2*(5), G3 ral threatened/endangered species (10); othe ed rank or qualifier] [exclude records which a e: migratory songbird/waterfowl (5); in-reserv	[1st order perennial or above] (3) ft (2 m) deep (5) oxbow, meander scar, etc.) (3) ool, stilted, shallow roots/tip-up, or pneumatophores (3) *(3) [*use higher rank where mixed rank or qualifier] r rare species with global rank G1*(10), G2*(5), G3*(3)
1	Metric 6. Plant Co	mmunities, Intersper	sion, Microtopography
max 20 pts. subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 l	lle. 0 = Absent or <0.1 ha   For BR/CM <0.04   1 = Present and either	(0.25 acre) contiguous acre ha (0.1 acre)] comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and lity, or comprises a small part and is of high quality rises a significant part or more of wetland's vegetation
	6b. Horizontal (plan view) intersp. Select only one.  High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CN  Low (1) [BR/CM (2)]	Iow = Low species divinative species and nonnative &/or can and species diversity (3)]  Iow = Low species divinative species and nonnative &/or can and species divinative w/o presence of	re dominant component of the vegetation, although isturbance tolerant native species can also be present, ersity moderate to moderately high, but generally rare, threatened or endangered species
	<ul><li> X  None (0)</li><li>6c. Coverage of invasive plants.</li></ul>	tolerant native s	of native species with nonnative sp &/or disturbance of absent or virtually absent, and high sp diversity and often the presence of rate, threatened, or endangered species
	Add or deduct points for coverage Extensive >75% cover (-5 Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	0 = Absent < 0.1 ha (0.3) 1 = Low 0.1 to <1 ha (0.1 to 0.5 acre)] (0) 2 = Moderate 1 to <4 li	ter Class Quality 25 acres) [For BR/CM <0.04 ha (0.1 acre)] 0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha at (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] as) or more [BR/CM 2 ha (5 acres) or more]
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	ale. socks cm (6 in.) 0 in.) dbh  None  Microtopography Cov 0 = Absent 1 = Present in very sm	all amounts or if more common of marginal quality
		amounts of highes	te amounts, but not of highest quality or in small t quality te or greater amounts and of highest quality
22	GRAND 1 (max 100	30- 59 = Category 2,	low wetland function, condition, quality** good/moderate wetland function, condition, quality** superior wetland function, condition, quality**

Site: PUB-013		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (Size) op (8	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 pi 25 to <50 acres (10.1 to < 10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 li  0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
max 14 pts. subtotal	NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of X LOW. Old field (>10 years MODERATELY HIGH. Re	th. Select only one and assign (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) around use. Select one or double clor older forest, prairie, savars), shrubland, young 2nd gro	gn score. Do not double check wetland perimeter (7) ft) around wetland perimeter 32 ft) around wetland perimeter and wetland perimeter (0) neck and average. nah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	<. (4) er (1)
max 30 pts. subtotal	Metric 3. Hydrolog  3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BF X Precipitation (1) [unless B X Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selector > 0.7 m (27.6 in.) (3)  X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/Cl 3e. Modifications to natural hydrom None or none apparent (1) Recovered (7)  X Recovering (3)  Recent or no recovery (1)	nat apply.  R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign scor  act only one and assign scor  blogic regime. Score one or (2)  Check all disturbance:	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat Ouble check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)
6	Metric 4. Habitat A	Alteration and [	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one None or none apparent (9) Recovered (6)	only one and assign score.  or double check and averag	Check all disturbances of   ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging

Site: PUB-013	Rater(s): (	DH:jb	Date: 03/2021
37 subtotal previous page			
max 10 pts. subtotal Metric 5. S	Special Wetland	s	
*If the documented	raw score for Metric 5 is 30	points or higher, the site is automatically co	ensidered a Category 3 wetland.
documentation for e Bog, fen, wet Assoc. forest Sensitive geol Vernal pool (5 Island wetland Braided chann Gross morph. Ecological cor Known occurr [*use higher Superior/enha	each selection (photos, checl prairie (10); acidophilic veg., mos (wetl. &/or adj. upland) incl. >0.2: logic feature such as spring/seep (b); isolated, perched, or slope we d >0.1 acre (0.04 ha) in reservoir nel or floodplain/terrace depressi adapt. in >5 trees >10 in. (25 cm mmunity with global rank (Nature rence state/federal threatened/en rank where mixed rank or qualificanced habitat/use: migratory song	ly in row, score row as single feature with holists, maps, resource specialist concurrences substrate >10 sq.m, sphagnum or other mose for acre (0.1 ha); old growth (10); mature >18 in. (40), sink, losing/underground stream, cave, waterfaltland (4); headwater wetland [1st order perennial, river, or perennial water >6 ft (2 m) deep (5) mos (floodplain pool, slough, oxbow, meander scan) dbh: buttress, multitrunk/stool, stilted, shallow reserve): G1*(10), G2*(5), G3*(3) [*use higher rand dangered species (10); other rare species with gler] [exclude records which are only "historic"] abird/waterfowl (5); in-reservoir buttonbush (4); ot 0 EITHER >80% cover of invasives OR nonveget.	De, data sources, references, etc).  (5); muck, organic soil layer (3)  (5); muck, organic soil layer (3)  (5); cm) dbh (5) [exclude pine plantation]  (II, rock outcrop/cliff (5)  or above] (3)  ar, etc.) (3)  oots/tip-up, or pneumatophores (3)  ox where mixed rank or qualifier]  lobal rank G1*(10), G2*(5), G3*(3)  ther fish/wildlife management/designation (3)
1 Metric 6. P	Plant Communit	ies, Interspersion, Micr	otopography
6a. Wetland vegeta Score all present us Aquatic bed Emergent Shrub Forest Mudflats Open water Moss/lichen	sing 0 to 3 scale. <20 acres (8 ha)	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contiguing [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a significant and either comprises a significant and is of high quality	Il part of wetland's vegetation and is of nificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality
6b. Horizontal (plan Select only one. High (5) Moderately I	h view) interspersion.  high (4) [BR/CM (5)] b)[BR/CM (5)] low (2) [BR/CM (3)]	Narrative Description of Vegetation Qu  low = Low species diversity &/or domina native species  mod = Native species are dominant comp	conent of the vegetation, although to native species can also be present, moderately high, but generally or endangered species
Moderate 25 Sparse 5-25		•	/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha s) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]
Coarse woo Standing de		None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if no content in moderate amounts, but no	Moderate Moderate High
20	RAND TOTAL	amounts of highest quality  3 = Present in moderate or greater amounts  0-29 = Category 1, low wetland function  30-59 = Category 2, good/moderate wetland for the second	unts and of highest quality  ion, condition, quality** etland function, condition, quality**

Site: PUB-014			Rater(s): JB; OH		Date: 03/2021
	2 btotal	Metric 1. Wetland	Area (Size)	ppen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
,		Select one size class and assign  >50 acres (>20.2 ha) (6 pi 25 to <50 acres (10.1 to < 10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 h  X 0.3 to <3 acres (0.1 to <1. 0.1 to <0.3 acre (0.04 to <0.1 acre (0.04 to <0.1 acre (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for GPS	size estimate (list):
	1	Metric 2. Upland E	Buffers and Su	rrounding Land	Use
max 14 pts. sub	ototal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to average <10 m (<32 ft) ar use. Select one or double or older forest, prairie, savas), shrubland, young 2nd gesidential, fenced pasture,	d wetland perimeter (7) 64 ft) around wetland perimeter <82 ft) around wetland perimet ound wetland perimeter (0) check and average. annah, wildlife area, etc. (7) rowth forest (5) park, conservation tillage, new	(4) er (1)
1	17	Metric 3. Hydrolog	ЭУ		
max 30 pts. sub	ototal	3a. Sources of water. Score all the light ph groundwater (5)   X Other groundwater (3) [BF X Precipitation (1) [unless B X Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selector > 0.7 m (27.6 in.) (3)   X 0.4 to 0.7 m (16 to 27.6 in.) <0.4 m (<16 in.) (1) [BR/Cl 3e. Modifications to natural hydrom None or none apparent (1) Recovered (7)   X Recovering (3)   Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) act only one and assign soc a.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 iologic regime. Score one o	Part of wetland/u Part of riparian or 3d. Duration inundation/s ore. Semi- to permane X Regularly inundat Seasonally inundat n.) (2)] Seasonally satura r double check and average.  es observed point source (non filling/grading road bed/RR trace dredging	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) sed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	6	Metric 4. Habitat A	Alteration and	Development	
max 20 pts. sub	btotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (2) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score or double check and avera	.  Check all disturbances of mowing  ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-014		Rater(s): OH:jb	Date: 03/2021				
26 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Vetlands					
,	*If the documented raw score for	ed raw score for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.					
	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, percl Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	(photos, checklists, maps, resource spophilic veg., mossy substrate >10 sq.m, sphpland) incl. >0.25 acre (0.1 ha); old growth the has spring/seep, sink, losing/underground shed, or slope wetland (4); headwater wetland ha) in reservoir, river, or perennial water					

Site: PUB-015		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	tric 1. Wetland	Area (Size) or	oen water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	to one size class and assign s >50 acres (>20.2 ha) (6 pts 25 to <50 acres (10.1 to <2 10 to <25 acres (4 to <10.1 3 to <10 acres (1.2 to <4 ha 0.3 to <3 acres (0.1 to <1.2 0.1 to <0.3 acre (0.04 to <1.2	s) (0.2 ha) (5) [BR/CM (6)] ha) (4) [BR/CM (6)] a) (3) [BR/CM (5)] ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
max 14 pts. subtotal 2a. C	alculate average buffer width WIDE. Buffers average 50 MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers a attensity of surrounding land u VERY LOW. 2nd growth or LOW. Old field (>10 years)	i. Select only one and assim (164 ft) or more around 25 m to <50 m (82 to <164 to <10 m to <25 m (32 ft to <10 m (32 ft) around (32 ft) aroun	4 ft) around wetland perimeter 82 ft) around wetland perimeter und wetland perimeter (0) theck and average.  nnah, wildlife area, etc. (7) owth forest (5) wark, conservation tillage, new	κ. (4) ∋r (1)
max 30 pts. subtotal  3a. S  3c. S  3e. M	ources of water. Score all that High pH groundwater (5) Other groundwater (3) [BR/K] Precipitation (1) [unless BR Seasonal/intermittent surfater of Perennial surface water (1) Maximum water depth. Selection (27.6 in.) (3) Out to 0.7 m (16 to 27.6 in.) (3) Out to 0.7 m (16 to 27.6 in.) (4) Out (10 in.) (1) [BR/CM of In.) (10 [BR/CM of	at apply.  (CM (5)]  (CM primary source (5)]  ce water (3)  lake or stream) (5)  t only one and assign sco  (2) [BR/CM (3)]  1 0.15 to 0.4 m (6 to <16 in order one or organization.	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane X Regularly inundat Seasonally inundat a.) (2)] Seasonally satura double check and average. es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) inturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)
6 <b>Me</b>	tric 4. Habitat A	Iteration and I	Development	
4b. H	ubstrate disturbance. Score of None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1) labitat development. Select of Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Poor (1) labitat alteration. Score one of None or none apparent (9)	nly one and assign score.	Check all disturbances of   ☑ mowing ☑ grazing □ clearcutting	observed ☑ shrub/sapling removal ☐ herbaceous/aquatic bed removal ☐ woody debris removal ☐ sedimentation
<u> </u>	None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)		selective cutting farming toxic pollutants	☐ dredging ☐ nutrient enrichment

Site: PUB-015	Rater(s): OH:jb	Date: 03/2021
31 subtotal previous page		
Metric 5. Special V	Wetlands	
*If the documented raw score for	Metric 5 is 30 points or higher, the site is a	utomatically considered a Category 3 wetland.
documentation for each selection  Bog, fen, wet prairie (10); aci Assoc. forest (wetl. &/or adj. of Sensitive geologic feature su Vernal pool (5); isolated, percomposed listend wetland >0.1 acre (0.0) Braided channel or floodplain Gross morph. adapt. in >5 tre Ecological community with gli Known occurrence state/fede [*use higher rank where mix Superior/enhanced habitat/us	dophilic veg., mossy substrate >10 sq.m, sphagnupland) incl. >0.25 acre (0.1 ha); old growth (10); ich as spring/seep, sink, losing/underground streathed, or slope wetland (4); headwater wetland [1st 4 ha) in reservoir, river, or perennial water >6 ft (2/terrace depressions (floodplain pool, slough, oxbes >10 in. (25 cm) dbh: buttress, multitrunk/stool, obal rank (NatureServe): G1*(10), G2*(5), G3*(3) ral threatened/endangered species (10); other rared rank or qualifier] [exclude records which are or ie: migratory songbird/waterfowl (5); in-reservoir b	alist concurrence, data sources, references, etc).  Im or other moss (5); muck, organic soil layer (3)  mature >18 in. (45 cm) dbh (5) [exclude pine plantation]  m, cave, waterfall, rock outcrop/cliff (5)  to order perennial or above] (3)  to m) deep (5)  ow, meander scar, etc.) (3)  stilted, shallow roots/tip-up, or pneumatophores (3)  [*use higher rank where mixed rank or qualifier]  to species with global rank G1*(10), G2*(5), G3*(3)
1 Metric 6. Plant Co	mmunities, Interspersi	on, Microtopography
6a. Wetland vegetation commun Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 Moss/lichen. Other	ale. 0 = Absent or <0.1 ha (0.2    For BR/CM <0.04 ha   1 = Present and either core	25 acre) contiguous acre
6b. Horizontal (plan view) intersponder only one.  High (5)  Moderately high (4) [BR/CM (5)]  Moderately low (2) [BR/CC Low (1) [BR/CM (2)]	ersion.    Narrative Description of Normalive Species diversity	degetation Quality by &/or dominance of nonnative or disturbance tolerant dominant component of the vegetation, although rbance tolerant native species can also be present, y moderate to moderately high, but generally be, threatened or endangered species hative species with nonnative sp &/or disturbance
6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5) Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	tolerant native sp ab but not always, the part of the	sent or virtually absent, and high sp diversity and often oresence of rate, threatened, or endangered species
6d. Microtopography.  Score all present using 0 to 3 so  Vegetated hummocks/tus  Coarse woody debris >15  Standing dead >25 cm (1  Amphibian breeding pools	ale. socks cm (6 in.) 0 in.) dbh  None Low  Microtopography Cover S 0 = Absent 1 = Present in very small a 2 = Present in moderate a amounts of highest qu	amounts or if more common of marginal quality mounts, but not of highest quality or in small lality
GRAND 1	0- 29 = Category 1, low 30- 59 = Category 2, goo	wetland function, condition, quality** d/moderate wetland function, condition, quality** erior wetland function, condition, quality**

Site: PUB-016	Pater(s): JB; OH			Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (Size)	pen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4   0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <1)  X <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
max 14 pts. subtotal	NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of X LOW. Old field (>10 years MODERATELY HIGH. Re	th. Select only one and ass o m (164 ft) or more around e 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to < average <10 m (<32 ft) arouse. Select one or double or or older forest, prairie, sava s), shrubland, young 2nd gr	ign score. Do not double check wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. nnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	k. (4) er (1)
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water sc. Maximum water depth. Selemon 1.00 per solution with the selemon 1.00 per solution 1.00 per solution with the selemon 1.00 per solution with the selemon 1.00 per solution	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign scc  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in old	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane X Regularly inundat Seasonally inundat a.) (2)] Seasonally satura double check and average.  es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) sed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat A	Alteration and	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and avera	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-016		Rater(s): OH:jb		Date: 03/2021
31 subtotal previous page				
max 10 pts. subtotal	etric 5. Special V	Vetlands		
	the documented raw score for	Metric 5 is 30 points or highe	r, the site is automatically co	onsidered a Category 3 wetland.
	Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perch Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	(photos, checklists, maps, reophilic veg., mossy substrate >1 pland) incl. >0.25 acre (0.1 ha); of has spring/seep, sink, losing/united, or slope wetland (4); headw. ha) in reservoir, river, or perenn terrace depressions (floodplain plass >10 in. (25 cm) dbh: buttress, bal rank (NatureServe): G1*(10), all threatened/endangered specied rank or qualifier] [exclude recoes: migratory songbird/waterfowl (	esource specialist concurrence of sq.m, sphagnum or other moss old growth (10); mature >18 in. (4 derground stream, cave, waterfalater wetland [1st order perennial ial water >6 ft (2 m) deep (5) ool, slough, oxbow, meander scamultitrunk/stool, stilted, shallow r G2*(5), G3*(3) [*use higher ranles (10); other rare species with girds which are only "historic"] 5); in-reservoir buttonbush (4); of	ce, data sources, references, etc). s (5); muck, organic soil layer (3) 45 cm) dbh (5) [exclude pine plantation] II, rock outcrop/cliff (5) or above] (3) ar, etc.) (3) roots/tip-up, or pneumatophores (3)
1 <b>M</b>	etric 6. Plant Co	mmunities, Inte	rspersion, Micr	otopography
	Wetland vegetation community ore all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 h	1e. 0 = Absent   [For BR]   1 = Present   moderal   2 = Present   is of mo   3 = Present   1   1   1   1   1   1   1   1   1	e quality, or comprises a signand either comprises a signand either comprises a signalerate quality, or comprises	uous acre  Il part of wetland's vegetation and is of nificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality part or more of wetland's vegetation
6b.	Horizontal (plan view) interspe	ersion. Narrative De	scription of Vegetation Qu	ality
	ect only one.  High (5)  Moderately high (4) [BR/CI  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CM  Low (1) [BR/CM (2)]  None (0)	low = Low s	pecies diversity &/or domina species e species are dominant competive &/or disturbance tolerant pecies diversity moderate to esence of rare, threatened of dominance of native species at native sp absent or virtuall	nce of nonnative or disturbance tolerant conent of the vegetation, although at native species can also be present, moderately high, but generally or endangered species with nonnative sp &/or disturbance ly absent, and high sp diversity and often
	Coverage of invasive plants. d or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover (Absent (1)	Mudflat and 0 = Absent 3  1 = Low 0.1 (0.1 to 0)  2 = Moderat	Open Water Class Quality <0.1 ha (0.25 acres) [For BR to <1 ha (0.25 to 2.5 acres) .5 acre)]	[BR/CM 0.04 to <0.2 ha s) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]
	. Microtopography. ore all present using 0 to 3 sca Vegetated hummocks/tuss Coarse woody debris >15 Standing dead >25 cm (10 Amphibian breeding pools	Hypothetica  ide.  iocks cm (6 in.) in.) dbh  None  Microtopogr 0 = Absent 1 = Present 2 = Present amounts	I Wetland for Estimating Do	Moderate Moderate High  more common of marginal quality ot of highest quality or in small
32	GRAND T (max 100	30-59 = C	ategory 1, low wetland functi ategory 2, good/moderate we ategory 3, superior wetland f	etland function, condition, quality**

Site: PUB-017		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (Size)	pen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4   0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <1)  X <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
3	Metric 2. Upland E	Buffers and Su	rrounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	o m (164 ft) or more around e 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to average <10 m (<32 ft) are use. Select one or double or older forest, prairie, sava s), shrubland, young 2nd gr	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. nnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selemon 1.00 per solution 1.00	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign scc  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in the color of the color	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane X Regularly inundat Seasonally inundat a.) (2)] Seasonally satura double check and average.  es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat A	Alteration and	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and avera	Check all disturbances o  ☑ mowing ☑ grazing	observed   shrub/sapling removal   herbaceous/aquatic bed removal   woody debris removal   sedimentation   dredging   nutrient enrichment

Site: PUB-017		Rater(s): OH:jb		Date: 03/2021		
22 subtotal previous page						
max 10 pts. subtotal	Metric 5. Special V	Vetlands				
*1	f the documented raw score for	Metric 5 is 30 points or high	er, the site is automatically co	onsidered a Category 3 wetland.		
	ocumentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perct Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.  nultiple values apply in row, score row as single feature with highest point value. Provide  tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3)  adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation]  e such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5)  perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3)  (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5)  alain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3)  5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3)  th global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier]  rederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3)  mixed rank or qualifier] [exclude records which are only "historic"]  at/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3)  <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1	Metric 6. Plant Co	mmunities, Inte	erspersion, Micr	otopography		
	a. Wetland vegetation communit score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats X Open water <20 acres (8 h	le. 0 = Absent	ate quality, or comprises a signt and either comprises a signoderate quality, or comprises	uous acre  Il part of wetland's vegetation and is of nificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality part or more of wetland's vegetation		
6	b. Horizontal (plan view) interspe	ersion. Narrative D	escription of Vegetation Qu	ality		
	Belect only one.  High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CM  Low (1) [BR/CM (2)]  None (0)	low = Low :   native   nativ	species diversity &/or domina e species e species are dominant comp ative &/or disturbance tolerar species diversity moderate to presence of rare, threatened of adominance of native species ant native sp absent or virtual	connent of the vegetation, although at native species can also be present, moderately high, but generally or endangered species with nonnative sp &/or disturbance ly absent, and high sp diversity and often		
	c. Coverage of invasive plants.  dd or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  X Nearly absent <5% cover (Absent (1))	Mudflat and 0 = Absent 3) 1 = Low 0.  (0.1 to 2 = Modera	Open Water Class Quality   <0.1 ha (0.25 acres) [For BR   to <1 ha (0.25 to 2.5 acres)   0.5 acre)	(CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha s) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]		
	Sd. Microtopography. Score all present using 0 to 3 sca Vegetated hummocks/tuss Coarse woody debris >15 Standing dead >25 cm (10 Amphibian breeding pools	Hypotheticalle. socks cm (6 in.) in.) dbh  None  Microtopog 0 = Absent 1 = Presen 2 = Presen amoun	Low Low raphy Cover Scale t in very small amounts or if r	egree of Interspersion  Moderate Moderate High  more common of marginal quality ot of highest quality or in small		
23	GRAND T (max 100	OTAL $30-59=0$	Category 1, low wetland funct category 2, good/moderate w category 3, superior wetland f	etland function, condition, quality**		

Metric 1. Wetland Area (size)    Motion   Metric 2. Wetland Area (size)	Site: PUB-018	Rater(s	):JB; OH		Date: 03/2021
Sol acres (1-20.2 ha) (6 pts)   Sol (First)   Sol acres (1-5 col acres (1-1 to -20.2 ha) (5) [BR/CM (6)]   10 to <25 acres (4 to -10.1 ha) (4) [BR/CM (6)]   3 to <10 acres (1.2 to -4 ha) (3) [BR/CM (6)]   GPS	max 6 pts. subtotal		(SIZE) oper	water body (excluding aquation	beds and seasonal mudflats) is >20 acres
2a. Calculate average buffer width. Select only one and assign score. Do not double check.    WIDE. Buffers average 50 m (164 ft) or more around wetland perimeter (7)   MRROW. Buffers average 50 m (164 ft) or more around wetland perimeter (4)   NARROW. Buffers average 25 m to -50 m (32 ft to -62 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 30 m to -50 m (82 ft to -62 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 30 m (164 ft) or not -61 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 10 m to -25 m (32 ft to -62 ft) around wetland perimeter (1)   VERY LOW. 2nd growth or older forest, prairie, savannah, wildlife area, etc. (7)   LOW. Old field (-10 years), shrubland, young 2nd growth forest (5)   MODERATELY HIGH. Residential, fenced pasture, park, conservation tillage, new fallow field (3)   High. Urban, industrial, open pasture, row cropping, mining, construction (1)    Metric 3. Hydrology   3a. Sources of water. Score all that apply.   High pH groundwater (5)   Other groundwater (3) [BR/CM (5)]   Perennial surface water (lake or stream) (5)   Seasonal/Intermitient surface water (3)   Perennial surface water (lake or stream) (5)   Seasonal/Intermitient surface water (3)   Active of the stream of the stream or valued corridor (1)   Part of wetland/upland (e.g., forest), complex (1)   P	2   1   3   0   X	50 acres (>20.2 ha) (6 pts) 5 to <50 acres (10.1 to <20.2 ha) (5) 0 to <25 acres (4 to <10.1 ha) (4) [B to <10 acres (1.2 to <4 ha) (3) [BR/ .3 to <3 acres (0.1 to <1.2 ha) (2) [B 0.1 to <0.3 acre (0.04 to <0.1 ha) (1)	BR/CM (6)] CM (5)] BR/CM (3)]	·	size estimate (list):
2a. Calculate average buffer width. Select only one and assign score. Do not double check.    WIDLE. Buffers average 55 m to <50 m (82 to <164 ft) around wetland perimeter (4)   NARROW. Buffers average 10 m to <25 m (32 to <164 ft) around wetland perimeter (9)   VERY NARROW. Buffers average > 10 m to <25 m (32 to <164 ft) around wetland perimeter (1)   VERY LOW. 2nd growth or older forest, prairie, savannah, wildlife area, etc. (7)   LOW. Old feld (-10 years), shrubband, young 2nd growth forest (5)   MODERATELY HIGH. Residential, fenced pasture, park, conservation tillage, new fallow field (3)   Wetric 3. Hydrology   Metric 3. Hydrology   Sa. Sources of water. Score all that apply.   Sources of water. Score all that apply.   High pH groundwater (3) [BR/CM (5)]   Percipitation (1) [Unless BR/CM (5)]   Percipitation (1) [Unless BR/CM (5)]   Part of ipparian or upland corridor (1)   Recovered (3)   Recent or no recovery (1)   Part of ipparian   Part of the pa		ic 2. Upland Buffers	s and Surr	ounding Land	Use
Sa. Sources of water. Score all that apply.   100-year floodplain (1)   100-year floodplain (1	2a. Calc V N N X V 2b. Inter L	VIDE. Buffers average 50 m (164 ft)  MEDIUM. Buffers average 25 m to <5  MEROW. Buffers average 10 m to <  MERY NARROW. Buffers average <1  mailtain stream of the stream	or more around w 50 m (82 to <164 f <25 m (32 ft to <82 10 m (<32 ft) arour one or double che est, prairie, savann d, young 2nd grow enced pasture, par	etland perimeter (7) t) around wetland perimeter th) around wetland perimete d wetland perimeter (0) ck and average. ah, wildlife area, etc. (7) th forest (5) k, conservation tillage, new	(4) er (1)
3a. Sources of water. Score all that apply.    High pH groundwater (5)   D0-year floodplain (1)   Between stream/lake and other human use (1)   Part of weltand/upland (e.g., forest), complex (1)   Part of priarian or upland corridor (1)   Part of weltand/upland (e.g., forest), complex (1)   Part of veltand/upland (e.g., forest), complex (1)   Part of veltand/upland (e.g., forest), complex (1)   Part of veltand/upland (e.g., forest), complex (1)   Part of weltand/upland (e.g., forest), complex (1)   Part of veltand/upland (e.g., forest), complex (1)   Part of veltand/upland (e.g., forest), complex (1)   Part of weltand/upland (e.g., forest), complex (1)   Part of weltand/upland (e.g., forest), complex (1)   Part of weltand/upland (e.g., forest), complex (1)   Part	13 Metr	ic 3. Hydrology			
### subtotal  4a. Substrate disturbance. Score one or double check and average.  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select only one and assign score.  Excellent (7)  Very good (6)  Good (5)  Moderately good (4)  Fair (3)  Check all disturbances observed	3a. Soul	digh pH groundwater (5) Other groundwater (3) [BR/CM (5)] Precipitation (1) [unless BR/CM prima deasonal/intermittent surface water (3) Perennial surface water (lake or streximum water depth. Select only one 0.7 m (27.6 in.) (3) 0.4 to 0.7 m (16 to 27.6 in.) (2) [BR/CD 0.4 m (<16 in.) (1) [BR/CM 0.15 to 0.00 one or none apparent (12) eccovered (7) Recovering (3) Recent or no recovery (1)	an) (5) and assign score.  (M (3)] 4 m (6 to <16 in.) be. Score one or do k all disturbances to be (including culverto ke	100-year floodplai   Between stream/la   Part of wetland/up   Part of riparian or   3d. Duration inundation/s   Semi- to permane   X Regularly inundation/s   Seasonally inundation/s   Seasonally satura buble check and average.   Deserved   point source (none)   filling/grading   road bed/RR track	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) aturation. Score one or dbl. check & avg. ntly inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)
4a. Substrate disturbance. Score one or double check and average.  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select only one and assign score. Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3)  Check all disturbances observed	6 <b>Me</b> tr	ic 4. Habitat Alterat	ion and D	evelopment	
Poor to fair (2)  X Poor (1)  4c. Habitat alteration. Score one or double check and average.  None or none apparent (9)  Recovered (6)  X Recovering (3)  ✓ mowing ✓ shrub/sapling removal ✓ grazing ─ learcutting ─ woody debris removal ─ selective cutting ─ farming ─ dredging ─ loxic pollutants ✓ nutrient enrichment	4a. Subs	lone or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1) tat development. Select only one an excellent (7) fery good (6) Good (5) Moderately good (4) fair (3) Foor to fair (2) Foor (1) tat alteration. Score one or double colone or none apparent (9) Recovered (6)	nd assign score.	Check all disturbances o  ☑ mowing ☑ grazing ☐ clearcutting ☐ selective cutting	

Site: PUB-018		Rater(s): OH:jb		Date: 03/2021			
21 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	<b>Vetlands</b>					
	*If the documented raw score for	Metric 5 is 30 points or higher, the	site is automatically co	onsidered a Category 3 wetland.			
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedel [*use higher rank where mixe Superior/enhanced habitat/us	www.score for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland. Where multiple values apply in row, score row as single feature with highest point value. Provide ch selection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). airie (10); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) etl. &/or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] gic feature such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) isolated, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) etc. 1 acre (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) et or floodplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) dapt. in >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) nunity with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] lice state/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) ink where mixed rank or qualifier] [exclude records which are only "historic"] sed habitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) quality): <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
max 20 pts. subtotal	Metric 6. Plant Co	mmunities, Inters <sub>l</sub>	persion, Mic	rotopography			
max 20 pts. Subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 l	to 3 scale.  0 = Absent or <0.1 ha (0.25 acre) contiguous acre  [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a small part of wetland's vegetation and is of moderate quality, or comprises a significant part but is of low quality  2 = Present and either comprises a significant part of wetland's vegetation and is of moderate quality, or comprises a small part and is of high quality  cres (8 ha)  3 = Present and comprises a significant part or more of wetland's vegetation					
	6b. Horizontal (plan view) intersponded to Select only one.		otion of Vegetation Quest diversity &/or domina				
	High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]	High (5)  Moderately high (4) [BR/CM (5)]  Moderately high (4) [BR/CM (5)]  Moderately low (2) [BR/CM (3)]  Moderately low (2) [BR/CM (3)]  Low (1) [BR/CM (2)]  Moderately low (2) [BR/CM (3)]  Low (1) [BR/CM (2)]  Moderately low (2) [BR/CM (3)]  Moderately low (3) [BR/CM (3)]  Moderately low (4) [BR/CM (3)]  Moderately low (5) [BR/CM (3)]  Moderately low (6) [BR/CM (3)]  Moderately low (7) [BR/CM (3)]  Moderately low (8) [BR/CM (3)]  Moderately low (9) [BR/CM (3)]  Moderate					
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5 Moderate 25-75% cover (-1) Sparse 5-25% cover (-1) X Nearly absent <5% cover Absent (1)	e. Mudflat and Ope  0 = Absent < 0.1  1 = Low 0.1 to < (0.1 to 0.5 ac)  (0)	n Water Class Quality ha (0.25 acres) [For BF I ha (0.25 to 2.5 acres) ere)] o <4 ha (2.5 to 9.9 acre	Ate, threatened, or endangered species  R/CM <0.04 ha (0.1 acre)]  [BR/CM 0.04 to <0.2 ha  as) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]  CM 2 ha (5 acres) or more]			
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Werale. socks cm (6 in.) 0 in.) dbh	cland for Estimating D	Degree of Interspersion  Moderate Moderate High			
		2 = Present in m amounts of h	oderate amounts, but nighest quality	more common of marginal quality not of highest quality or in small nunts and of highest quality			
23	GRAND 1 (max 100	30- 59 = Catego	ory 2, good/moderate w	tion, condition, quality** vetland function, condition, quality** function, condition, quality**			

Page 2 of 2

Site: PUB-019	JB-019 Rater(s): JB; OH			Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
2	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average X NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savants), shrubland, young 2nd grossidential, fenced pasture, pasidential, pasi	wetland perimeter (7) ft) around wetland perimeter (2 ft) around wetland perimeter (3 ft) around wetland perimeter (4 wetland perimeter (0) (5 eck and average. (7) (6 math, wildlife area, etc. (7) (7) (8 wth forest (5) (8 conservation tillage, new	(4) er (1)
13	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) cct only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or of (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and D	)evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovered (6) Recovered (6) Recovering (3)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-019	Rater(s):	OH:jb	Date: 03/2021			
23 subtotal previous page						
max 10 pts. subtotal	Special Wetland	S				
	raw score for Metric 5 is 30	points or higher, the site is automaticall	y considered a Category 3 wetland.			
documentation for a Bog, fen, wet Assoc. forest Sensitive geo Vernal pool (E Island wetland Braided chand Gross morph.  Ecological cod Known occurr  [*use higher Superior/enha	each selection (photos, chec prairie (10); acidophilic veg., mo (wetl. &/or adj. upland) incl. >0.2 logic feature such as spring/seep 5); isolated, perched, or slope wed >0.1 acre (0.04 ha) in reservoir nel or floodplain/terrace depressi adapt. in >5 trees >10 in. (25 cmmunity with global rank (Nature rence state/federal threatened/er rank where mixed rank or qualifianced habitat/use: migratory song	e for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.  multiple values apply in row, score row as single feature with highest point value. Provide  ction (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  gracidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3)  adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation]  e such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5)  perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3)  (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5)  blain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3)  5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilled, shallow roots/tip-up, or pneumatophores (3)  th global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier]  federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3)  e mixed rank or qualifier] [exclude records which are only "historic"]  at/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3)  <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 Metric 6. P	Plant Communit	ies, Interspersion, Mi	crotopography			
6a. Wetland vegeta Score all present u Aquatic bed Emergent Shrub Forest Mudflats Open water Moss/lichen	sing 0 to 3 scale.	2 = Present and either comprises a sis of moderate quality, or compri	ntiguous acre			
6b. Horizontal (plar	n view) interspersion.	Narrative Description of Vegetation	Quality			
Select only one.  High (5)  Moderately  Moderate (3	high (4) [BR/CM (5)] 3)[BR/CM (5)] low (2) [BR/CM (3)]	low = Low species diversity &/or dominative species  mod = Native species are dominant of nonnative &/or disturbance tole and species diversity moderate w/o presence of rare, threaten high = A predominance of native species tolerant native speabsent or vir	omponent of the vegetation, although erant native species can also be present, to moderately high, but generally ed or endangered species cies with nonnative sp &/or disturbance tually absent, and high sp diversity and often			
Moderate 25		Mudflat and Open Water Class Qua 0 = Absent <0.1 ha (0.25 acres) [For 1 = Low 0.1 to <1 ha (0.25 to 2.5 acres) (0.1 to 0.5 acre)]	BR/CM <0.04 ha (0.1 acre)] tes) [BR/CM 0.04 to <0.2 ha teres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]			
6d. Microtopograpi Score all present u Vegetated h Coarse woo	,	Hypothetical Wetland for Estimatin  None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts o  2 = Present in moderate amounts, b amounts of highest quality  3 = Present in moderate or greater a	g Degree of Interspersion  Moderate Moderate High  r if more common of marginal quality ut not of highest quality or in small			
0.4	RAND TOTAL nax 100 pts)	0- 29 = Category 1, low wetland ft 30- 59 = Category 2, good/moderat 60-100 = Category 3, superior wetla	e wetland function, condition, quality**			

Site: PUB-020	Rater(s): JB; OH Date: 03/2021		Date: 03/2021	
max 6 pts. subtotal	Metric 1. Wetland	Area (Size) op	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
·	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  X 0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland B	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	O m (164 ft) or more around e 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) around use. Select one or double clor older forest, prairie, savars), shrubland, young 2nd grostdential, fenced pasture, pasidential, pasiden	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter ft) around wetland perimeter ft) ft) around wetland perimeter ft)	(4) er (1)
21	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)	R/CM (5)]  R/CM primary source (5)]  face water (3)  (lake or stream) (5)  ect only one and assign scor  a.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  blogic regime. Score one or (2)  Check all disturbances	Part of wetland/up X Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally satura double check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) aturation. Score one or dbl. check & avg. intly inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and [	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (S) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and average)	Check all disturbances of   ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: PUB-020	Rater(	s): OH:jb	Date: 03/2021			
30 subtotal previous page						
max 10 pts. subtotal	c 5. Special Wetla	nds				
*If the do	cumented raw score for Metric 5 is	30 points or higher, the site is automatically co	onsidered a Category 3 wetland.			
documen  Bo As Se Ve Isl Br Gr Er Sr	tation for each selection (photos, or g, fen, wet prairie (10); acidophilic veg soc. forest (wetl. &/or adj. upland) incl. insitive geologic feature such as spring irrnal pool (5); isolated, perched, or slop and wetland >0.1 acre (0.04 ha) in reseaided channel or floodplain/terrace deposs morph. adapt. in >5 trees >10 in. (2000) ological community with global rank (Nown occurrence state/federal threaten use higher rank where mixed rank or opperior/enhanced habitat/use: migratory	for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland. nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] a such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) alain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) the strees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] at/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 Metri	c 6. Plant Commu	nities, Interspersion, Mici	rotopography			
Score all According Errors Share Moreover	and vegetation communities. present using 0 to 3 scale. quatic bed mergent nrub prest udflats pen water <20 acres (8 ha) poss/lichen. Other	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contig [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a significant and either comprises a significant and is of high quality.	all part of wetland's vegetation and is of gnificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality			
6b. Horiz	ontal (plan view) interspersion.	Narrative Description of Vegetation Qu	uality			
Select on Hi Mi Mi Mi Lo	. , .	low = Low species diversity &/or dominance of nonnative or disturbance tolerar native species  4) [BR/CM (5)] mod = Native species are dominant component of the vegetation, although nonnative &/or disturbance tolerant native species can also be present, and species diversity moderate to moderately high, but generally				
Add or de Ex M Sp X Ne	rage of invasive plants. educt points for coverage. ktensive >75% cover (-5) poderate 25-75% cover (-3) parse 5-25% cover (-1) early absent <5% cover (0) posent (1)	Mudflat and Open Water Class Quality  0 = Absent <0.1 ha (0.25 acres) [For BF  1 = Low 0.1 to <1 ha (0.25 to 2.5 acres)  (0.1 to 0.5 acre)]  2 = Moderate 1 to <4 ha (2.5 to 9.9 acres)  3 = High 4 ha (9.9 acres) or more [BR/C	R/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha es) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]			
6d. Micro Score all Ve	otopography. present using 0 to 3 scale. egetated hummocks/tussocks parse woody debris >15 cm (6 in.) anding dead >25 cm (10 in.) dbh aphibian breeding pools	Hypothetical Wetland for Estimating D  None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if the second process of highest quality  3 = Present in moderate or greater amounts.	Moderate Moderate High  more common of marginal quality not of highest quality or in small			
31	GRAND TOTAL (max 100 pts)	0- 29 = Category 1, low wetland functions 30- 59 = Category 2, good/moderate w 60-100 = Category 3, superior wetland	etland function, condition, quality**			

Site: W-PEM-	Site: W-PEM-01			Date: 03/2021
Max 6 pts. subtotal	Metric 1. Wetland	Area (Size) op (8	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  X <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland I	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	O m (164 ft) or more around be 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double clor older forest, prairie, savans), shrubland, young 2nd grostidential, fenced pasture, pasidential, pasid	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) neck and average. Inah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
5	Metric 3. Hydrolog	gy		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BI X Precipitation (1) [unless E Seasonal/intermittent surform Perennial surface water ac. Maximum water depth. Selector 20.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in X < 0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydrom None or none apparent (1 Recovered (7) Recovering (3)  X Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign scor  a.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  blogic regime. Score one or (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane Regularly inundat Seasonally inundat Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) isaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and [	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovering (2) Poor (1)  4c. Habitat alteration. Score one	only one and assign score.	Check all disturbances of   ☑ mowing ☑ grazing	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal
	Ac. Habitat alteration. Score one None or none apparent (S Recovered (6) Recovering (3) Recent or no recovery (1)	9)	e. clearcutting selective cutting farming toxic pollutants	

Site: W-PEM-1		Rater(s): OH:jb		Date: 03/2021		
12 subtotal previous page						
max 10 pts. subtotal	letric 5. Special V	Vetlands				
*If	the documented raw score for	Metric 5 is 30 points or hi	gher, the site is automatical	y considered a Category 3 wetland.		
	Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perch Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	for Metric 5 is 30 points or higher, the site is automatically considered a Category 3 wetland.  ultiple values apply in row, score row as single feature with highest point value. Provide ion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3)  dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation]  such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5)  erched, or slope wetland (4); headwater wetland [1st order perennial or above] (3)  0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5)  ain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3)  trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3)  adderal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3)  mixed rank or qualifier] [exclude records which are only "historic"]  fuse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3)  acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 M	letric 6. Plant Co	mmunities, In	terspersion, M	crotopography		
	N. Wetland vegetation community core all present using 0 to 3 sca Aquatic bed X Emergent Shrub Forest Mudflats Open water <20 acres (8 h	le. 0 = Absi   For   1 = Pres   mod	erate quality, or comprises a sent and either comprises a moderate quality, or compri	ntiguous acre		
6b.	. Horizontal (plan view) interspe	ersion. Narrative	Description of Vegetation	Quality		
	elect only one.  High (5)  Moderately high (4) [BR/CI  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CN  X Low (1) [BR/CM (2)]  None (0)	M (5)] mod = Lo mad = Na nod =	w species diversity &/or dor tive species ative species are dominant of nnative &/or disturbance toll d species diversity moderate to presence of rare, threaten predominance of native spe erant native sp absent or vii	omponent of the vegetation, although erant native species can also be present, eto moderately high, but generally ed or endangered species cies with nonnative sp &/or disturbance tually absent, and high sp diversity and often		
	c. Coverage of invasive plants. dd or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover ( Absent (1)	9. Mudflat a 0 = Absor 3) 1 = Low (0.1 0) 2 = Mod	and Open Water Class Quant <0.1 ha (0.25 acres) [Fo 0.1 to <1 ha (0.25 to 2.5 acres)] erate 1 to <4 ha (2.5 to 9.9 acres)	Ility BR/CM <0.04 ha (0.1 acre) es) [BR/CM 0.04 to <0.2 ha acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] R/CM 2 ha (5 acres) or more]		
	d. Microtopography. core all present using 0 to 3 sca Vegetated hummocks/tuss Coarse woody debris >15 Standing dead >25 cm (10 Amphibian breeding pools	Hypother lle. locks cm (6 in.) in.) dbh  None  Microtop 0 = Abst 1 = Pres 2 = Pres amo	Low Low  ography Cover Scale ent ient in very small amounts of ent in moderate amounts, bunts of highest quality	g Degree of Interspersion  Moderate Moderate High  r if more common of marginal quality ut not of highest quality or in small		
13	GRAND T (max 100	OIAL 30-59		unction, condition, quality** e wetland function, condition, quality** nd function, condition, quality**		

Site: W-PEM-02		Rater(s): JB; OH		Date: 03/2021	
max 6 pts. subto		Metric 1. Wetland	Area (SIZE) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign  >50 acres (>20.2 ha) (6 pi  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 h  0.3 to <3 acres (0.1 to <1.  X 0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sui	rounding Land	Use
max 14 pts. subto	total	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around a 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <164 ge 10 m to <25 m (32 ft) arouse. Select one or double corrolder forest, prairie, savars), shrubland, young 2nd grostdential, fenced pasture, p	wetland perimeter (7) If the around wetland perimeter (32 ft) around wetland perimeter (32 ft) around wetland perimeter (40) If the around wetland perimeter (5) If the around wetland average. If the around average (7) If the around around around around around average (7) If the around aro	(4) er (1)
12		Metric 3. Hydrolog	ЭУ		
max 30 pts. subte	total	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surformation of the Seasonal/intermittent of the Seasonal Seasonal of the Sea	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) ct only one and assign scor .) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in ologic regime. Score one or 2) Check all disturbance	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally saturat double check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	6	Metric 4. Habitat	Alteration and I	Development	
max 20 pts. subte	total	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovered (6) Recovered (6) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score. or double check and averag	Check all disturbances o  ☑ mowing ☑ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PEM-2	Rater(s): OH:jb	Date: 03/2021				
19 subtotal previous page						
Metric 5. Special \	Metric 5. Special Wetlands					
*If the documented raw score for	Metric 5 is 30 points or higher, the site is auto	omatically considered a Category 3 wetland.				
documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, perci Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with gld Known occurrence state/fedet [*use higher rank where mixe Superior/enhanced habitat/use	lophilic veg., mossy substrate >10 sq.m, sphagnum pland) incl. >0.25 acre (0.1 ha); old growth (10); math as spring/seep, sink, losing/underground stream, hed, or slope wetland (4); headwater wetland [1st oid ha) in reservoir, river, or perennial water >6 ft (2 matherial terrace depressions (floodplain pool, slough, oxbow sex >10 in. (25 cm) dbh: buttress, multitrunk/stool, stipbal rank (NatureServe): G1*(10), G2*(5), G3*(3) [*ural threatened/endangered species (10); other rare sed rank or qualifier] [exclude records which are only	st concurrence, data sources, references, etc). or other moss (5); muck, organic soil layer (3) ature >18 in. (45 cm) dbh (5) [exclude pine plantation] cave, waterfall, rock outcrop/cliff (5) rder perennial or above] (3) n) deep (5) n, meander scar, etc.) (3) iilted, shallow roots/tip-up, or pneumatophores (3) use higher rank where mixed rank or qualifier] species with global rank G1*(10), G2*(5), G3*(3) "historic"] tonbush (4); other fish/wildlife management/designation (3)				
1 Metric 6. Plant Co	mmunities, Interspersio	n, Microtopography				
6a. Wetland vegetation community Score all present using 0 to 3 scar Aquatic bed  X Emergent Shrub Forest Mudflats Open water <20 acres (8 b)	lle. 0 = Absent or <0.1 ha (0.25    For BR/CM <0.04 ha (0)   1 = Present and either comp   moderate quality, or com   2 = Present and either comp   is of moderate quality, or com   and   3 = Present and comprises	acre) contiguous acre				
Moss/lichen. Other  6b. Horizontal (plan view) interspond in the second one.  High (5)  Moderately high (4) [BR/CM (5)]  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CM (2)]  None (0)	M (3)]  low = Low species diversity native species  mod = Native species are dornonnative &/or disturbed and species diversity recommendation with the model of the model.	getation Quality &/or dominance of nonnative or disturbance tolerant minant component of the vegetation, although ance tolerant native species can also be present, moderate to moderately high, but generally threatened or endangered species tive species with nonnative sp &/or disturbance				
6c. Coverage of invasive plants.  Add or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover  Absent (1)	tolerant native sp absorbut not always, the present of the present	ent or virtually absent, and high sp diversity and often esence of rate, threatened, or endangered species				
6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Ale. Socks cm (6 in.) O in.) dbh  None  Low  Microtopography Cover Sca  0 = Absent 1 = Present in very small am  2 = Present in moderate am  amounts of highest qual	nounts or if more common of marginal quality counts, but not of highest quality or in small				
20 GRAND 7	30- 59 = Category 2, good/	vetland function, condition, quality** (moderate wetland function, condition, quality** ior wetland function, condition, quality**				

Metric 1. Wetland Area (size)    Metric 2. Wetland Area (size)	Site: W-PEM-03	Rater(s): JB; OH		Date: 03/2021
Select one size class and assign score.    So sares (20.2 ho 6) parts (10.1 to -20.2 ho (6 pats)		Area (SIZe) ope	en water body (excluding aquation	beds and seasonal mudflats) is >20 acres
2a. Calculate average buffer width. Select only one and assign score. Do not double check.    WIDE. Buffers average 50 m (164 ft) or more around wetland perimeter (7)   MARROW. Buffers average 50 m (164 ft) or more around wetland perimeter (4)   NARROW. Buffers average 25 m to <50 m (32 ft to <52 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 30 m to <50 m (32 ft to <52 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 30 m (164 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 50 m (32 ft to <52 ft) around wetland perimeter (1)   VERY LOW. 20 growth or older forest, prairie, assumansh, widthis area, etc. (7)   LOW. Old field (-10 years), shrubland, young 2nd growth forest (5)   MoDERATELY HIGH. Residential, lenced pasture, park, conservation tillage, new fallow field (3)   High. Urban, industrial, open pasture, row cropping, mining, construction (1)    Metric 3. Hydrology   3a. Sources of water. Score all that apply.   High pH groundwater (5)   Other groundwater (5)   Other groundwater (3) [BR/CM (6)]   Perennial surface water (3) (5)   Seasonal/Internitient surface water (3)   Other groundwater (3) [BR/CM (6)]   Perennial surface water (a) (5)   As of the continual surface water (3) (5)	Select one size class and assign  >50 acres (>20.2 ha) (6 pt  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 h  0.3 to <3 acres (0.1 to <1.  X 0.1 to <0.3 acre (0.04 to <	rs) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	·	size estimate (list):
a. Calculate average buffer width. Select only one and assign score. Do not double check.    WIDLE. Buffers average 50 m (14 ft) or more around wetland perimeter (1)   MEDIUM. Buffers average 50 m to <50 m (82 to <164 ft) around wetland perimeter (4)   NARROW. Buffers average 10 m to <25 m (32 to <164 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 10 m to <25 m (32 to <164 ft) around wetland perimeter (1)   VERY NARROW. Buffers average 10 m (<32 ft) around wetland perimeter (1)   VERY LOW. 2nd growth or older forest, prairie, savannah, wildlife area, etc. (7)   LOW. Old field (-10 years), shrubland, young 2nd growth forest (5)   MODERATELY HIGH. Residential, fenced pasture, park, conservation tillage, new fallow field (3)   Wetric 3. Hydrology   Wetric 3. Sources of water. Score all that apply.   Wetric 3. Sources of water. Score all that apply.   Wetric 3. Sources of water (3)   Wetric 4. Wetric 3. Sources of water (3)   Wetric 4. Wetric 3. Sources of water (3)   Wetric 4. Wetric 4. Wetric 4. Wetric 4. Wetric 5. Sources of water (4)   Wetric 5. Sources of water (5)   Wetric 6. Sources of water (6)   Wetric 6. Sources of water (7)   Wetric 6. Sources of water (8)   Wetric 7. Sources of water (8)   Wetric 7. Sources of water (8)   Wetric 7. Sources of water (8)   Wetric 8. Sources of water (8)   Wetric 8. Sources of water (8)   Wetric 8. Sources of water (8)   Wetric 7. Sources of water (8)   Wetric 8. Sources of water (8)   Wetric 7. Sources of water (8)   Wetric 7	-	Buffers and Sur	rounding Land	Use
a. Sources of water. Score all that apply.    High pH groundwater (3) [BR/CM (5)]   Precipitation (1) [unless BR/CM primary source (5)]   Seasonal/intermittent surface water (3)   Pernial surface water (8)   Part of riparian or upland corridor (1)   Part of water or upland corridor (1)   Part of riparian or upland corridor (1)   Part of riparian or upland corridor (1)   Part of riparian or upland corridor (1)   Part of water or upland corridor (1)   Par	2a. Calculate average buffer widt  WIDE. Buffers average 50  MEDIUM. Buffers average  NARROW. Buffers average  VERY NARROW. Buffers  2b. Intensity of surrounding land of the company of t	o m (164 ft) or more around version to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <8 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savant), shrubland, young 2nd grossidential, fenced pasture, pa	vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter nd wetland perimeter (0) eck and average. nah, wildlife area, etc. (7) wth forest (5) rk, conservation tillage, new	(4) er (1)
3a. Sources of water. Score all that apply.    High pH groundwater (5)   Other groundwater (5)   Descriptation (1) [unless BR/CM (5)]   Part of wetland/upland (e.g., forest), complex (1)   P	12 Metric 3. Hydrolog	ЭУ		
### subtotal  4a. Substrate disturbance. Score one or double check and average.    None or none apparent (4)     Recovered (3)     Recovering (2)     Recent or no recovery (1)     4b. Habitat development. Select only one and assign score.    Excellent (7)     Very good (6)     Good (5)     Moderately good (4)     Fair (3)     Poor to fair (2)     Check all disturbances observed     mowing   shrub/sapling removal	3a. Sources of water. Score all th     High pH groundwater (5)     Other groundwater (3) [BF     X Precipitation (1) [unless B     Seasonal/intermittent surface     Perennial surface water     3c. Maximum water depth. Sele     >0.7 m (27.6 in.) (3)     0.4 to 0.7 m (16 to 27.6 in.)     X<0.4 m (<16 in.) (1) [BR/CN     3e. Modifications to natural hydro     None or none apparent (1)     Recovered (7)     Recovering (3)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) ct only one and assign score  1) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  alogic regime. Score one or decent of the control of the contr	X 100-year floodplai     Between stream/li     Part of wetland/up     Part of riparian or     3d. Duration inundation/se.     Semi- to permane     Regularly inundat     Seasonally inundat     Seasonally satura     Observed     point source (non     filling/grading     road bed/RR track     dredging	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) aturation. Score one or dbl. check & avg. ntly inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
4a. Substrate disturbance. Score one or double check and average.  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select only one and assign score. Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2)  Check all disturbances observed  mowing shrub/sapling removal	6 Metric 4. Habitat A	Alteration and D	evelopment	
4c. Habitat alteration. Score one or double check and average.  None or none apparent (9)  Recovered (6)    Clearcutting   woody debris removal   selective cutting   sedimentation   dredging   dredging	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) Ab. Habitat development. Select (1) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) Ac. Habitat alteration. Score one (2) None or none apparent (9)	only one and assign score.  or double check and average	Check all disturbances of mowing grazing clearcutting selective cutting	☐ shrub/sapling removal ☐ herbaceous/aquatic bed removal ☐ woody debris removal ☐ sedimentation

Site: W-PEM-3		Rater(s): OH:jb	Date: 03/2021				
18							
subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Wetlands					
max vo pie.	*If the documented raw score for	Metric 5 is 30 points or higher, the si	site is automatically considered a Category 3 wetland.				
raw score*	Select all that apply. Where multi documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature sud Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/fedet [*use higher rank where mixt Superior/enhanced habitat/us	ple values apply in row, score row as (photos, checklists, maps, resource tophilic veg., mossy substrate >10 sq.m, supland) incl. >0.25 acre (0.1 ha); old growth as spring/seep, sink, losing/undergrounhed, or slope wetland (4); headwater wetlaha) in reservoir, river, or perennial water (terrace depressions (floodplain pool, sloues >10 in. (25 cm) dbh: buttress, multitrun balar rank (NatureServe): G1*(10), G2*(5), ral threatened/endangered species (10); ced rank or qualifier] [exclude records whice: migratory songbird/waterfowl (5); in-res	c 5 is 30 points or higher, the site is automatically considered a Category 3 wetland. alues apply in row, score row as single feature with highest point value. Provide tos, checklists, maps, resource specialist concurrence, data sources, references, etc). c veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) or slope wetland (4); headwater wetland [1st order perennial or above] (3) n reservoir, river, or perennial water >6 ft (2 m) deep (5) e depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) 0 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) ink (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] satened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) k or qualifier] [exclude records which are only "historic"] ratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) 4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1	Metric 6. Plant Co	mmunities, Interspe	ersion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 in the most of th	ide.  0 = Absent or <0.1    For BR/CM <0.1    For BR/CM <0.1  1 = Present and either moderate qualities of moderate of moderat	ha (0.25 acre) contiguous acre 0.04 ha (0.1 acre)] ther comprises a small part of wetland's vegetation and is of ity, or comprises a significant part but is of low quality ther comprises a significant part of wetland's vegetation and quality, or comprises a small part and is of high quality omprises a significant part or more of wetland's vegetation quality  ion of Vegetation Quality diversity &/or dominance of nonnative or disturbance tolerant es es are dominant component of the vegetation, although or disturbance tolerant native species can also be present, diversity moderate to moderately high, but generally es of rare, threatened or endangered species unce of native species with nonnative sp &/or disturbance ye sp absent or virtually absent, and high sp diversity and often yes, the presence of rate, threatened, or endangered species  Water Class Quality				
	Extensive >75% cover (-5  Moderate 25-75% cover (-5  Sparse 5-25% cover (-1)  Nearly absent <5% cover  Absent (1)	1 = Low 0.1 to <1 h (0.1 to 0.5 acre 2 = Moderate 1 to <	a (0.25 acres) [For BR/CM <0.04 ha (0.1 acre)] ha (0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha e)] <4 ha (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] acres) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetland ale. socks cm (6 in.) 0 in.) dbh  None  Microtopography C 0 = Absent 1 = Present in very 2 = Present in modulamounts of high	And for Estimating Degree of Interspersion  Low Moderate Moderate High  Cover Scale  A small amounts or if more common of marginal quality derate amounts, but not of highest quality or in small				
19	GRAND 1 (max 100	30- 59 = Category	y 1, low wetland function, condition, quality** y 2, good/moderate wetland function, condition, quality** y 3, superior wetland function, condition, quality**				

Site: W-PEM-04		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	2 subtotal	Metric 1. Wetland	Area (Size)	pen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  X 0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland B	<b>Buffers and Su</b>	rrounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	O m (164 ft) or more around e 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to < average <10 m (<32 ft) aro use. Select one or double or or older forest, prairie, sava s), shrubland, young 2nd gr esidential, fenced pasture, p	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. nnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
	12	Metric 3. Hydrolog	ду		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surfix Perennial surface water 3c. Maximum water depth. Seleston 100 on 1	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign sco  a.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in ologic regime. Score one or 2)  Check all disturbance of the control of the con	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane X Regularly inundat Seasonally inundat a.) (2)] Seasonally satura double check and average.  es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)
	6	Metric 4. Habitat	Alteration and	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select  Excellent (7)  Very good (6)  Good (5)  Moderately good (4)  Fair (3)  Poor to fair (2)  Recond (5)  Moderately good (4)  Fair (3)  Poor to fair (2)  None or none apparent (5)	only one and assign score.  or double check and avera	Check all disturbances of   ☑ mowing ☑ grazing ge. □ clearcutting □ selective cutting	☐ shrub/sapling removal ☐ herbaceous/aquatic bed removal ☐ woody debris removal ☐ sedimentation
		Recovered (6) Recovering (3) Recent or no recovery (1)		☐ farming☐ toxic pollutants	☐ dredging ☑ nutrient enrichment

Site: W-PEM-4	Rater(s): OH:jb	Date: 03/2021				
18 subtotal previous page						
Metric 5. Special	Wetlands					
*If the documented raw score fo  select all that apply. Where mult documentation for each selection  Bog, fen, wet prairie (10); ac  Assoc. forest (wetl. &/or adj.  Sensitive geologic feature st  Vernal pool (5); isolated, per Island wetland >0.1 acre (0.1)  Braided channel or floodplain  Gross morph. adapt. in >5 tr  Ecological community with good Known occurrence state/fed  [*use higher rank where milting superior/enhanced habitat/u	r Metric 5 is 30 points or higher, the site tiple values apply in row, score row as s n (photos, checklists, maps, resource sidophilic veg., mossy substrate >10 sq.m, spl upland) incl. >0.25 acre (0.1 ha); old growth uch as spring/seep, sink, losing/underground ched, or slope wetland (4); headwater wetland (4) in reservoir, river, or perennial water > n/terrace depressions (floodplain pool, sloughees >10 in. (25 cm) dbh: buttress, multitrunk/lobal rank (NatureServe): G1*(10), G2*(5), othe ked rank or qualifier] [exclude records which is se: migratory songbird/waterfowl (5); in-reser	is 30 points or higher, the site is automatically considered a Category 3 wetland.  see apply in row, score row as single feature with highest point value. Provide see, checklists, maps, resource specialist concurrence, data sources, references, etc).  seg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3)  cl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation]  ng/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5)  ope wetland (4); headwater wetland [1st order perennial or above] (3)  servoir, river, or perennial water >6 ft (2 m) deep (5)  servoir, river, or perennial water >6 ft (2 m) deep (5)  servoir, dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3)  (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier]  sened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3)  requalifier] [exclude records which are only "historic"]  ory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3)  has AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
Metric 6. Plant Co  aux 20 pts.  Metric 6. Plant Co  aux 20 pts.  6a. Wetland vegetation commun Score all present using 0 to 3 so  Aquatic bed  Emergent Shrub Forest Mudflats Open water <20 acres (8 Moss/lichen. Other  High (5) Moderately high (4) [BR/CM (5)] Moderately low (2) [BR/CM (2)] None (0)  6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-1) Moderate 25-75% cover  Moderate 25-75% cover Sparse 5-25% cover (-1) Nearly absent <5% cove Absent (1)  6d. Microtopography. Score all present using 0 to 3 s Vegetated hummocks/tu Coarse woody debris >1 Standing dead >25 cm (-1) Amphibian breeding poo	Vegetation Communicate Nationale.  Vegetation Communicate Nationale.  Vegetation Communicate Nationale.  O = Absent or <0.1 has present and either moderate quality. The present and either is of moderate quality. The present and communicate of high quarties of high quarties pecies of the native species of th	a (0.25 acre) contiguous acre 4 ha (0.1 acre)] er comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality er comprises a significant part of wetland's vegetation and ality, or comprises a small part and is of high quality prises a significant part or more of wetland's vegetation ality  of Vegetation Quality pressity &/or dominance of nonnative or disturbance tolerant are dominant component of the vegetation, although disturbance tolerant native species can also be present, pressity moderate to moderately high, but generally for are, threatened or endangered species e of native species with nonnative sp &/or disturbance esp absent or virtually absent, and high sp diversity and often the presence of rate, threatened, or endangered species  ater Class Quality 0.25 acres) [For BR/CM < 0.04 ha (0.1 acre)] (0.25 to 2.5 acres) [BR/CM < 0.04 to < 0.2 ha  ha (2.5 to 9.9 acres) [BR/CM 0.04 to < 0.2 ha  ha (2.5 to 9.9 acres) [BR/CM 0.2 to < 0.2 ha (0.5 to 5 acre)] res) or more [BR/CM 2 ha (5 acres) or more]  If for Estimating Degree of Interspersion  Moderate  mall amounts or if more common of marginal quality ate amounts, but not of highest quality or in small				
19 GRAND (max 100	707AL 0- 29 = Category 1 30- 59 = Category 2	ate or greater amounts and of highest quality  , low wetland function, condition, quality** , good/moderate wetland function, condition, quality** , superior wetland function, condition, quality**				

Site: W-PEM-05		Rater(s): JB; OH		Date: 03/2021	
max 6 pts. su	2 ubtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
mak o poi.		Select one size class and assign  >50 acres (>20.2 ha) (6 pi  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 h  X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	is) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. su	ubtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around 2 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double concluder forest, prairie, savance), shrubland, young 2nd grosidential, fenced pasture, pasidential, pasiden	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter ft) around wetland perimeter ft) ft) around wetland perimeter ft)	(4) er (1)
	11	Metric 3. Hydrolog	ЭУ		
max 30 pts. su	ubtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF X Precipitation (1) [unless B X Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selector > 0.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) X<0.4 m (<16 in.) (1) [BR/Cl 3e. Modifications to natural hydror None or none apparent (1) Recovered (7)  Recovering (3)  X Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) ct only one and assign scor .) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in. blogic regime. Score one or o	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane Regularly inundat Seasonally inundat O(2)] X Seasonally saturat Couble check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	6	Metric 4. Habitat	Alteration and D	Development	
max 20 pts. su	ubtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Recovered (6) Recovered (6) Recovered (6) Recovering (3) Recent or no recovery (1)	) only one and assign score. or double check and averag	Check all disturbances o	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PEM-05	Rater(s): OH:jb	Date: 03/2021			
20 subtotal previous page					
Metric 5. Speci	al Wetlands				
*If the documented raw sco	e for Metric 5 is 30 points or higher, the site is a	automatically considered a Category 3 wetland.			
documentation for each sele  Bog, fen, wet prairie (10  Assoc. forest (wetl. &/o  Sensitive geologic featt  Vernal pool (5); isolated  Island wetland >0.1 acr  Braided channel or floo  Gross morph. adapt. in  Ecological community v  Known occurrence statt  [*use higher rank whe]  Superior/enhanced hab	nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] e such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) dain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] it/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 Metric 6. Plant	Communities, Interspersi	on, Microtopography			
6a. Wetland vegetation com Score all present using 0 to Aquatic bed X Emergent Shrub Forest Mudflats Open water <20 acre Moss/lichen. Other	3 scale. 0 = Absent or <0.1 ha (0.    For BR/CM <0.04 ha	25 acre) contiguous acre			
6b. Horizontal (plan view) ir	terspersion. Narrative Description of	Vegetation Quality			
Select only one.  High (5)  Moderately high (4)  Moderate (3)[BR/CN  Moderately low (2) [I  X Low (1) [BR/CM (2)]  None (0)	low = Low species diversity &/or dominance of nonnative or disturbance native species  BR/CM (5)] mod = Native species are dominant component of the vegetation, althout nonnative &/or disturbance tolerant native species can also be pro-				
6c. Coverage of invasive pla Add or deduct points for cov Extensive >75% cov Moderate 25-75% cov Sparse 5-25% cover Nearly absent <5% of Absent (1)	Mudflat and Open Water       er (-5)     0 = Absent < 0.1 ha (0.25)       ver (-3)     1 = Low 0.1 to <1 ha (0.2)       (-1)     (0.1 to 0.5 acre)       over (0)     2 = Moderate 1 to <4 has	Class Quality acres) [For BR/CM <0.04 ha (0.1 acre)] 5 to 2.5 acres) [BR/CM 0.04 to <0.2 ha  (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] or more [BR/CM 2 ha (5 acres) or more]			
6d. Microtopography. Score all present using 0 to Vegetated hummock Coarse woody debri Standing dead >25 o	Hypothetical Wetland for  3 scale. s/tussocks s > 15 cm (6 in.) m (10 in.) dbh pools  None  Low  Microtopography Cover 0 = Absent 1 = Present in very small 2 = Present in moderate amounts of highest qu	Estimating Degree of Interspersion  Low Moderate Moderate High  Scale  amounts or if more common of marginal quality amounts, but not of highest quality or in small			
l 21 II -	30- 59 = Category 2, god	v wetland function, condition, quality** od/moderate wetland function, condition, quality** perior wetland function, condition, quality**			

Site: W-PEM-06		Rater(s): JB; OH		Date: 03/2021	
max 6 pts. sul	1 lbtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign	is) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. sul	ibtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around 25 m to <50 m (82 to <164 pe 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arouse. Select one or double clar older forest, prairie, savar (), shrubland, young 2nd grosidential, fenced pasture, pasidential, pasidenti	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) heck and average. hah, wildlife area, etc. (7) howth forest (5) ark, conservation tillage, new	(4) er (1)
	12	Metric 3. Hydrolog	ЭУ		
max 30 pts. sul	ibtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF   Precipitation (1) [unless B   Seasonal/intermittent surf.	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) ct only one and assign scor .) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in allogic regime. Score one or allogic regime.	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally saturat double check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	3	Metric 4. Habitat A	Alteration and [	Development	
max 20 pts. sul	ıbtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (1) Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) X Recent or no recovery (1)	) only one and assign score. or double check and averag	Check all disturbances of   ☑ mowing ☐ grazing	observed

Site:W-PEM-06	Rater(s): OH:jb	Date: 03/2021			
17 subtotal previous page					
Metric 5. Specia	l Wetlands				
*If the documented raw score	for Metric 5 is 30 points or higher, the site is auto	omatically considered a Category 3 wetland.			
documentation for each select Bog, fen, wet prairie (10); Assoc. forest (wetl. &/or a Sensitive geologic feature Vernal pool (5); isolated, Island wetland >0.1 acre Braided channel or flood; Gross morph. adapt. in >0 Ecological community wit Known occurrence state/fi [*use higher rank where Superior/enhanced habita	nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) lain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) is trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] truse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
<sup>1</sup> Metric 6. Plant 0	Communities, Interspersion	n, Microtopography			
6a. Wetland vegetation comm Score all present using 0 to 3 Aquatic bed X Emergent Shrub Forest Mudflats Open water <20 acres Moss/lichen. Other	scale. 0 = Absent or <0.1 ha (0.25    For BR/CM <0.04 ha (0.01 ha (0.025 ha (0.01 ha	acre) contiguous acre			
6b. Horizontal (plan view) inte	rspersion. Narrative Description of Ve	getation Quality			
Select only one.  High (5)  Moderately high (4) [B  Moderate (3)[BR/CM (  Moderately low (2) [BF  X Low (1) [BR/CM (2)]  None (0)	low = Low species diversity &/or dominance of nonnative or disturbance to native species  R/CM (5)] mod = Native species are dominant component of the vegetation, although nonnative &/or disturbance tolerant native species can also be presented.				
6c. Coverage of invasive plar Add or deduct points for cove  Extensive >75% cover  Moderate 25-75% cov  Sparse 5-25% cover ( Nearly absent <5% co Absent (1)	ts. rage. (-5)  er (-3) 1)  ver (0)  Mudflat and Open Water Cla 0 = Absent < 0.1 ha (0.25 acr 1 = Low 0.1 to <1 ha (0.25 to (0.1 to 0.5 acre)] 2 = Moderate 1 to <4 ha (2.5	ass Quality res) [For BR/CM <0.04 ha (0.1 acre)] o 2.5 acres) [BR/CM 0.04 to <0.2 ha to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] more [BR/CM 2 ha (5 acres) or more]			
6d. Microtopography. Score all present using 0 to 3 Vegetated hummocks. Coarse woody debris: Standing dead >25 cm Amphibian breeding p	Hypothetical Wetland for Est tussocks >15 cm (6 in.) (10 in.) dbh pols  Microtopography Cover Sca 0 = Absent 1 = Present in very small am 2 = Present in moderate amamounts of highest quali	Low Moderate Moderate High  ale  nounts or if more common of marginal quality ounts, but not of highest quality or in small			
18 GRANE (max 10	30- 59 = Category 2, good/i	etland function, condition, quality** moderate wetland function, condition, quality** ior wetland function, condition, quality**			

Site:	W-PEM	-07	Rater(s): JB; OH		Date: 03/2021
max 6 pts.	2 subtotal	Metric 1. Wetland	Area (Size) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
max 14 pts	1 subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	th. Select only one and assign (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <164 average <10 m (<32 ft) arouse. Select one or double corrolder forest, prairie, savars), shrubland, young 2nd grostidential, fenced pasture, p	gn score. Do not double check wetland perimeter (7)  If the around wetland perimeter (82 ft) around wetland perimeter (0)  wetland perimeter (0)  heck and average.  nah, wildlife area, etc. (7)  with forest (5)  ark, conservation tillage, new	k. (4) er (1)
max 30 pts	5 subtotal	Metric 3. Hydrolog  3a. Sources of water. Score all the High pH groundwater (5) Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Selection 90.7 m (27.6 in.) (3) 0.4 to 0.7 m (16 to 27.6 in. X<0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydrom None or none apparent (1) Recovered (7) Recovering (3) X Recent or no recovery (1)	nat apply.  R/CM (5)]  R/CM primary source (5)] face water (3) (lake or stream) (5) foct only one and assign score.)  1.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in blogic regime. Score one or 2)  Check all disturbance  ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s e. Semi- to permane Regularly inundat X Seasonally inundat O(2) Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ated in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	6	Metric 4. Habitat		Development	
max 20 pts	subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and averag	Check all disturbances o  ☑ mowing ☐ grazing	observed

Site: W-PEM-07		Rater(s): OH:jb	Date: 03/2021			
14 subtotal previous page						
max 10 pts. subtotal	letric 5. Special V	Vetlands				
*If 1	the documented raw score for	Metric 5 is 30 points or higher, the sit	e is automatically considered a Category 3 wetland.			
	cumentation for each selection  Bog, fen, wet prairie (10); acid  Assoc. forest (wetl. &/or adj. u  Sensitive geologic feature suc  Vernal pool (5); isolated, perch Island wetland >0.1 acre (0.04  Braided channel or floodplain/  Gross morph. adapt. in >5 tree  Ecological community with glo  Known occurrence state/feder  [*use higher rank where mixe  Superior/enhanced habitat/use	ultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) dj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) 0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) lain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) is trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) in global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] truse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 M	letric 6. Plant Co	mmunities, Interspe	rsion, Microtopography			
	. Wetland vegetation communit ore all present using 0 to 3 sca Aquatic bed X Emergent Shrub Forest Mudflats Open water <20 acres (8 h	le. 0 = Absent or <0.1 h    For BR/CM <0.0   1 = Present and eith   moderate quality   2 = Present and eith   is of moderate q	na (0.25 acre) contiguous acre  04 ha (0.1 acre)]  ther comprises a small part of wetland's vegetation and is of the comprises a significant part but is of low quality there comprises a significant part of wetland's vegetation and the comprises a small part and is of high quality the prises a significant part or more of wetland's vegetation			
6b.	. Horizontal (plan view) interspe	ersion. Narrative Descriptio	on of Vegetation Quality			
	High (5) High (5) Moderately high (4) [BR/C Moderately low (2) [BR/CM Moderately low (2) [BR/CM X Low (1) [BR/CM (2)] None (0)	low = Low species diversity &/or dominance of nonnative or disturbance native species  [BR/CM (5)] mod = Native species are dominant component of the vegetation, althoug nonnative &/or disturbance tolerant native species can also be present and species diversity moderate to moderately high, but generally				
	Coverage of invasive plants.  d or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover (Absent (1)	Mudflat and Open W 0 = Absent <0.1 ha ( 3) 1 = Low 0.1 to <1 ha ( 0.1 to 0.5 acre) 2 = Moderate 1 to <4	(0.25 acres) [For BR/CM <0.04 ha (0.1 acre)] a (0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha			
	d. Microtopography.  core all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetlan  ale. socks cm (6 in.) in.) dbh  None  Low  Microtopography Co 0 = Absent 1 = Present in very s 2 = Present in mode amounts of higher	Low Moderate Moderate High  were Scale  Small amounts or if more common of marginal quality erate amounts, but not of highest quality or in small			
15	GRAND T (max 100	30- 59 = Category 2	1, low wetland function, condition, quality** 2, good/moderate wetland function, condition, quality** 3, superior wetland function, condition, quality**			

Site: W-PEM-08		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	3 subtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign  >50 acres (>20.2 ha) (6 pi  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  X 3 to <10 acres (1.2 to <4 h  0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double conduction of the conduction of th	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) neck and average. Inah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
	7	Metric 3. Hydrolog	ЭУ		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Seleston 20.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in X<0.4 m (<16 in.) (1) [BR/Cl Se. Modifications to natural hydrom None or none apparent (1) Recovered (7)  Recovering (3)  X Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) ct only one and assign scor  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane X Regularly inundat Seasonally inundat O(2)] Seasonally saturations and average. Sobserved point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) auturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)]  stormwater)
	6	Metric 4. Habitat A	Alteration and <b>D</b>	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select (1) Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score. or double check and averag	Check all disturbances of   ☑ mowing ☑ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PEM-08	Rater(s): O	H:jb		Date: 03/2	021	
17 subtotal previous page						
Metric 5. Spec	ial Wetlands					
*If the documented raw so	ore for Metric 5 is 30 po	ints or higher, the site is a	utomatically co	onsidered a Cate	egory 3 wetland	d.
documentation for each so Bog, fen, wet prairie ( Assoc. forest (wetl. & Sensitive geologic fee Vernal pool (5); isolat Island wetland >0.1 a Braided channel or fl Gross morph. adapt. Ecological community Known occurrence st [*use higher rank wh Superior/enhanced he	election (photos, checklis 10); acidophilic veg., mossy or adj. upland) incl. >0.25 a ture such as spring/seep, s ed, perched, or slope wetlar cre (0.04 ha) in reservoir, riv codplain/terrace depressions n >5 trees >10 in. (25 cm) o with global rank (NatureSe ate/federal threatened/enda ere mixed rank or qualifier] sbitat/use: migratory songbi	nultiple values apply in row, score row as single feature with highest point value. Provide tion (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] e such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) alain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) 5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) h global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ederal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) mixed rank or qualifier] [exclude records which are only "historic"] tituse: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)				
1 Metric 6. Plan	Communitie	es, Interspersi	on, Micr	otopogra	aphy	
6a. Wetland vegetation of Score all present using 0 Aquatic bed  X Emergent Shrub Forest Mudflats Open water <20 at Moss/lichen. Other	o 3 scale. 0 1 2 res (8 ha) 3	/egetation Community Co D = Absent or <0.1 ha (0.2   For BR/CM <0.04 ha D = Present and either cormoderate quality, or collist of moderate quality. D = Present and comprise and is of high quality	5 acre) contiguous (0.1 acre)] mprises a smalemprises a signification or comprises	Il part of wetland nificant part but ificant part of we a small part and	is of low qualitetland's vegeta I is of high qua	ty ition and ility
6b. Horizontal (plan view)	interspersion.	Varrative Description of V	/egetation Qu	alitv		
Select only one.  High (5)  Moderately high (4)  Moderate (3)[BR/C  Moderately low (2)  X Low (1) [BR/CM (2)  None (0)	[BR/CM (5)] n M (5)] [BR/CM (3)]	low = Low species diversity &/or dominance of nonnative or disturbance t native species  [R/CM (5)] mod = Native species are dominant component of the vegetation, although nonnative &/or disturbance tolerant native species can also be pres				ugh resent, / ance / and often
6c. Coverage of invasive   Add or deduct points for c  Extensive >75% co  Moderate 25-75%  Sparse 5-25% cov  Nearly absent <5%  Absent (1)	over age. Nover (-5) 00 00 00 00 00 00 00 00 00 00 00 00 00	but not always, the public form of the following form of the follo	Class Quality acres) [For BR to 2.5 acres)	/CM <0.04 ha (0 [BR/CM 0.04 to s) [BR/CM 0.2 to	0.1 acre)] <0.2 ha o <02 ha (0.5 to	
6d. Microtopography. Score all present using 0 Vegetated hummo Coarse woody deb Standing dead >25 Amphibian breedir	to 3 scale. cks/tussocks ris >15 cm (6 in.) cm (10 in.) dbh g pools	None Low  Microtopography Cover S  Present in very small a  Present in moderate a amounts of highest qu  B = Present in moderate o	Estimating Do	Moderate  nore common of ot of highest quarte	Moderate  marginal qual ality or in small	
1 10 11 -	ND TOTAL 100 pts)	0- 29 = Category 1, low 30- 59 = Category 2, goo 60-100 = Category 3, sup	d/moderate we	etland function,	condition, quali	ity**

Site: W-PEM-10		Rater(s): JB; OH		Date: 03/2021
2 max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) ope	en water body (excluding aquation	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  X 0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
1	Metric 2. Upland I	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers averag NARROW. Buffers averag VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	O m (164 ft) or more around ve 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <8 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savans), shrubland, young 2nd groesidential, fenced pasture, pastidential, fenced pastidential,	vetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) seck and average. hah, wildlife area, etc. (7) wth forest (5) irk, conservation tillage, new	(4) er (1)
12	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [Blook Precipitation (1) [unless Extended Seasonal/intermittent surfold Perennial surface water ac. Maximum water depth. Selector Season Perennial surface water sc. Maximum water depth. Selector No.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) (1) [BR/C]  3e. Modifications to natural hydromaphology (1) [BR/C]  Recovered (7)  X Recovering (3)  Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  face water (3)  (lake or stream) (5)  ect only one and assign score  1.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  blogic regime. Score one or of (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/s Semi- to permane X Regularly inundat Seasonally inundat (2)] Seasonally satura louble check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
6	Metric 4. Habitat	Alteration and D	evelopment	
max 20 pts. subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) X Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) X Poor (1) 4c. Habitat alteration. Score one None or none apparent (5) Recovered (6) X Recovering (3) Recent or no recovery (1)	only one and assign score.  or double check and average)	Check all disturbances o  ☑ mowing ☑ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PEM-10	Rater(s):	OH:jb	Date: 03/2021
21 subtotal previous page			
max 10 pts. subtotal Metric	5. Special Wetland	ds	
	mented raw score for Metric 5 is 30	o points or higher, the site is automatically co	onsidered a Category 3 wetland.
documentat  Bog, Assor Sens Verna Island Gross Ecolo Know  [*usr	ion for each selection (photos, che fen, wet prairie (10); acidophilic veg., mo. forest (wetl. &/or adj. upland) incl. >0. tive geologic feature such as spring/seal pool (5); isolated, perched, or slope will wetland >0.1 acre (0.04 ha) in reservoed channel or floodplain/terrace depress morph. adapt. in >5 trees >10 in. (25 cigical community with global rank (Naturn occurrence state/federal threatened/eachigher rank where mixed rank or qualitior/enhanced habitat/use: migratory so	ply in row, score row as single feature with he cklists, maps, resource specialist concurrences y substrate >10 sq.m, sphagnum or other most 25 acre (0.1 ha); old growth (10); mature >18 in. (4 ep, sink, losing/underground stream, cave, waterfaretland (4); headwater wetland [1st order perennial ir, river, or perennial water >6 ft (2 m) deep (5) sions (floodplain pool, slough, oxbow, meander sciently of the buttress, multitrunk/stool, stilted, shallow reserve): G1*(10), G2*(5), G3*(3) [*use higher rander and angered species (10); other rare species with gifter] [exclude records which are only "historic"] ingbird/waterfowl (5); in-reservoir buttonbush (4); of the property of the	ce, data sources, references, etc). s (5); muck, organic soil layer (3) 45 cm) dbh (5) [exclude pine plantation] II, rock outcrop/cliff (5) or above] (3) ar, etc.) (3) roots/tip-up, or pneumatophores (3) k where mixed rank or qualifier] lobal rank G1*(10), G2*(5), G3*(3) ther fish/wildlife management/designation (3)
	6. Plant Communi	ties, Interspersion, Micr	otopography
Score all pro Aqua X Eme Shru Fore Mudi	b st	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contig	Il part of wetland's vegetation and is of inificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality
6b. Horizon Select only High Mod Mod	al (plan view) interspersion. one. (5) erately high (4) [BR/CM (5)] erate (3)[BR/CM (5)] erately low (2) [BR/CM (3)] (1) [BR/CM (2)]	Narrative Description of Vegetation Quality low = Low species diversity &/or dominative species  mod = Native species are dominant companitive &/or disturbance tolerant and species diversity moderate to w/o presence of rare, threatened of high = A predominance of native species	conent of the vegetation, although at native species can also be present, moderately high, but generally or endangered species
Add or dedu Exte Mod X Spar Near	e of invasive plants. ct points for coverage. nsive >75% cover (-5) erate 25-75% cover (-3) se 5-25% cover (-1) ly absent <5% cover (0) ent (1)	•	/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha s) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]
☐ Vege ☐ Coal ☐ Stan	pography. esent using 0 to 3 scale. etated hummocks/tussocks se woody debris >15 cm (6 in.) ding dead >25 cm (10 in.) dbh hibian breeding pools	Hypothetical Wetland for Estimating Down None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if no	Moderate Moderate High
	GRAND TOTAL	2 = Present in moderate amounts, but no amounts of highest quality 3 = Present in moderate or greater amounts 0-29 = Category 1, low wetland funct 30-59 = Category 2, good/moderate we	ot of highest quality or in small  unts and of highest quality  ion, condition, quality** etland function, condition, quality**
22	(max 100 nts)	60-100 = Category 3, superior wetland f	

Site: W-PFO-01		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	1 subtotal	Metric 1. Wetland		pen water body (excluding aquati-	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign   >50 acres (>20.2 ha) (6 p   25 to <50 acres (10.1 to <   10 to <25 acres (4 to <10.   3 to <10 acres (1.2 to <4 l   0.3 to <3 acres (0.1 to <1.   X 0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	3	Metric 2. Upland E	Buffers and Su	rrounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average X NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	o m (164 ft) or more around 2 25 m to <50 m (82 to <16 3 9 9 10 m to <25 m (32 ft to <16 4 average <10 m (<32 ft) around 5 use. Select one or double or 6 or older forest, prairie, sava 6), shrubland, young 2nd gr	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. linnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
	4	Metric 3. Hydrolog	ЭУ		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Seleston 100 pc. 1	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) act only one and assign sco act.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in blogic regime. Score one or	Part of wetland/up Part of riparian or 3d. Duration inundation/s ore. Semi- to permane Regularly inundat Seasonally inundat Seasonally satura of double check and average.  es observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] sted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	6	Metric 4. Habitat	Alteration and	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select  Excellent (7)  Very good (6)  Good (5)  Moderately good (4)  Fair (3)  Poor to fair (2)	)	Check all disturbances o  ☑ mowing	☐ shrub/sapling removal
		X Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) X Recovering (3) Recent or no recovery (1)	)	ge.	

Site: W-PFO-(	)1	Rater(s): OH:jb	Date: 03/2021				
14 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special \	Wetlands					
	*If the documented raw score for	Metric 5 is 30 points or higher, the site is	automatically considered a Category 3 wetland.				
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. t. Sensitive geologic feature suc Vernal pool (5); isolated, perc Island wetland >0.1 acre (0.04 Braided channel or floodplain, Gross morph. adapt. in >5 tre Ecological community with gld Known occurrence state/fedel [*use higher rank where mix Superior/enhanced habitat/us	e multiple values apply in row, score row as single feature with highest point value. Provide lection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  0); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] ture such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) dd, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) are (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) adplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) an >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] te/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) ere mixed rank or qualifier] [exclude records which are only "historic"] bitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) v) : <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
3	Metric 6. Plant Co	mmunities, Interspers	sion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communi Score all present using 0 to 3 sca Aquatic bed Emergent Shrub X Forest Mudflats Open water <20 acres (8 Moss/lichen. Other	lle. 0 = Absent or <0.1 ha   [For BR/CM <0.04] 1 = Present and either moderate quality, conduction   2 = Present and either is of moderate quality.	0.25 acre) contiguous acre na (0.1 acre)] comprises a small part of wetland's vegetation and is of r comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and ity, or comprises a small part and is of high quality ises a significant part or more of wetland's vegetation				
	6b. Horizontal (plan view) intersp	ersion. Narrative Description	of Vegetation Quality				
	Select only one.  High (5)  Moderately high (4) [BR/C  Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/CI  Low (1) [BR/CM (2)]  None (0)	low = Low species diversity &/or dominance of nonnative or disturbance native species  #I [BR/CM (5)] mod = Native species are dominant component of the vegetation, althout nonnative &/or disturbance tolerant native species can also be proposed in the pr					
	6c. Coverage of invasive plants.  Add or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover  Absent (1)	Mudflat and Open Wate  0 = Absent < 0.1 ha (0.3)  1 = Low 0.1 to <1 ha (0.4)  (0.1 to 0.5 acre)  2 = Moderate 1 to <4 has to <	er Class Quality 25 acres) [For BR/CM <0.04 ha (0.1 acre)] 25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha a (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] 3) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sc:  Vegetated hummocks/tus:  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	Hypothetical Wetland for socks cm (6 in.)  O in.) dbh  None Low  Microtopography Cove  0 = Absent  1 = Present in very sm.  2 = Present in moderat amounts of highest	Low Moderate Moderate High  Tr Scale  all amounts or if more common of marginal quality amounts, but not of highest quality or in small				
17	GRAND 1 (max 100	30-59 = Category 2. o	ow wetland function, condition, quality** lood/moderate wetland function, condition, quality** uperior wetland function, condition, quality**				

Site: W-PFO-02		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	2 subtotal	Metric 1. Wetland	Area (SIZe) op	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
mat o po		Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10.  3 to <10 acres (1.2 to <4 l  X 0.3 to <3 acres (0.1 to <1.  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) 20.2 ha) (5) [BR/CM (6)] 1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] 2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	1	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average NARROW. Buffers average X VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double conduction of the conduction of th	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) neck and average. Inah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
	8	Metric 3. Hydrolog	ЭУ		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BF   Precipitation (1) [unless B   X Seasonal/intermittent surform   Perennial surface water   3c. Maximum water depth. Selector   >0.7 m (27.6 in.) (3)   0.4 to 0.7 m (16 to 27.6 in.)   X<0.4 m (<16 in.) (1) [BR/C] 3e. Modifications to natural hydrom   None or none apparent (1   Recovered (7)   Recovering (3)   X   Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5) ct only one and assign scor  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.) clogic regime. Score one or (2)  Check all disturbances	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane Regularly inundat X Seasonally inundat Ouble check and average. s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) auturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)]  stormwater)
	6	Metric 4. Habitat A	Alteration and D	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recent or no recovery (1) Recent or no recovery (1) Ab. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Poor to fair (2) Poor (1) Ac. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	only one and assign score. or double check and averag	Check all disturbances of   ☑ mowing ☐ grazing	observed shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PFO-	04	Rater(s): OH:jb	Date: 03/2021
17 subtotal previous page			
max 10 pts. subtotal	Metric 5. Special V	Vetlands	
·	*If the documented raw score for	Metric 5 is 30 points or higher, the site is	automatically considered a Category 3 wetland.
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u Sensitive geologic feature suc Vernal pool (5); isolated, percl Island wetland >0.1 acre (0.04 Braided channel or floodplain/ Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder [*use higher rank where mixe Superior/enhanced habitat/use	(photos, checklists, maps, resource speophilic veg., mossy substrate >10 sq.m, sphagpland) incl. >0.25 acre (0.1 ha); old growth (10 h as spring/seep, sink, losing/underground strued, or slope wetland (4); headwater wetland [4 ha) in reservoir, river, or perennial water >6 fterrace depressions (floodplain pool, slough, cas >10 in. (25 cm) dbh: buttress, multitrunk/stobal rank (NatureServe): G1*(10), G2*(5), G3*(al threatened/endangered species (10); other are rank or qualifier] [exclude records which are exemigratory songbird/waterfowl (5); in-reservo	1st order perennial or above] (3) (2 m) deep (5) xbow, meander scar, etc.) (3) ol, stilted, shallow roots/tip-up, or pneumatophores (3) 3) [*use higher rank where mixed rank or qualifier] rare species with global rank G1*(10), G2*(5), G3*(3)
3	Metric 6. Plant Co	mmunities, Interspers	ion, Microtopography
max 20 pts. subtotal	6a. Wetland vegetation communit Score all present using 0 to 3 sca Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 h	0 = Absent or <0.1 ha (	2.25 acre) contiguous acre (a (0.1 acre)]  comprises a small part of wetland's vegetation and is of comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and ty, or comprises a small part and is of high quality sees a significant part or more of wetland's vegetation
	Moss/lichen. Other  6b. Horizontal (plan view) intersperson Select only one.  High (5)  Moderately high (4) [BR/C Moderate (3)[BR/CM (5)]	Iow = Low species divernative species  M (5)] mod = Native species are nonnative &/or dis	f Vegetation Quality sity &/or dominance of nonnative or disturbance tolerant e dominant component of the vegetation, although turbance tolerant native species can also be present,
	Moderately low (2) [BR/CM X Low (1) [BR/CM (2)] None (0)	w/o presence of rachingh = A predominance of tolerant native sp	sity moderate to moderately high, but generally are, threatened or endangered species  of native species with nonnative sp &/or disturbance absent or virtually absent, and high sp diversity and often be presence of rate, threatened, or endangered species
	6c. Coverage of invasive plants.  Add or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover (-1)  Absent (1)	Mudflat and Open Water  0 = Absent <0.1 ha (0.2  1 = Low 0.1 to <1 ha (0  (0.1 to 0.5 acre)]  2 = Moderate 1 to <4 ha	
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  Amphibian breeding pools	ale. socks cm (6 in.)	or Estimating Degree of Interspersion
		Microtopography Cove 0 = Absent 1 = Present in very sma 2 = Present in moderate amounts of highest	Il amounts or if more common of marginal quality amounts, but not of highest quality or in small
21	GRAND T	30- 59 = Category 2, g	ow wetland function, condition, quality** cod/moderate wetland function, condition, quality** uperior wetland function, condition, quality**

Site: W-PFO-03		Rater(s): JB; OH		Date: 03/2021
Max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) opi	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4  0.3 to <3 acres (0.1 to <1  0.1 to <0.3 acre (0.04 to  X <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] ha) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
11	Metric 2. Upland I	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers 2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savans), shrubland, young 2nd gro	wetland perimeter (7) ft) around wetland perimeter ft) around wetland perimeter and wetland perimeter (0) neck and average. nah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
10	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all the High pH groundwater (5)  X Other groundwater (3) [BI Precipitation (1) [unless Eseasonal/intermittent surform Perennial surface water ac. Maximum water depth. Seleston 20.4 to 0.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) (3)  None or none apparent (1)  Recovered (7)  Recovered (7)  Recovering (3)  X Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign score  1.) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  Clogic regime. Score one or complete the stream of the	Part of wetland/up Part of riparian or 3d. Duration inundation/se.  X Semi- to permane Regularly inundat Seasonally inundat Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) upland corridor (1) isaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] ted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
41	Metric 4. Habitat	Alteration and E	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  X None or none apparent (4 Recovered (3) Recovering (2) Recent or no recovery (1) 4b. Habitat development. Select Excellent (7) Very good (6) Good (5)	<b>(</b> )	verage.	
	Moderately good (4)  X Fair (3) Poor to fair (2) Poor (1)  4c. Habitat alteration. Score one X None or none apparent (5 Recovered (6) Recovering (3) Recent or no recovery (1)	9)	Check all disturbances of mowing grazing clearcutting selective cutting farming toxic pollutants	bbserved shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PFO-04	4	Rater(s): OH	:jb			Date: 03/2	2021	
subtotal previous page								
max 10 pts. subtotal	Metric 5. Special V	Vetlands						
	*If the documented raw score for I	Metric 5 is 30 point	s or higher, the	site is auto	matically co	nsidered a Cat	egory 3 wetlar	nd.
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acide Assoc. forest (wetl. &/or adj. up Sensitive geologic feature such Vernal pool (5); isolated, perch Island wetland >0.1 acre (0.04 Braided channel or floodplain/t Gross morph. adapt. in >5 tree Ecological community with glol Known occurrence state/federe [*use higher rank where mixe Superior/enhanced habitat/use	re multiple values apply in row, score row as single feature with highest point value. Provide election (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). 10); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) /or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] ature such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) ed, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) cre (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) codplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) in >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ate/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) lere mixed rank or qualifier] [exclude records which are only "historic"] abitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) by) : <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)						
6	Metric 6. Plant Co	mmunities	s, Intersp	ersio	n, Micr	otopogr	aphy	
max 20 pts. subtotal	6a. Wetland vegetation communities Score all present using 0 to 3 scales Aquatic bed Emergent Shrub Forest Mudflats Open water <20 acres (8 h	e. $0 = \frac{1}{1} = \frac{1}{2} = \frac{1}{1}$	Present and	1 ha (0.25 and 1 ha (	acre) contiguacre)] rises a smaluprises a siguirises a si	ll part of wetlan nificant part bu ficant part of w a small part an	t is of low quali etland's vegeta d is of high qua	ity ation and ality
	6b. Horizontal (plan view) interspe Select only one.		rrative Descrip				ve or disturban	ce tolerant
	High (5)  Moderately high (4) [BR/CI  Moderately high (4) [BR/CI  Moderately low (2) [BR/CN  Low (1) [BR/CM (2)]  None (0)	M (5)] mo	native spec d = Native spec nonnative & and species w/o present h = A predomin tolerant nat	ies bies are dor k/or disturba s diversity n ce of rare, t ance of nat ive sp abse	minant comp ance toleran noderate to hreatened o tive species ent or virtuall	onent of the vert native species moderately high rendangered swith nonnative y absent, and l	egetation, altho s can also be p h, but generall species sp &/or disturb nigh sp diversit	ough present, y pance by and often
	6c. Coverage of invasive plants.  Add or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover ( Absent (1)	$ \begin{array}{c} 0 = \\ 3) & 1 = \\ 0) & \underline{2} = \\ \end{array} $	dflat and Oper Absent <0.1 h Low 0.1 to <1 (0.1 to 0.5 ac Moderate 1 to High 4 ha (9.5)	Mater Clana (0.25 acre) ha (0.25 tore)] o <4 ha (2.5	res) [For BR 0 2.5 acres) 5 to 9.9 acres	[BR/CM 0.04 to s) [BR/CM 0.2	(0.1 acre)] o <0.2 ha to <02 ha (0.5	· 
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10  X Amphibian breeding pools	Hy le. ocks cm (6 in.) in.) dbh	pothetical Wet	land for Es	Estimating De			High
		0 = 1 = 2 =	Absent Present in we Present in mo amounts of hi Present in mo	ry small am oderate amo ghest quali	ounts or if mounts, but not	ot of highest qu	ality or in smal	
47	GRAND T (max 100	OIAL 3	0- 29 = Catego 0- 59 = Catego 0-100 = Catego	ry 2, good/r	moderate we	etland function,	condition, qua	ality**

Site: W-PFO-04		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	1 subtotal	Metric 1. Wetland	Area (Size)	pen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign   >50 acres (>20.2 ha) (6 p   25 to <50 acres (10.1 to <   10 to <25 acres (4 to <10.   3 to <10 acres (1.2 to <4 l   0.3 to <3 acres (0.1 to <1.   X 0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	4	Metric 2. Upland E	<b>Buffers and Su</b>	rrounding Land	Use
max 14 pts. s	subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	o m (164 ft) or more around e 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to < average <10 m (<32 ft) aro use. Select one or double or or older forest, prairie, sava s), shrubland, young 2nd gr	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter bund wetland perimeter (0) check and average. nnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
	27	Metric 3. Hydrolog	ду		
max 30 pts. s	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BF X Precipitation (1) [unless B Seasonal/intermittent surf X Perennial surface water 3c. Maximum water depth. Selectory 10.4 to 0.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in.) (3)  4.0.4 m (<16 in.) (1) [BR/C]  3e. Modifications to natural hydromy X None or none apparent (1)  Recovered (7)  Recovering (3)  Recent or no recovery (1)	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) cot only one and assign sco .) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 ir ologic regime. Score one or 2) Check all disturbance	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane Regularly inundat X Seasonally inundat on.) (2)] Seasonally saturate double check and average.  se observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] sted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	16	Metric 4. Habitat	Alteration and	Development	
max 20 pts. s	subtotal	4a. Substrate disturbance. Score  None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1)  4b. Habitat development. Select Excellent (7) Very good (6) Good (5) Medosrately good (4)	)	everage.	
		Moderately good (4)  X Fair (3) Poor to fair (2) Poor (1)  4c. Habitat alteration. Score one X None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	))	Check all disturbances of mowing grazing clearcutting selective cutting farming toxic pollutants	observed  shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PFO-04	Rater(s):	OH:jb	Date: 03/2021			
47 subtotal previous page						
-10 Metric 5. S	pecial Wetland	ls				
*If the documented	raw score for Metric 5 is 30	points or higher, the site is automatically co	onsidered a Category 3 wetland.			
documentation for e Bog, fen, wet p Assoc. forest ( Sensitive geole Vernal pool (5) Island wetland Braided chann Gross monph. Ecological cool Known occurre [*use higher   Superior/enhar	Where multiple values apply in row, score row as single feature with highest point value. Provide ch selection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc). airie (10); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) etl. &/or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] gic feature such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) isolated, perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) 0.1 acre (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) or floodplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) dapt. in >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) nunity with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] ce state/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) nk where mixed rank or qualifier] [exclude records which are only "historic"] evel habitat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) quality): <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
	lant Communit	ies, Interspersion, Micr	otopography			
6a. Wetland vegetat Score all present us Aquatic bed Emergent Shrub X Forest Mudflats Open water Moss/lichen.	sing 0 to 3 scale.	Vegetation Community Cover Scale  0 = Absent or <0.1 ha (0.25 acre) contig         [For BR/CM <0.04 ha (0.1 acre)]  1 = Present and either comprises a sma         moderate quality, or comprises a sign         is of moderate quality, or comprises  3 = Present and comprises a significant and is of high quality	Il part of wetland's vegetation and is of nificant part but is of low quality ificant part of wetland's vegetation and a small part and is of high quality			
6b. Horizontal (plan Select only one.	view) interspersion.	Narrative Description of Vegetation Quality low = Low species diversity &/or domina				
☐ High (5) ☐ Moderately h ☐ Moderate (3)	ow (2) [BR/CM (3)]	native species  mod = Native species are dominant comp nonnative &/or disturbance toleran and species diversity moderate to w/o presence of rare, threatened of high = A predominance of native species tolerant native sp absent or virtual	are dominant component of the vegetation, although disturbance tolerant native species can also be present, ersity moderate to moderately high, but generally fare, threatened or endangered species e of native species with nonnative sp &/or disturbance sp absent or virtually absent, and high sp diversity and often			
☐ Moderate 25	s for coverage. 75% cover (-5) i-75% cover (-3)	Mudflat and Open Water Class Quality 0 = Absent < 0.1 ha (0.25 acres) [For BR 1 = Low 0.1 to < 1 ha (0.25 to 2.5 acres) (0.1 to 0.5 acre)] 2 = Moderate 1 to < 4 ha (2.5 to 9.9 acre 3 = High 4 ha (9.9 acres) or more [BR/C	/CM <0.04 ha (0.1 acre)] [BR/CM 0.04 to <0.2 ha s) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)]			
6d. Microtopograph Score all present us Vegetated hu Coarse wood Standing dea		Hypothetical Wetland for Estimating Down None Low Low  Microtopography Cover Scale  0 = Absent  1 = Present in very small amounts or if many amounts of highest quality	Moderate Moderate High			
1 27 II	RAND TOTAL nax 100 pts)	3 = Present in moderate or greater amounts  0- 29 = Category 1, low wetland funct 30- 59 = Category 2, good/moderate we 60-100 = Category 3, superior wetland f	ion, condition, quality** etland function, condition, quality**			

Site: W-PFO-05		Rater(s): JB; OH		Date: 03/2021
max 6 pts. subtotal	Metric 1. Wetland	Area (SIZe) opi	en water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
	Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <1)  <0.1 acre (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
4	Metric 2. Upland E	Buffers and Sur	rounding Land	Use
max 14 pts. subtotal	MEDIUM. Buffers average  NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years MODERATELY HIGH. Re	o m (164 ft) or more around to 25 m to <50 m (82 to <164 ge 10 m to <25 m (32 ft to <6 average <10 m (<32 ft) arounds. Select one or double char older forest, prairie, savans), shrubland, young 2nd gro	wetland perimeter (7) ft) around wetland perimeter 2 ft) around wetland perimeter and wetland perimeter (0) neck and average. nah, wildlife area, etc. (7) wth forest (5) ark, conservation tillage, new	(4) er (1)
10	Metric 3. Hydrolog	ду		
max 30 pts. subtotal	3a. Sources of water. Score all th High pH groundwater (5)  Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surf Perennial surface water 3c. Maximum water depth. Sele >0.7 m (27.6 in.) (3)  0.4 to 0.7 m (16 to 27.6 in X<0.4 m (<16 in.) (1) [BR/C 3e. Modifications to natural hydro None or none apparent (1 Recovered (7) Recovering (3) X Recent or no recovery (1)	R/CM (5)]  R/CM primary source (5)]  ace water (3) (lake or stream) (5)  act only one and assign score  .) (2) [BR/CM (3)]  M 0.15 to 0.4 m (6 to <16 in.)  logic regime. Score one or complete to the complet	Part of wetland/up Part of riparian or 3d. Duration inundation/se. Semi- to permane Regularly inundat Seasonally inundat Seasonally satura double check and average.  s observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) iaturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] tted in upper 30 cm (12 in.) (1) [BR/CM (2)
11	Metric 4. Habitat A	Alteration and D	Development	
max 20 pts. subtotal	4a. Substrate disturbance. Score  None or none apparent (4)  Recovered (3)  Recovering (2)  Recent or no recovery (1)  4b. Habitat development. Select  Excellent (7)  Very good (6)  Good (5)	)	verage.	
	Moderately good (4) Fair (3) Poor to fair (2) Poor (1) 4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)	)	Check all disturbances of mowing grazing e. clearcutting selective cutting farming toxic pollutants	bbserved shrub/sapling removal herbaceous/aquatic bed removal woody debris removal sedimentation dredging nutrient enrichment

Site: W-PFO-(	05	Rater(s): OH:jb	Date: 03/2021				
25 subtotal previous page							
max 10 pts. subtotal	Metric 5. Special V	Vetlands					
	*If the documented raw score for I	Metric 5 is 30 points or higher, the site is	s automatically considered a Category 3 wetland.				
raw score*	documentation for each selection  Bog, fen, wet prairie (10); acid Assoc. forest (wetl. &/or adj. u  Sensitive geologic feature sucl Vernal pool (5); isolated, perch Island wetland >0.1 acre (0.04 Braided channel or floodplain/M Gross morph. adapt. in >5 tree Ecological community with glo Known occurrence state/feder: [*use higher rank where mixe Superior/enhanced habitat/use	emultiple values apply in row, score row as single feature with highest point value. Provide ection (photos, checklists, maps, resource specialist concurrence, data sources, references, etc).  (b); acidophilic veg., mossy substrate >10 sq.m, sphagnum or other moss (5); muck, organic soil layer (3) or adj. upland) incl. >0.25 acre (0.1 ha); old growth (10); mature >18 in. (45 cm) dbh (5) [exclude pine plantation] ure such as spring/seep, sink, losing/underground stream, cave, waterfall, rock outcrop/cliff (5) d., perched, or slope wetland (4); headwater wetland [1st order perennial or above] (3) e (0.04 ha) in reservoir, river, or perennial water >6 ft (2 m) deep (5) deplain/terrace depressions (floodplain pool, slough, oxbow, meander scar, etc.) (3) >5 trees >10 in. (25 cm) dbh: buttress, multitrunk/stool, stilted, shallow roots/tip-up, or pneumatophores (3) with global rank (NatureServe): G1*(10), G2*(5), G3*(3) [*use higher rank where mixed rank or qualifier] e/federal threatened/endangered species (10); other rare species with global rank G1*(10), G2*(5), G3*(3) re mixed rank or qualifier] [exclude records which are only "historic"] intat/use: migratory songbird/waterfowl (5); in-reservoir buttonbush (4); other fish/wildlife management/designation (3) 1: <1 acre (0.4 ha) AND EITHER >80% cover of invasives OR nonvegetated on mined/excavated land (-10)					
3	Metric 6. Plant Co	mmunities, Interspers	sion, Microtopography				
max 20 pts. subtotal	6a. Wetland vegetation communit Score all present using 0 to 3 scal Aquatic bed Emergent Shrub X Forest Mudflats Open water <20 acres (8 h	e. 0 = Absent or <0.1 ha (    For BR/CM <0.04   1 = Present and either moderate quality, or   2 = Present and either is of moderate qual	0.25 acre) contiguous acre ha (0.1 acre)] comprises a small part of wetland's vegetation and is of r comprises a significant part but is of low quality comprises a significant part of wetland's vegetation and ity, or comprises a small part and is of high quality ises a significant part or more of wetland's vegetation				
	6b. Horizontal (plan view) interspe Select only one. High (5)	ew) interspersion.  Narrative Description of Vegetation Quality  low = Low species diversity &/or dominance of nonnative or disturbance to native species					
	Moderately high (4) [BR/CI Moderate (3)[BR/CM (5)] Moderately low (2) [BR/CN X Low (1) [BR/CM (2)] None (0)	nonnative &/or di and species dive w/o presence of I high = A predominance tolerant native sp	re dominant component of the vegetation, although sturbance tolerant native species can also be present, rsity moderate to moderately high, but generally rare, threatened or endangered species of native species with nonnative sp &/or disturbance absent or virtually absent, and high sp diversity and often				
	6c. Coverage of invasive plants.  Add or deduct points for coverage  Extensive >75% cover (-5)  Moderate 25-75% cover (-1)  Sparse 5-25% cover (-1)  Nearly absent <5% cover (Absent (1)	Mudflat and Open Wate 0 = Absent <0.1 ha (0.2) 3)	er Class Quality 25 acres) [For BR/CM <0.04 ha (0.1 acre)] 0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha a (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] s) or more [BR/CM 2 ha (5 acres) or more]				
	6d. Microtopography.  Score all present using 0 to 3 sca  Vegetated hummocks/tuss  Coarse woody debris >15  Standing dead >25 cm (10	Hypothetical Wetland for ocks cm (6 in.)	for Estimating Degree of Interspersion				
	Amphibian breeding pools	None Low	Low Moderate Moderate High				
		2 = Present in moderat amounts of highest	all amounts or if more common of marginal quality e amounts, but not of highest quality or in small				
30	GRAND T (max 100	30- 59 = Category 2, g	ow wetland function, condition, quality** good/moderate wetland function, condition, quality** superior wetland function, condition, quality**				

Page 2 of 2

Site: W-PFO-06		Rater(s): JB; OH		Date: 03/2021	
max 6 pts.	2 subtotal	Metric 1. Wetland	Area (Size) or	pen water body (excluding aquati	Blue Ridge and Cumberland Mountains. If an c beds and seasonal mudflats) is >20 acres a) of it to the wetland size for Metric 1.
		Select one size class and assign  >50 acres (>20.2 ha) (6 p  25 to <50 acres (10.1 to <  10 to <25 acres (4 to <10  3 to <10 acres (1.2 to <4    X 0.3 to <3 acres (0.1 to <1)  0.1 to <0.3 acre (0.04 to <0.1 acres (0.04 to <0.1 acres (0.04 ha) (0)	ts) :20.2 ha) (5) [BR/CM (6)] .1 ha) (4) [BR/CM (6)] na) (3) [BR/CM (5)] .2 ha) (2) [BR/CM (3)]	Sources/assumptions for s	size estimate (list):
	4	Metric 2. Upland E	Buffers and Su	rrounding Land	Use
max 14 pts.	subtotal	MEDIUM. Buffers average NARROW. Buffers average VERY NARROW. Buffers  2b. Intensity of surrounding land VERY LOW. 2nd growth of LOW. Old field (>10 years	o m (164 ft) or more around a 25 m to <50 m (82 to <16 ge 10 m to <25 m (32 ft to < average <10 m (<32 ft) arouse. Select one or double cor older forest, prairie, savas), shrubland, young 2nd gresidential, fenced pasture, p	wetland perimeter (7) 4 ft) around wetland perimeter 82 ft) around wetland perimeter und wetland perimeter (0) check and average. nnah, wildlife area, etc. (7) owth forest (5) park, conservation tillage, new	(4) er (1)
	7	Metric 3. Hydrolog	ду		
max 30 pts.	subtotal	3a. Sources of water. Score all the High pH groundwater (5)  Other groundwater (3) [BI X Precipitation (1) [unless B Seasonal/intermittent surform Perennial surface water 3c. Maximum water depth. Seleston 100 -	R/CM (5)] R/CM primary source (5)] ace water (3) (lake or stream) (5) act only one and assign sco act.) (2) [BR/CM (3)] M 0.15 to 0.4 m (6 to <16 in ologic regime. Score one or 2)  Check all disturbance in ditch	Part of wetland/up Part of riparian or 3d. Duration inundation/s re. Semi- to permane Regularly inundat X Seasonally inundat on.) (2)] Seasonally satura double check and average.  so observed point source (non	in (1) ake and other human use (1) bland (e.g., forest), complex (1) upland corridor (1) saturation. Score one or dbl. check & avg. ently inundated/saturated (4) ed/saturated (3) [BR/CM (4)] ated (2) [BR/CM (4)] sted in upper 30 cm (12 in.) (1) [BR/CM (2)] stormwater)
	7	Metric 4. Habitat	Alteration and I	Development	
max 20 pts.	subtotal	4a. Substrate disturbance. Score None or none apparent (4) Recovered (3) Recovering (2) Recent or no recovery (1) Ab. Habitat development. Select Excellent (7) Very good (6) Good (5) Moderately good (4) Fair (3) Recovering (2)	)	Check all disturbances o	observed □ shrub/sapling removal
		Poor (1)  4c. Habitat alteration. Score one None or none apparent (9) Recovered (6) X Recovering (3) Recent or no recovery (1)	))	grazing	herbaceous/aquatic bed removal   woody debris removal   sedimentation   dredging   nutrient enrichment

Site: W-PFO-0	06	Rater(s): OH:jb	Date: 03/2021
20 subtotal previous page			
max 10 pts. subtotal	Metric 5. Special	Wetlands	
max 10 ptc.	*If the documented raw score for	Metric 5 is 30 points or higher, the site	is automatically considered a Category 3 wetland.
raw score*	documentation for each selection  Bog, fen, wet prairie (10); aci Assoc. forest (wetl. &/or adj. Sensitive geologic feature su Vernal pool (5); isolated, per Island wetland >0.1 acre (0.0 Braided channel or floodplair Gross morph. adapt. in >5 tre Ecological community with gl Known occurrence state/fede [*use higher rank where mix Superior/enhanced habitat/us	n (photos, checklists, maps, resource s dophilic veg., mossy substrate >10 sq.m, spi upland) incl. >0.25 acre (0.1 ha); old growth ch as spring/seep, sink, losing/underground ched, or slope wetland (4); headwater wetland (4) in reservoir, river, or perennial water y/terrace depressions (floodplain pool, slough ese >10 in. (25 cm) dbh: buttress, multitrunk/obal rank (NatureServe): G1*(10), G2*(5), Garal threatened/endangered species (10); othed rank or qualifier] [exclude records which ese: migratory songbird/waterfowl (5); in-reservence.	d [1st order perennial or above] (3) 6 ft (2 m) deep (5) 7, oxbow, meander scar, etc.) (3) 8 tool, stilted, shallow roots/tip-up, or pneumatophores (3) 8 (3) [*use higher rank where mixed rank or qualifier] 8 er rare species with global rank G1*(10), G2*(5), G3*(3)
3	Metric 6. Plant Co	ommunities, Interspe	sion, Microtopography
max 20 pts. subtotal	6a. Wetland vegetation commun Score all present using 0 to 3 sc. Aquatic bed Emergent Shrub X Forest Mudflats Open water <20 acres (8 Moss/lichen. Other	ale. 0 = Absent or <0.1 ha  [For BR/CM <0.0] 1 = Present and eithe  moderate quality, 2 = Present and eithe is of moderate qu	(0.25 acre) contiguous acre  ha (0.1 acre)] r comprises a small part of wetland's vegetation and is of or comprises a significant part but is of low quality r comprises a significant part of wetland's vegetation and ality, or comprises a small part and is of high quality or see a significant part or more of wetland's vegetation
	6b. Horizontal (plan view) intersp Select only one.	persion. Narrative Description	of Vegetation Quality rersity &/or dominance of nonnative or disturbance tolerant
	High (5)  Moderately high (4) [BR/C Moderate (3)[BR/CM (5)]  Moderately low (2) [BR/C Low (1) [BR/CM (2)]  None (0)	CM (5)]  mod = Native species nonnative &/or M (3)]  mod = Native species nonnative &/or and species div w/o presence of high = A predominance	are dominant component of the vegetation, although disturbance tolerant native species can also be present, ersity moderate to moderately high, but generally rare, threatened or endangered species of native species with nonnative sp &/or disturbance ap absent or virtually absent, and high sp diversity and often
	6c. Coverage of invasive plants. Add or deduct points for coverage Extensive >75% cover (-5) Moderate 25-75% cover (-5)	ee. <u>Mudflat and Open Water</u> 5) 0 = Absent < 0.1 ha (0	the presence of rate, threatened, or endangered species  ter Class Quality  .25 acres) [For BR/CM <0.04 ha (0.1 acre)]  (0.25 to 2.5 acres) [BR/CM 0.04 to <0.2 ha
	X Sparse 5-25% cover (-1) Nearly absent <5% cover (0) Absent (1)		ha (2.5 to 9.9 acres) [BR/CM 0.2 to <02 ha (0.5 to 5 acre)] res) or more [BR/CM 2 ha (5 acres) or more]
	6d. Microtopography.  Score all present using 0 to 3 sc  Vegetated hummocks/tus  Coarse woody debris >15  Standing dead >25 cm (1  Amphibian breeding pools	sale. socks 5 cm (6 in.) 0 in.) dbh s None Low  Microtopography Co 0 = Absent	for Estimating Degree of Interspersion  Low Moderate Moderate High  ver Scale  mall amounts or if more common of marginal quality
		2 = Present in moder amounts of highe	ate amounts, but not of highest quality or in small
23	GRAND (max 100	30- 59 = Category 2	low wetland function, condition, quality** good/moderate wetland function, condition, quality** superior wetland function, condition, quality**

\*\*Based on ORAM Score Calibration Report for the scoring breakpoints between wetland categories: http://www.epa.state.oh.us/dsw/401/401.html

Page 2 of 2



CREATE AMAZING.

Burns & McDonnell World Headquarters 9400 Ward Parkway Kansas City, MO 64114 •• 816-333-9400 •• 816-333-3690 •• www.burnsmcd.com



# ONSITE AND OFFSITE DRAINAGE REPORT

# Golden Triangle Solar Project Phase 2

**Lowndes County, Mississippi** 

# PREPARED FOR

MS Solar 6, LLC

800 Brickell Ave, Suite 1100 Miami, Florida 33131 Contact: Daniel Ferrell Director of Project Development Origis Energy USA (786) 757 4332

# PREPARED BY



421 Fayetteville St., Suite 600 Raleigh, NC 27601 Contact: John Barefoot (919) 653-5843 June 2021

# TABLE OF CONTENTS

1	INTF	RODUCTION	2
	1.1	Project Background	2
	1.2	Site Description	2
2	Eng	SINEERING METHODOLOGY	2
	2.1	Model Selection	3
	2.2	Model Topography and Extents	3
	2.3	Soil Data	3
	2.4	Precipitation	3
	2.5	Hydrologic Analysis	3
	2.6	Hydraulic Analysis	
	2.7	Inundation Depth Calculation Summary	4
	2.8	Velocity Calculation Summary	4
	2.9	Scour Depth Calculation Summary	4
3	Con	ICLUSIONS	5
4	Ref	ERENCES	6

# LIST OF FIGURES

- Figure 1. Site Location Map
- Figure 2. Topographic Work Map
- Figure 3. HEC-RAS Study Map
- Figure 4. Pre-Development 100-Year Depth Map
- Figure 5. Pre-Development 100-Year Flow Velocity Map
- Figure 6. Pre-Development 100-Year Scour Depth Map
- Figure 7. USGS Quadrangle Map
- Figure 8. FEMA FIRM Map
- Figure 9. NRCS WSS Site Soil Map

# LIST OF APPENDICES

Appendix A. Figures

# 1 INTRODUCTION

# 1.1 PROJECT BACKGROUND

Kimley-Horn was retained by MS Solar 6, LLC to conduct a hydrologic analysis for the proposed solar photovoltaic (PV) generation facility Golden Triangle Solar in Lowndes County, Mississippi (Figure 1). The project site is located approximately 10 miles east of the city of Starkville, Mississippi and 2 miles west of Golden Triangle Regional Airport. This report summarizes the results of the Phase 2 existing conditions drainage analysis in and around the project site. The drainage analysis estimates the existing conditions 100-year inundation depths, flow velocities, and scour depths in and around the project site.

# 1.2 SITE DESCRIPTION

The project site is approximately 1,835 acres located in Lowndes County, Mississippi east of Catalpa Creek. It is bounded by Artesia Road to the north and Swedenburg Road to the south. The existing site land cover consists primarily of pasture, agricultural land, wetlands, and forest.

The latest USGS Quad Map (Figure 7) shows a perennial stream, Gilmer Creek, to the north of the site. North Branch Magowah Creek passes through the southern section of the site, and there are also numerous intermittent streams located within the site that are shown as dashed blue lines.

The project site is located within **FEMA FIRM Panels 28087C025K and 28087C0250K** (Figure 8). **FEMA unshaded Zone X** contains areas determined to be outside the 0.2% (500-year) annual chance floodplain. **FEMA Zone A** contains areas determined by FEMA to be inside the 1% annual chance special flood hazard area via Approximate or Limited Detail methods. **FEMA Zone AE** contains areas determined to be in the 1% annual chance special flood hazard area via Detailed methods, with provided base flood elevations. **FEMA Zone AE Floodway** is the portion of the floodplain reserved for the passage of the base flood.

Most of the site lies within **FEMA unshaded Zone X.** The southern portion of the site lies within **FEMA Zone A** associated with North Branch Magowah Creek.

The existing topography is generally moderately sloped with some steep areas. The slope ranges from 0 to 10 percent within most of the site, with slopes over 10 percent located along ridges and steeper channel banks. The majority of the site has less than a 3 percent slope. The elevation ranges from 205 to 280 feet, with the northern portion of the site draining to the west, while the rest of the site drains to North Branch Magowah Creek to the south and east.

Kimley-Horn received survey of specific hydraulic structures within and adjacent to the project site from Survey and Mapping, LLC (SAM) on April 29th, 2021. A total of 26 culverts were confirmed and measured during the site visit.

# 2 ENGINEERING METHODOLOGY

The project site is at risk from onsite flooding due to extreme onsite precipitation events. This analysis modeled an extreme 100-year precipitation event occurring over the onsite area, and adjacent offsite area. This methodology captures the initial flooding resulting from onsite and immediate offsite precipitation and subsequent runoff.

#### 2.1 MODEL SELECTION

The topography within the project site creates drainage patterns that vary in direction across the land surface outside of defined channels. 2-dimensional (2-D) hydrodynamic models are ideal for modeling the runoff and drainage patterns of flat areas without strong one-dimensional channel definition. The modeling in this analysis utilizes the 2-D version of the U.S Army Corps of Engineers' Hydrologic Center River Analysis System (HEC-RAS) version 5.0.7.

# 2.2 MODEL TOPOGRAPHY AND EXTENTS

The model topography is derived from LiDAR elevation data obtained from the Mississippi Automated Resource Information System website combined with onsite survey topographic data (Figure 2). All mapping was done using vertical datum NAVD 88. Onsite topography was used to define the onsite 2-D modeling boundary just beyond the estimated runoff boundary to allow the model to calculate the direction of runoff (Figure 3).

#### 2.3 SOIL DATA

The soil data obtained from the NRCS Web Soil Survey indicates the site is composed mostly of silty clay and silty clay loam with assigned Hydrologic Soil Groups C, D, and C/D (Figure 9). Due to the extreme nature of the modeled 100-year rainfall event and the resulting inundation conditions that follow, soils with dual classification are assumed to be undrained and are modeled as group D. Assuming that the soils are undrained in the modeled scenario produces conservative runoff volumes and inundation depths.

# 2.4 PRECIPITATION

The precipitation depth for the 24-hour, 1-percent (100-year) annual chance event was obtained from the NOAA Atlas 14 Precipitation-Frequency Estimates. The rainfall distribution hyetograph utilized for onsite hydrology modeling was created by distributing the 24-hour rainfall total over the SCS Type II synthetic rainfall distribution curve. The SCS Type II distribution was selected based on geographic boundaries for SCS rainfall distributions provided in Appendix B of the USDA's Technical Release 55.

#### 2.5 HYDROLOGIC ANALYSIS

Onsite hydrology was simulated as excess rainfall runoff applied directly to the onsite 2-D modeling area. The total 24-hour rainfall depth for the 100-year annual chance event is 9.45 inches. Kimley-Horn used the USACE's Hydrologic Modeling System software (HEC-HMS) version 4.7 to produce a timeseries of the precipitation excess from the rainfall distribution hyetograph. The precipitation excess timeseries is the amount of rainfall that is converted to runoff after subtracting the rainfall captured in microtopography on the surface and losses due to infiltration. Kimley-Horn used the Soil Conservation Service (SCS) curve number loss method within HEC-HMS to calculate the losses over the 2-D modeling area based on land cover and soil data. National Land Cover Database (NLCD) 2016 data and aerial imagery were used to delineate the existing land cover within the 2-D modeling area. The resulting precipitation excess timeseries was applied equally across the 2-D modeling area.

#### 2.6 HYDRAULIC ANALYSIS

In HEC-RAS, the 2-D modeling boundary was divided into a 2-D computational grid with a maximum cell size of 100-feet (Figure 3). The combined LiDAR and onsite topographic survey model terrain surface was used to develop a family of rating tables for the stage storage volume in each computational grid cell. HEC-RAS defines cross sections at each cell face of the computational grid to more accurately simulate flow between cells.

The computational grid was further refined by adding breaklines to subdivide cells along important hydraulic features such as channels, culverts, roads, and berms (Figure 3). The minimum computational grid cell size along breaklines is 4 feet.

Culvert measurements received from survey and remote observation were incorporated into the 2-D hydraulic model. Survey information and professional judgement were used to assign Manning's n, entrance loss, and exit loss values during the modeling process. Aerial imagery and LiDAR were referenced to assign culvert lengths and invert elevations.

Roughness coefficients were applied to the grid to account for energy losses to water flow due to friction based on existing landcover data.

Outflow boundary conditions for the model were defined at the boundary of the 2-D computational grid. The boundary condition type was set to normal depth and assigned a friction slope ranging from 0.5 to 2.5 percent.

# 2.7 INUNDATION DEPTH CALCULATION SUMMARY

Predicted inundation depths for existing conditions are calculated based on the 2-D modeling simulation and are mapped within the study area. Results are shown in Figure 4. The mapping reflects depths that are expected to equal or exceed 0.25 feet. The direct rainfall method produces depth in every grid cell of the mesh. Shallower depths are removed from the mapping as is typically done for models based on the direct rainfall method.

The inundation depths on the project parcels range from 0 to 13 feet, with the greatest depths occurring within North Branch Magowah Creek. The regions with over 2 feet of expected depth are mostly restricted to channels, established FEMA flood zones, or areas that are impounded by a downstream change in elevation.

#### 2.8 VELOCITY CALCULATION SUMMARY

Predicted flow velocities were examined in the HEC-RAS model with results shown in Figure 5. Flow velocities that are shown to be greater than 1 foot per second are associated with defined flow paths and their associated overflow areas.

# 2.9 SCOUR DEPTH CALCULATION SUMMARY

Pier scour depth is calculated within the project parcels using guidance in *HEC-18: Evaluating Scour at Bridges, Fifth Edition* (FHWA, 2012). The calculated pier scour depth is based on the modeled maximum flow velocity and maximum flow depth. While the flow depth and flow velocity may not reach their maximum levels at the same time and location, this assumption produces a conservative estimate of scour for the entire site. The model indicates portions of the site will experience potential scour depths in excess of 1 foot, generally along defined flow paths (channels and draws). Anticipated scour depths within the project parcels are shown in Figure 6.



# 3 CONCLUSIONS

The inundation depths generally range from 0 to 13 feet in most areas onsite. The deepest inundation is shown within the southernmost parcels and is associated with depths within the channel of North Branch Magowah Creek. Areas of inundation within the other parcels are generally associated with existing onsite channels and local depression ponding.

# Recommendations:

- Based on this analysis, we recommend avoiding array development within areas of higher inundation depth, flow velocity, and scour, particularly within FEMA Zone A (Figures 4 and 8).
- Based on this analysis, we recommend final system design consider placing equipment outside of areas with inundation depths exceeding 1 foot or provide countermeasures.
   Countermeasures may include extending posts to elevate equipment, installing additional culverts to provide inundation relief, or grading modifications to eliminate local depressions where ponding may occur.
- Based on this analysis, we recommend final system design consider placing equipment outside of areas with probable scour depths exceeding 1 foot or provide countermeasures.
   Countermeasures may include extending posts deeper, applying erosion control matting, or grading to smooth out local areas with a high slope.

# 4 REFERENCES

Chow, V.T., 1959. Open Channel Hydraulics. McGraw-Hill, New York.

Cook, L.M. and McCuen, R.H. (2013). Hydrologic Response of Solar Impacts. Journal of Hydrologic Engineering, 18(5), 536-541.

http://ascelibrary.org/doi/abs/10.1061/%28ASCE%29HE.1943-5584.0000530?journalCode=jhyeff&

Folmar, N.D., and A.C. Miller. 2008. Development of an empirical lag time equation. Amer. Soc. of Civil Eng. Journal of Irrigation and Drainage, Vol. 134, No. 4. https://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=27002.wba

NOAA Atlas 14, Point Precipitation Frequency (PF) Data Server.. https://hdsc.nws.noaa.gov/hdsc/pfds/

U.S. Army Corps of Engineers (USACE, 2016). 2016. CPD-68: HEC-RAS River Analysis System User's Manual

http://www.hec.usace.army.mil/software/hec-ras/documentation/HEC-RAS%205.0%20Users%20Manual.pdf

U.S. Army Corps of Engineers (USACE, 2016). 2016. CPD-68A: HEC-RAS River Analysis System 2D Modeling User's Manual

http://www.hec.usace.army.mil/software/hec-ras/documentation/HEC-RAS%205.0%202D%20Modeling%20Users%20Manual.pdf

U.S. Army Corps of Engineers (USACE, 2016). 2016. CPD-70: HEC-RAS River Analysis System Application Guide

http://www.hec.usace.army.mil/software/hec-ras/documentation/HEC-RAS%205.0%20Applications%20Guide.pdf

U.S. Army Corps of Engineers (USACE, 2018). 2018. CPD-74A: Hydrologic Modeling System HEC-HMS User's Manual

https://www.hec.usace.army.mil/software/hec-hms/documentation/HEC-HMS Users Manual 4.3.pdf

U.S. Department of Agriculture (USDA, 1986). 1986. TR-55: Urban Hydrology for Small Watersheds


https://www.nrcs.usda.gov/Internet/FSE DOCUMENTS/stelprdb1044171.pdf


- U.S. Department of Agriculture, Natural Resources Conservation Service. Part 654 Stream Restoration Design National Engineering Handbook. Chapter 8, Threshold Channel Design. August 2007.
- U.S. Department of Agriculture, Natural Resources Conservation Service. Part 630 Hydrology National Engineering Handbook Chapter 15, Time of Concentration. May 2010. <a href="https://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=27002.wba">https://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=27002.wba</a>
- U.S. Department of Transportation. Federal Highway Administration (2012). Evaluating Scour at Bridges Fifth Edition (Publication No. FHWA-HIF-12-003) https://www.fhwa.dot.gov/engineering/hydraulics/pubs/hif12003.pdf
- Yang, L., Jin, S., Danielson, P., Homer, C.G., Gass, L., Bender, S.M., Case, A., Costello, C., Dewitz, J.A., Fry, J.A., Funk, M., Granneman, B.J., Liknes, G.C., Rigge,


Kimley » Horn


M.B., and Xian, G., 2018, A new generation of the United States National Land Cover Database—Requirements, research priorities, design, and implementation strategies: ISPRS Journal of Photogrammetry and Remote Sensing, v. 146, p. 108–123, at <a href="https://doi.org/10.1016/j.isprsjprs.2018.09.006">https://doi.org/10.1016/j.isprsjprs.2018.09.006</a>.


# Appendix A – Figures





















# Figure 9. NRCS WSS Site Soil Map



# **Hydrologic Soil Group**

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
BrA	Brooksville silty clay, 0 to 1 percent slopes	D	1,132.1	12.8%
BrB	Brooksville silty clay, 1 to 3 percent slopes	D	1,445.8	16.4%
Ср	Catalpa silty clay	C/D	615.0	7.0%
DeC2	Demopolis-Binnsville complex, 2 to 8 percent slopes, eroded	D	608.5	6.9%
Gr	Griffith silty clay	D	187.5	2.1%
KpB2	Kipling silty clay loam, 2 to 5 percent slopes, eroded	D	62.7	0.7%
KpC2	Kipling silty clay loam, 5 to 8 percent slopes, eroded	D	2.8	0.0%
Le	Leeper silty clay, 0 to 2 percent slopes, occasionally flooded	D	473.1	5.4%
OkA	Okolona silty clay, 0 to 1 percent slopes	D	317.6	3.6%
OkB	Okolona silty clay, 1 to 3 percent slopes	D	919.0	10.4%
SuB2	Sumter silty clay loam, 2 to 5 percent slopes, eroded	С	178.1	2.0%
SuC2	Sumter silty clay loam, 5 to 12 percent slopes, eroded	С	269.4	3.1%
SvD3	Sumter-Demopolis- Chalk outcrop complex, 5 to 20 percent slopes, severely eroded		44.6	0.5%
VaA	Vaiden silty clay, 0 to 2 percent slopes	D	1,196.3	13.6%
VaB2	Vaiden silty clay, 2 to 5 percent slopes, eroded	D	1,247.9	14.1%
VaC2	Vaiden silty clay, 5 to 8 percent slopes, eroded	D	13.1	0.1%
W	Water		110.1	1.2%
Totals for Area of Inte	rest		8,823.8	100.0%

# **Description**

Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The soils in the United States are assigned to four groups (A, B, C, and D) and three dual classes (A/D, B/D, and C/D). The groups are defined as follows:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

If a soil is assigned to a dual hydrologic group (A/D, B/D, or C/D), the first letter is for drained areas and the second is for undrained areas. Only the soils that in their natural condition are in group D are assigned to dual classes.

# **Rating Options**

Aggregation Method: Dominant Condition

Component Percent Cutoff: None Specified

Tie-break Rule: Higher





# Protected Species Habitat Assessment Report



MS Solar 6, LLC

Golden Triangle II Solar Project

Revision 4.0 12/8/2021

# Protected Species Habitat Assessment Report

prepared for

MS Solar 6, LLC Golden Triangle II Solar Project Lowndes County, Mississippi

12/8/2021

prepared by

Burns & McDonnell Engineering Company, Inc. Atlanta, Georgia

COPYRIGHT © 2021 BURNS & McDONNELL ENGINEERING COMPANY, INC.

# **TABLE OF CONTENTS**

			Page No.
1.0	INTR	ODUCTION	1-1
2.0	MET	HODOLOGY	2-1
	2.1	Review Existing Information	
	2.2	Environmental Field Surveys	
3.0	RES	ULTS	3-1
	3.1	Land Use Types	3-1
		3.1.1 Active Agriculture and Pasture	3-1
		3.1.2 Forest	
		3.1.3 Wetlands	
		3.1.4 Streams	
	3.2	Protected Species Information	
		3.2.1 Species Protected under the Endangered Species Act	
		3.2.2 Bald Eagles and Migratory Birds	
		3.2.3 TVA Natural Heritage Database Results	3-11
4.0 5.0		CLUSIONS	
APPE APPE	NDIX NDIX	A - PROTECTED SPECIES INFORMATION B - MAPS AND FIGURES C - REPRESENTATIVE PHOTOGRAPHS D - ACOUSTIC BAT SURVEY REPORT	
		LIST OF TABLES	Page No.
Table 3		Mississippi Department of Wildlife, Fisheries, and Parks - Natural Heritage Program -Protected Species and Species of Special Concern Potentially Occurring within two miles of the Survey Area	3-6
Table 3	3-3:	TVA Heritage Database Plants Potentially Occurring within the Project Site	3-11

# LIST OF ABBREVIATIONS

Abbreviation Term/Phrase/Name

BCC Birds of Conservation Concern

BGEPA Bald and Golden Eagle Protection Act

Burns & McDonnell Engineering Company, Inc.

dbh diameter at breast height

ESA Endangered Species Act of 1973

GIS Geographic Information System

Hwy Highway

HUC Hydrologic Unit Code

IPaC Information for Planning and Conservation

MBTA Migratory Bird Protection Act

MDWFP Mississippi Department of Wildlife, Fisheries, and Parks

MS Mississippi

MW megawatt

NLEB Northern long-eared bat

NRCS Natural Resources Conservation Service

PEM palustrine emergent

PFO palustrine forested

PSS palustrine scrub shrub

PUB palustrine unconsolidated bottom

Project Golden Triangle Solar Project

ROW Right-of-way

SR State Route

TVA Tennessee Valley Authority

US United States

USEPA U.S. Environmental Protection Agency

USFWS U.S. Fish and Wildlife Service

USGS U.S. Geological Survey

MS Solar 6, LLC i Burns & McDonnell

# 1.0 INTRODUCTION

MS Solar 6, LLC, is evaluating a site in Lowndes County, Mississippi (MS) for potential development of a 150 megawatt (MW) solar energy facility (Golden Triangle Solar II Project or Project). Burns & McDonnell Engineering Company, Inc. (Burns & McDonnell) was contracted by MS Solar 6, LLC, to conduct a habitat assessment for the proposed Project, specifically potential parcels upon which solar array facility sites may be installed, in Lowndes County, MS (**Figure 1, Appendix A**). The Survey Area consists of approximately 1,525 acres of land that is predominantly used for pasture, hay production, and agriculture with fragmented forested areas throughout. The Project is east of the US Hwy Alt 45, south and adjacent to Gilmer Wilburn Road, North of Swedenburg Road, and west of Hardy Billups Road in Artesia, MS, which is approximately 8 miles east of Starkville, MS and 10 miles west of Columbus, MS. The Project was surveyed for ecological resources on March 8 through March 13, 2021.

Burns & McDonnell ecologists evaluated characteristics of the Project limits to determine presence or absence of suitable habitat for federal and/MDWFP Natural Heritage Database rare species. The habitat assessment survey was conducted throughout all parcels being considered for the proposed Project (Survey Area) as identified by MS Solar 6, LLC. Based on results of this protected species habitat assessment, in addition to results from other environmental and civil surveys, MS Solar 6, LLC will implement a project design that minimizes environmental impacts to the greatest extent practicable.

# 2.0 METHODOLOGY

Key methods used for determining the presence or absence of potential and suitable protected species habitats within the Survey Area are: Review of existing publicly available information published by federal and states agencies such as the U.S. Fish and Wildlife Service (USFWS), Mississippi Department of Wildlife, Fisheries, and Parks (MDWFP), Tennessee Valley Authority's (TVA) Natural Heritage Database, and performing pedestrian surveys.

# 2.1 Review Existing Information

Prior to conducting field work, Burns & McDonnell biologists reviewed USFWS Information for Planning and Consultation (IPaC) results letter (Appendix A) sent to Burns & McDonnell on April 20, 2021. The IPaC provides information regarding special status species with potential to occur within the Survey Area (Consultation Code: 04EM1000-2021-SLI-0745, Event Code: 04EM1000-2021-E-01680). Using the information provided in the IPaC, Burns & McDonnell's field team assessed whether the proposed Project had potential to affect certain species that are protected under the Endangered Species Act of 1973 (ESA) specifically those species that are federally listed, proposed and/or candidate species), as well as bald eagles (*Haliaeetus leucocephalus*), golden eagles (*Aquila chrysaetos*), migratory birds (including raptor species), and the occurrence of suitable habitat for these species within the Survey Area. Additionally, consultation was conducted with MDWFP's National Heritage Program which provided a list of state protected species and state ranked species of special concern with potential to occur within two miles of the Survey Area. U.S. Fish and Wildlife's Range Wide Indiana Bat Survey Guidelines (USFWS, 2019) was also reviewed. Lastly, review of TVA's natural heritage data was reviewed which provides known occurrences of rare species within three and five miles of the Survey Area for terrestrial and botanical species, respectively.

Field maps were created using the available Geographic Information System (GIS) data including U.S. Geological Survey (USGS) topographic data and Natural Resources Conservation Service (NRCS) soil survey data. This information was thoroughly reviewed to determine which protected species could occur within the Survey Area. In addition to federal and state agency sources, a literature review was conducted on each species for pertinent information regarding species' distinct physical characteristics, vegetative preferences, diet, motility, home range requirements, reproductive needs, and sensitivity to anthropogenic disturbances.

Based on a review of available information (Appendix A), it was determined suitable habitat may occur within the Survey Area for three federally protected species, one of which is also a state-listed species.

Additionally, based on data provided by MWDFP Natural Heritage Program through consultation, there are 15 state ranked species of special concern with potential to occur within two miles of the Survey Area.

# 2.2 Environmental Field Surveys

Field surveys were conducted from March 8 - 13, 2021. The Survey Area begins on the south side of US Hwy 82/State Route (SR) 12 and extends south to Gilmer-Wilburn Road (Vicinity Map, Appendix B). The survey was conducted within five parcels under consideration for the proposed Project. The Survey Area encompassed approximately 1,525 acres (Survey Area Map; Appendix B).

During the field surveys, data was collected on existing vegetative cover/land use, potentially suitable habitats of protected species, and general wildlife observations. The field investigation consisted of pedestrian surveys within the Survey Area depicted in the Survey Area Map for components of the overall Project.

Additionally, in accordance with TVA's agreement with the USFWS, it was determined that presence/absence surveys should be conducted to assess potential occurrence of the northern long-eared bat (NLEB). Accordingly, Burns & McDonnell conducted presence/absence surveys for the NLEB using acoustic methods on August 3 through August 10, 2021. The survey protocol followed the most current (2020) U.S. Fish and Wildlife (USFWS) *Rangewide Indiana Bat Survey Guidelines* (Guidelines).

# 3.0 RESULTS

# 3.1 Land Use Types

The Project is within the U.S. Environmental Protection Agency (USEPA) Blackland Prairie Ecoregion (Level 4) and is within the Middle Tombigbee River [Hydrologic Unit Code (HUC) 03160106] and Tibbee Creek (HUC 03160104) watersheds.

Four dominant vegetative cover/land use communities were observed within the Survey Area, including active agriculture/pasture, bottomland hardwood forest, and upland forest (Figures 3A-3N: Habitat Maps, Appendix B). Descriptions of these communities are provided below and representative photographs of each habitat community type are provided in Appendix C.

# 3.1.1 Active Agriculture and Pasture

Active agricultural and pasture is the primary land use community found in the Survey Area and composed approximately 73.64 percent (1123 acres) of the Survey Area. Areas identified as active agriculture include cattle pasture, hay production, and row crop fields. Vegetation in these communities is maintained in an early successional state due to crop growth/harvesting, and regular management/mowing as well as repeated cattle grazing. Corn is planted in late spring and covers the row crop fields on approximately 125 acres. Vegetation observed in pastures consists of primarily tall fescue grass (*Festuca arundinacea*), Johnson grass (*Sorghum halepense*), annual bluegrass (*Poa annua*), couch grass (*Elymus repens*), bermuda (*Cynodon dactylon*), annual ragweed (*Ambrosia atemisiifolia*), cheatgrass (*Bromus tectorum*), perennial ryegrass (*Lolium perenne*), green foxtail (*Setaria viridis*), butterweed (Packera glabella), bulbous bittercress (*Cardamine bulbosa*), soft rush (*Juncus effusus*), Cherokee sedge (*Carex cherokeensis*), fox sedge (*Carex vulpinoidea*), red sorrel (*Rumex acetosella*), curly dock (*Rumex crispus*), prairie fleabane (*Erigeron strigosus*), jimsonweed (*Datura stramonium*), Carolina horsenettle (*Solanum carolinense*), bull thistle (*Cirsum vulgare*), wild violet (*Viola sororia*), field garlic (*Allium vineale*), and sensitive partridge pea (*Chamaecrista nictitans*).

Observations of garter snake (*Thamnophis sirtalis*), blue bird (*Sialia sialis*), killdeer (*Charadrius vociferus*) and field sparrow (*Spizella pusilla*) were documented in open pasture and active agriculture habitats. Flyovers of American crow (*Corvus brachyrhynchos*), black vulture (*Coragyps astratus*), and red-tailed hawk (*Buteo jamaicensis*) also were observed in these areas.

# **3.1.2** Forest

Approximately 17.6 percent (268 acres) of the Survey Area is composed of both upland and bottomland forested areas..

# **Bottomland Forest**

Bottomland hardwood forest communities in the Survey Area are composed of a canopy age ranging from approximately 20 to 70+ years old. Dominant vegetation observed consisted of water hickory (*Carya aquatica*), willow oak (*Quercus phellos*), cherrybark oak (*Quercus pagoda*), swamp chestnut oak (*Quercus michauxii*), silky dogwood (*Cornus amomum*), osage orange (*Maclura pomifera*), green ash (*Fraxinus pernnsylvanica*), eastern red cedar (*Juniperus virginiana*), water locust (*Gledista aquatica*), shagbark hickory (*Carya ovata*), box elder (*Acer negundo*), red maple (*Acer rubrum*), American sycamore (*Platanus occidentalis*), sugarberry (*Celtis laevigata*), Cherokee sedge (*Carex cherokeensis*), wild petunia (*Ruellia humilis*), tall fescue, poison ivy (*Toxicodendron radicans*), greenbrier (*Smilax rotundifolia*), Virginia spiderwort (*Tradescantia virginiana*), Virginia creeper (*Parthenocissus quinquefolia*), prairie ironweed (*Vernonia fasciculata*), hairy buttercup (*Ranunculus sardous*), and resurrection fern (*Pleopeltis polypodioides*),

# **Upland Forest**

Upland forest communities in the Survey Area and are composed of a canopy age ranging from approximately 20 to 70 years old. Dominant vegetation observed consisted of white oak (*Quercus alba*), southern red oak (*Quercus falcata*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), mockernut hickory (*Carya tomentosa*), shagbark hickory, pignut hickory (*Carya glabra*), loblolly pine (*Pinus taeda*), eastern red cedar, American elm (*Ulmus americana*), honey locust (*Gleditsia triacanthos*), osage orange, Chinese privet, Devil's walkingstick (*Aralia spinosa*), Christmas fern (*Polystichum acrostichoides*), Japanese honeysuckle (*Lonicera japonica*), multiple greenbrier species (*Smilax spp.*), muscadine (*Vitus rotundifolia.*), Virginia creeper, blackberry (*Rubus spp.*), wooly panic grass (*Dichanthelium acuminatum*), and wild violet (*Viola sororia*).

Observations of American Robin (*Turdis migratorius*), northern mockingbird (*Mimus polyglottos*), cedar waxwing (*Bombycilla cedrorum*), downy woodpecker (*Dryobates pubescens*), great-horned owl (*Bubo virginianus*), upland chorus frog (*Pseudacris feriarum*), and eastern king snake (*Lampropeltis getula*) were noted in both of the upland and bottomland forest areas.

# 3.1.3 Wetlands

During the field surveys, a total of 20 open waters and 19 wetlands were identified within the Survey Area of the Project. Palustrine unconsolidated bottom wetlands (PUB) accounted for approximately 14.15 acres (less than one percent) of the total Survey Area and were primarily located in open fields. Common vegetation associated with the banks and riparian areas of the PUBs consisted of tall fescue, white clover, tall buttercup, little blue stem, annual bluegrass, scutch grass, curly dock, spike rush, an unknown sedge (*Carex sp.*), *Rubus sp.*, Osage orange, and black willow. Representative photographs of wetland habitat community types are provided in Appendix C.

Palustrine forested wetlands (PFO) accounted for less than one percent (2.8 acres) of the total Survey Area. The common overstory and understory vegetation consists of sugarberry, osage orange, green ash, American sycamore, red maple, Chinese privet; and common vines, shrubs, and herbaceous vegetation consisting of poison ivy, saw-tooth blackberry, butterweed, alligator weed, Virginia creeper, Virginia wild rye, and soft rush.

Palustrine emergent wetlands (PEM) accounted for less than one percent (11.89 acres) of the total Survey Area. Common vegetation observed within the emergent wetlands included black willow, butterweed, shallow sedge, rough cockleburr, woolgrass (*Scirpus cyperinus*), curly dock, yellow nutsedge, blackberries, soft rush, and alligator weed.

Palustrine scrub/shrub wetlands (PSS) accounted for less than one percent (2.49 acres) of the total Survey Area. Common vegetation observed within the scrub/shrub wetlands included, green ash, water locust, butterweed, curly dock, and soft rush.

Several observations of garter snakes, red-eared sliders (*Trachemys scripta*), upland chorus frogs, *a* cottonmouth (*Agkistrodon piscivorus*), and banded water snake (*Nerodia fasciata*) were documented in wetlands within the Survey Area. Wading birds included great blue heron (*Ardea herodius*) and great egret (*Ardea alba*), which were observed near the norther portion of the Survey Area near the aquaculture ponds (across from Gilmer Wilburn Rd). Mallard ducks (*Anas platyrhynchos*) and gambusia sp. were observed in many of the ponds within the Survey Area. Evidence of North American beaver (*Castor canadensis*) activity was also observed in several locations within the Survey Area.

#### 3.1.4 **Streams**

Of the 59 streams mapped within the Survey Area, there were 6 perennial streams (totaling 25,525 linear feet), 22 intermittent streams (totaling 13,138 linear feet), and 31 ephemeral streams (totaling 12,295

linear feet). At the time of survey, most ephemeral streams did not contain water. Most perennial streams had limited flow and fragmented sections of water throughout the reaches.

Observations of gambusia were documented during field surveys. However, no fish or macroinvertebrate sampling was conducted. No other observations of aquatic species were documented. Representative photographs of each stream flow regime type are provided in Appendix C.

# 3.2 Protected Species Information

# 3.2.1 Species Protected under the Endangered Species Act

The USFWS IPaC for the Survey Area was reviewed to determine the potential occurrence of species listed by the USFWS as threatened, endangered, or species of special concern (**Appendix A**). Three federally listed species were identified as potentially occurring within the Survey Area: the northern long-eared bat (*Myotis septentrionalis*), the wood stork (*Mycteria americana*), and Price's potato-bean (*Apios priceana*). Critical habitat for federally protected species has not been designated within Lowndes County, Mississippi for any of these species.

# 3.2.1.1 Wood Stork

The wood stork (*Mycteria americana*) is a federally threatened bird found primarily in freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches. Wood storks typically develop roosting sites in medium to tall trees that occur in stands either in swamps or on islands surrounded by relatively broad expanses of open water; primarily bald cypress (*Taxodium distichum*) swamps. Nesting usually occurs in bald cypress trees in swamps. According to Cornell Lab's wood stork sightings map from 2016-2021, there have historically been sightings near the Survey Area; however, wood stork is primarily a non-migratory bird in the United States and typically occurs in South Carolina, Georgia, and Florida (Cornell, 2021). No nesting records are known from this area of Mississippi (USFWS 2020b). Wetlands within the Survey Area were small, isolated, and low quality (Appendix C). Therefore, no suitable roosting habitat for wood storks was observed within the Survey Area; however, marginal foraging habitat for the species was observed within most of the open waters throughout the Survey Area. Proposed solar array layouts would be designed to minimize impacts to wetlands that could potentially provide wood stork foraging habitat.

# 3.2.1.2 Northern Long-eared Bat

The northern long-eared bat is a federally threatened mammal that roosts in tree cavities and under exfoliating bark, but this species has also been found in buildings and behind window shutters. During the

winter, northern long-eared bats hibernate in tight crevices in caves and mines. Foraging is done primarily on forested hillsides and ridges.

#### Potentially Suitable Northern Long-Eared Bat Summer Roosting Habitat

In the USFWS's Range-Wide Indiana Bat Survey Guidelines (USFWS, 2019), suitable summer habitat for NLEB is defined as follows:

"Suitable summer habitat consists of a wide variety of forested/wooded habitats where they roost, forage, and travel and may also include adjacent and interspersed non-forested habitats such as emergent wetlands and adjacent edges of agricultural fields, old fields and pastures. This includes forests and woodlots containing potential roosts (i.e., live trees and/or snags ≥3 inches diameter at breast height [dbh] that have exfoliating bark, cracks, crevices, and/or cavities), as well as linear features such as fencerows, riparian forests, and other wooded corridors. These wooded areas may be dense or loose aggregates of trees with variable amounts of canopy closure. Individual trees may be considered suitable habitat when they exhibit characteristics of suitable roost trees and are within 1,000 feet of forested/wooded habitat. NLEB has also been observed roosting in human-made structures, such as buildings, barns, bridges, and bat houses; therefore, these structures should also be considered potential summer habitat. NLEBs typically occupy their summer habitat from mid-May through mid-August each year; and the species may arrive or leave some time before or after this period.

#### Examples of unsuitable habitat:

- Individual trees that are greater than 1,000 feet from forested/wooded areas.
- Trees found in highly developed urban areas (e.g., street trees, downtown areas); and
- A pure stand of less than 3-inch dbh trees that are not mixed with larger trees."

Suitable roosting habitat for the NLEB was observed within most of the forested areas, and suitable foraging habitat was observed within the perennial stream corridors throughout the Survey Area (**Habitat Assessment Map; Appendix B**). Suitable roosting habitat was observed as mixed hardwood forest with an even representation of shagbark hickory with exfoliating bark and scattered snags throughout. No caves or mines were identified on the Project Site during field surveys. Additionally, in speaking with landowners during site visits, no known caves or mines exist in proximity to Survey Area. Photographs representing examples of suitable bat habitat with the Survey Area are provided in (**Appendix C**). According to TVA's natural heritage database, there are no known records of NLEB within ten miles of the Survey Area.

#### 3.2.1.3 Price's Potato-bean

Price's potato-bean (*Apios priceana*) is a federally threatened plant that prefers lightly disturbed areas such as forest openings, wood edges and where bluffs descend to streams. It also grows along highway ROWs and powerline corridors. There are no bluffs that descend to streams within this Survey Area. One area of potentially suitable habitat for the species was observed where there is minor north facing slope towards a perennial stream shown in (Habitat Assessment Map; Appendix B). However, no individual Price's potato-bean were observed during field surveys. If identified suitable habitat for Price's potato-bean cannot be avoided/removed from the project area, species specific surveys would be conducted by a qualified botanist during the USFWS' recommended optimal survey window (late July – August).

# 3.2.2 Mississippi Natural Heritage Program State Protected Species and Species of Special Concern

The Mississippi Natural Heritage Program is managed under the MDWFP, Museum of Natural Science. Consultation was initiated with the MDWFP on February 25, 2021, and database results were issued on March 18, 2021, which indicate occurrences of state listed species or species of special concern that are known to occur within two miles of the Survey Area. Results of that correspondence are in Appendix A. According to the results, one state-listed species, the federally threatened wood stork, and 15 state ranked species of special concern have potential to occur within the Survey Area. Furthermore, MDWFP concluded that if best management practices are properly implemented, monitored, and maintained, the proposed project likely poses no threat to listed species or their habitats. Each of the species and a description of its preferred habitat is provided below. The state protected species identified are listed in Table 3-1 and discussed in further detail below.

Table 3-1: Mississippi Department of Wildlife, Fisheries, and Parks - Natural Heritage Program - Protected Species and Species of Special Concern Potentially Occurring within two miles of the Golden Triangle II Project Survey Area

Common Name	Scientific Name	Federal Status	State Status	Preferred Habitat Description	
Eastern Spotted Skunk	Spilogale putorius	-	<b>S</b> 1	Open prairies, brushy lands, and cultivated areas (MDC 2021)	
Wood Stork	Mycteria americana	LT	LE	Freshwater wetlands, including ponds, bayheads, flooded pastures, oxbow lakes, and ditches (Cornell 2019a)	
Painted Bunting	Passerina ciris	-	S3B	Dense weedy habitats as well as the dense understory of semi-open forests (Cornell 2019b)	
Scissor-tail Flycatcher	Tyrannus forficatus	-	S1B; S1N	Open habitats in the southern Great Plains and sout Texas, often around sparsely scattered trees or utilit lines (Cornell 2019c)	
Big-head Evax	Evax <i>Prolifera</i>	-	S1	Meadows, grasslands, sagebrush steppe, and plains (MNHP 2021)	

Common Name	Scientific Name	Federal Status	State Status	Preferred Habitat Description	
Slender Sedge	Carex gracilescens	-	S1	Mesic deciduous forest; often in sandy, rocky, and sometimes calcareous soils (WDNR 2020)	
Stiff Greenthread	Thelesperma filifolium	-	S1	Disturbed sites in dry sandy or gravely soil (UTA 2018)	
Nebraska Sedge	Carex jamesii	-	S1S2	rich mesic woodlands, wooded slopes, wooded groves (where the ground vegetation is infrequently mowed underneath large trees), and edges of woodland paths (Hilty 2020a)	
Big Shellbark Hickory	Carya laciniosa	-	S2	Rich alluvium of floodplains and rich mesic soil along riverbanks and marshes (NYNHP 2021)	
Bur Oak	Quercus macrocarpa	-	S2	Moist woodlands and bottomland forests to prairies and sandhills (USFS 2021)	
Ohio Buckeye	Aesculus glabra	-	S2	Moist to mesic deciduous woodlands, wooded valleys along rivers, and rocky wooded slopes in sheltered areas (Hilty 2020b)	
Prairie Parsley	Polytaenia nuttallii	-	S2	Upland prairies, hill prairies, limestone glades, che glades, thinly wooded bluffs, and savannas (Hilty 2018).	
Prairie Pleatleaf	Nemastylis geminiflora	1	S2	Glades, prairies, and rocky slopes, on calcareous substrates – Occasionally on glades in the shade of eastern red cedars or Ashe's junipers (MDC 2020)	
Small-toothed Sedge	Carex microdonta	ı	S2	Dry to moist, calcareous substrates in open rocky of wet prairies, swales, seeps, and ditches - Relic patches of Blackland prairies, limestone glades, and chalk openings (FNA 2020)	
Mead's Sedge	Carex meadii	-	S3S4	moist to dry black soil prairies, dolomite prairies, hill prairies, savannas with sparse ground vegetation, sedge meadows, limestone glades, and areas along railroads (Hilty 2019a)	
Carya myristiciformis	Nutmeg Hickory	-	S3S4	Banks of rivers and swamps in rich moist soils, occasionally on higher ground and often on limestone (PFAF 2021)	

Key: Statuses are LE= Listed Endangered, LT= Listed Threatened

State Rank S1 - Critically imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

Sources: USFWS, 2021 and Mississippi Natural Heritage Program, 2021.

### 3.2.3 Bald Eagles and Migratory Birds

In Mississippi, the bald eagle is protected under the Bald and Golden Eagle Protection Act (BGEPA) and the Migratory Bird Protection Act (MBTA). No bald eagles or nests were observed during the environmental field surveys within the Survey Area. One potential bald eagle nest was observed from a distance in the adjacent parcel outside of the Survey Area, approximately 1,700 ft away but could not be confirmed. The potential nest is within private property outside of the Survey Area. Photographs were attempted but were unable to provide value due to distance and obstruction. Status of the nest is unknown,

S2 - Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation. S3 - Rare or uncommon in Mississippi (on the order of 21 to 100 occurrences).

and no individuals were observed. Bald eagles would be most likely found near the Project Site in December and January. There are aquaculture ponds directly north and south of the Project Site which could provide a source of food for the bald eagle. No nests were observed within the Survey Area during field surveys; however, bald eagles could exist near the Project Site due to the aquaculture ponds. According to TVA's natural heritage data, there are no known bald eagles within three miles of the Survey Area.

According to the USWFS IPaC report, there are no migratory birds of concern protected under the MBTA within the vicinity of the Survey Area (USFWS 2021). Migratory birds of conservation concern identified in TVA's natural heritage data include American kestrel, bald eagle, cerulean warbler, dunlin, Kentucky warbler, lesser yellowlegs, marbled godwit, prothonotary warbler, red-headed woodpecker, rusty blackbird, short-billed dowitcher, swallow-tailed kite, willet, and wood thrush. Additionally, MDWFP indicated that scissor-tail flycatcher and the painted bunting could also occur within two miles of the Survey Area. Both species are state species of concern.

The American Kestrel is a small member of the falcon family and prefers habitats consisting of open farmland, wood edges, and cities. It breeds from April through August most likely in areas outside the Project Site (Cornell 2019d). It is most commonly found in Lowndes County, Mississippi during December, and January. No individuals or nests were observed during field surveys; however suitable nesting and foraging habitat does occur within the Survey Area.

Ospreys (*Pandion haliaetus*) are large birds of prey that typically nest in elevated nest sites near rivers, lakes, reservoirs, lagoons, swamps, and marshes with an adequate supply of accessible fish within about twelve miles of the nest (Cornell 2019e). According to TVA's natural heritage data, there is a record from 2011 of an osprey's nest approximately 2.7 miles from the Survey Area. No nests were observed during field surveys; however, osprey could exist near the Survey Area due to the proximity of the aquaculture ponds.

Lesser yellowlegs (*Tringa flavipes*) is a small "piper" bird that breeds in meadows and open woodlands in Canada. In North America, lesser yellowlegs typically occurs in marshes, shallow wetlands, shorelines, and flooded fields during migration. Most often, they prefer vegetated wetlands rather than bare habitats (Cornell 2019b). Marginal foraging habitat is present at PEM wetlands throughout the Project Site. PEM wetlands within the Survey area were low quality, isolated, and are regularly disturbed during crop and hay harvesting (Appendix C).

Marbled godwit (*Limosa fedoa*) is a small shorebird with a two-toned, long, slightly upcurved bill that breeds in the shortgrass prairies near wetlands in Canada and the northern central U.S. While migrating south, marbled godwits typically inhabit native grass prairies with green needle grass, western wheatgrass, blue gram, needle-and-thread, and blue stem. Wintering grounds typically consist of coastal mudflats, estuaries, and sandy beaches (Cornell 2019c). The migration corridor for the marbled godwit is further west of the Survey Area, closer to Texas and Arkansas. Due to lack of suitable habitat and the location of the Survey Area outside of the migration corridor, marbled godwit is unlikely to occur within the Survey Area.

Red-headed woodpecker (*Melanerpes erythrocephalus*) is a medium-sized woodpecker with a bright red head, white buff, and black back. Red-headed woodpeckers breed in river bottoms, beaver swamps, burned areas, recent clearing, deciduous woodlands with oak or beech and groves of dead or dying trees. Dead or partially dead trees are important for nest cavities in areas where they breed. In Mississippi, red-headed woodpeckers typically inhabit pine and pine-oak forests (Cornell 2019d). Most of the undisturbed forested areas within the Survey Area would provide suitable habitat for red-headed woodpeckers.

As its name entails, the rusty blackbird (*Zanata Canadiense*) has distinctively rusty feather edges and pale-yellow eyes. Rusty blackbirds typically breed in wet forests, swamps, bogs, and beaver ponds. They also winter in swamps, woodlands, and pond edges. This species is considered a common bird in steep decline due to habitat loss (Cornell 2019e). Potentially suitable habitat for rusty blackbird is present near the pond edges and riparian woodlands to the open ponds.

Short billed dowitcher (*Limnodromus griseus*) is a full-bodied orange, brown, and golden shorebird with a long bill used for rhythmic probing of mudflats and flooded field for food. Short-billed dowitchers breed in the taiga shield ecotone, nest in wetlands usually near edges of bogs, small lakes, or wet meadows. During winter months, short-billed dowitchers are founds in saltwater and brackish environments such as estuaries and lagoon with tidal influences (Cornell 2019f). Marginal foraging habitat for migrant short-billed dowitchers is present in some of the shallow ponds and potentially flooded farm fields when excessive precipitation occurs.

Willet (*Tringa semipalmata*) is a straight billed, mottled brown, shorebird with distinct wing markings and a piercing call. Willets typically inhabit open beaches, marshes, mudflats, and rocky coastal zones. During the breeding season eastern willets are typically found in saltmarshes, barrier islands, and barrier beaches (Cornell 2019g). Marginal foraging habitat for migrant willets is present in some of the shallow ponds and potentially flooded farm fields from excessive precipitation.

MS Solar 6, LLC 3-9 Burns & McDonnell

MDWFP's Natural Heritage Data also identified two migratory birds with potential to occur within two miles of the Survey Area:

Scissor tail flycatcher (*Tyrannus forficatus*) was identified by the MDWFP as potentially occurring within two miles of the Survey Area. Scissor tail flycatchers prefer open habitats in the southern Great Plains and south Texas, often around sparsely scattered trees, or utility lines (Cornell 2019c). Open pasture within the Survey Area provides marginal foraging habitat for the scissor-tailed flycatcher.

Painted Bunting (*Passerina ciris*) was also identified by the MDWFP as potentially occurring within two miles of the Survey Area. Painted bunting prefers dense weedy habitats as well as the dense understory of semi open forests (Cornell 2019b). Although they are typically more often found further south and west, most of the forested areas within the Survey Area provide foraging habitat.

If tree clearing activities associated with construction of the Project overlap with the primary nesting season (March 15 – September 15), short-term inadvertent impacts could occur on bird species that nest in or near the construction areas. To the extent practicable, Project construction will be scheduled to minimize potential effects to bird species by minimizing clearing activities within forested areas during nesting season. Other mitigation measures MS Solar 6, LLC plans to implement include:

- Designing Project facilities to avoid sensitive resources where possible.
- Maximizing locations where the Project utilizes agricultural areas.
- Limiting the construction and operation workspaces to the minimum necessary.
- Conducting mitigation for effects to sensitive resources (e.g., wetlands) through agency permit conditions.
- Avoiding forested areas, to the greatest extent practicable.
- Minimizing routine mowing/maintenance during the bird nesting season once the Project is complete (generally March 15 through September 15 in the general Project vicinity) to the extent feasible.

On Dec. 22, 2017, the U.S. Department of the Interior (DOI) revised its guidance on incidental take of migratory birds in Memorandum M-370501, which specifies that incidental take prohibitions apply only to actions that have *as their purpose* the taking or killing of migratory birds. Under the Biden administration, the Department of the Interior withdrew its 2017 legal opinion which preceded and form the basis of the rule. The new administration may return policies initiated under the Obama administration, which considered a proposal to develop an incidental take program. Because MS Solar 6, LLC's purpose is the lawful construction of a clean energy facility, and not the intentional take of

migratory birds, MS Solar 6, LLC does not anticipate further coordination with USFWS regarding migratory birds, however, legal developments to MBTA should be monitored.

#### 3.2.4 TVA Natural Heritage Database Results

Terrestrial species documented in TVA's natural heritage data included northern long-eared bat, wood stork, osprey and bald eagle which are discussed above. Mitchell's satyr (*Neonnympha mitchellii*) was also identified has having the potential to occur within the Survey Area, however, that species has not been documented within three miles of the project area or from Lowndes County, Mississippi. Additionally, Mitchell's satyr habitat is restricted to fens which are rare wetlands comprised of low nutrient systems that receive carbonate-rich ground water from seeps and springs which do not occur within the Survey Area (USFWS 2021b).

As of 2018, there are 495 plants of special concern in Mississippi. The designation of plants of special concern in Mississippi was developed for flagging sensitive species that may be adversely affected by proposed projects, determining protection priorities of natural areas that contain special plants, and determining priorities for inventory and protection for special plants to include proposing special plants for federal protection (Mississippi Natural Heritage Program, 2018).

Twenty-eight rare plant species were included in the results of the TVA Heritage Database search as potentially occurring within five miles of the Survey Area. Because MS Solar 6 is avoiding most of the wetland and stream riparian areas and chalk outcroppings and associated buffers from the construction and operation footprint, suitable habitat only exists for 12 of these plant species within the Project Site. The table below provides additional details for these 12 species.

Table 3-2: TVA Natural Heritage Database - Plants Potentially Occurring within the Survey Area

Common Name	Scientific Name	Rank	Preferred Habitat Description
Ridgestem false foxglove	Agalinis oligophylla	S2	Typically found from east Texas to southwest Louisiana on coastal plains primarily in prairies – Prefers moist to dry sand or clay soils and also found in pine hardwood forest and pine savannahs (USGS 2020)
Wild hyacinth	Camassia scilloides	S2	Black soil prairies, moist savannas, moist open woodlands along stream banks, rocky wooded slopes, and limestone glades - Typically prefer moist conditions to full sun to light shade in rich loamy soil (Hilty 2020c)
James's Sedge	Carex jamesii	S1S2	Mesic woodlands, wooded slopes, wooded groves, and edges of woodland paths - May also occur in upland woods and swampy woodlands and occasionally found in degraded woodland habitats (Hilty 2020a)

MS Solar 6, LLC 3-11 Burns & McDonnell

Common Name	Scientific Name	Rank	Preferred Habitat Description
Small-toothed sedge	Carex microdonta	<b>S</b> 3	Dry to moist, calcareous substrates in open rocky or wet prairies, swales, seeps, and ditches - Relic patches of Blackland prairies, limestone glades, and chalk openings (FNA 2020)
Kingnut Hickory	Carya laciniosa	<b>S</b> 2	Bottomland woodlands, upland woodlands, swamps, savannas, and limestone glades. Usually not far from rivers (Hilty 2020d)
Scarlet Indian- paintbrush	Castilleja coccinea	<b>S</b> 1	Sandy soils in prairies and open woods (USFS 2020)
Pumpkin ash	Fraximus profunda	<b>S</b> 3	Moist wet conditions often in swamps or floodplains, wet bottomlands, river valleys, and low areas. Often in bald cypress swamps, cottonwood, and tupelo swamps (MBG 2020a)
Canada moonseed	Menispermum canadense	<b>S</b> 3	Deciduous woodlands, woodland borders, thickets, semi- shaded riverbanks, cleared powerline ROWs, overgrown fencerows, and hedges (Hilty 2020e)
Prairie pleatleaf	Nemastylis geminiflora	S2	Glades, prairies, and rocky slopes, on calcareous substrates – Occasionally on glades in the shade of eastern red cedars or Ashe's junipers (MDC 2020)
Bur oak	Quercus macrocarpa	S2	Bottomland soils in woodland and stream valleys (MBG 2020b)
Heath aster	Symphyotrichum ericoides	S2	Mesic to dry black soil prairies, gravel prairies, dolomite prairies, hill prairies, savannas, openings in dry rocky forests, limestone glades, roadsides and railroad sides, and pastures (Hilty, 2020f)
Southern Meadow Rue	Thalictrum debile	S1S2	Floodplain forests over calcareous substrates – prefers rich, rocky limestone woods often near streams (Chafin 2007).
Great Plains Ladies'-tresses	Spiranthes magnicamporum	S2	Open prairie and higher pH soils in both low and, moist areas and higher, gravelly sites (Minnesotawildflowers 2021)

S1 = Critically Imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation.

Source: TVA's Heritage Database "Botany within 5 Miles"

During field surveys, one species of state listed tree was observed in the forested along the fence line of the southeast border of the Survey Area: bur oak (*Quercus macrocarpa*) (state ranked S2). Bur oak observations were limited to two individuals in lower elevations along a stream in the same general area. The two individuals were mature trees and had a diameter at breast height (dbh) of three to four feet. Representative photos are provided under Appendix C.

MS Solar 6, LLC 3-12 Burns & McDonnell

S2 = Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.

S1S2 = Rank range between S1 and S2.

S3 = Vulnerable in Mississippi due to a restricted range (on the order of 21 to 100 occurrences), relatively few populations or occurrences, recent and widespread declines, or other factors making it vulnerable to extirpation.

Two areas of open chalk outcrops and associated buffers were also mapped in the southwest portion of the Survey Areas as areas that could potentially support sensitive plant species and will be recommended as avoidance areas. No rare plants were observed at or near the chalk outcrops during surveys.

Table 3-4: TVA Natural Heritage Database – Aquatic Species Potentially Occurring within 10-miles of Survey Area.

Common Name	Scientific Name	Rank/Status	Preferred Habitat Description
Blue Sucker	Cycleptus elongatus	S3	Large rivers with deep channels and swift currents that have substrates consisting of sand, gravel or rubble bottoms. (MDNR 2021a)
Crystal Darter	Crystallaria asprella	S1/LE	Medium to large rivers, usually with moderate to swift currents and clean sandy or gravel bottoms (MDNR 2021b)
Cylinder Elimia	Elimia cylindracea	S2	Riffle habitats often in areas with rocky legdes (NatureServe 2021)
Flat Pigtoe	Pleurobema marshalli	SX/LE	Medium to large rivers with moderate to fast current and sandy gravel to gravel cobble substrates and clean water (NatureServe 2021b)
Frecklebelly Madtom	Noturus munitus	S2/LE	Fast moving streams associated with larger rivers that provide firm gravel substrates with pebbles and rock (USFWS 2020).
Freckled Darter	Percina lenticula	S2	Medium to large rivers with deep and swift currents (NatureServe 2021c)
Heavy Pigtoe	Pleurobema taitianum	SX/LE	Riffles and shoals of small to large rivers with sandy gravel to gravel-cobble substrates and moderate to fast currents (NatureServe 2021d).
Southern Clubshell	Pleurobema decisum	S1	Highly oxygenated small to medium rivers with loose sand and gravel substrates (NatureServe 2021e).
Southern Combshell	Epioblasma penita	S1/LE	Medium sized rivers with high gradient and riffles with sandy gravel to gravel cobble substrates in moderate to swift currents (NatureServe 2021f).
Stirrupshell	Quadrula stapes	SX/LE	Medium to large rivers with moderate to fast current and sandy gravel to gravel cobble substrates and clean water (NatureServe 2021g)
Striped Bass	Morone saxatilis	SH	Coastal estuaries; migrates upstream to medium to large rivers during spawning typically within 6 km of the shore (NatureServe 2021h).
White Heelsplitter	Lasmigona complanata	S3	Rivers and stream of various sizes with substrates of gravel, sand, or mud (NatureServe 2021i).

MS Solar 6, LLC 3-13 Burns & McDonnell

Common Name	Scientific Name	Rank/Status	Preferred Habitat Description

- S1 = Critically Imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or very few remaining individuals or acres) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation.
- S2 = Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.
- S3 = Vulnerable in Mississippi due to a restricted range (on the order of 21 to 100 occurrences), relatively few populations or occurrences, recent and widespread declines, or other factors making it vulnerable to extirpation.
- SH = Possibly Extirpated Known from only historical records in Mississippi, but still some home of rediscovery. There is evidence that the species or ecosystem may no longer be present in the jurisdiction, but not enough to state this with certainty.
- SX = Presumed Extirpated Species or ecosystem is believed to be extirpated from Mississippi. Not located despite intensive searches of historical sites and other appropriate habitat, and virtually no likelihood that it will be rediscovered.

The six perennial streams identified within the Survey Area had limited flow and fragmented sections of water throughout the reaches. Perennial streams within the Survey Area were characterized by the presence of a substantial volume of flow at the time of the site visit as well as secondary characteristics such as observance of fish and rooted aquatic fauna, indicating that water flows year-round. Perennial streams were approximately 6 to 25 feet in width at the OHWM with bank heights ranging from 3 to 25 feet. At the time of the delineation, the depth of water observed was 0.5 to 2.5 feet. Perennial streams flowed through upland fields, agricultural fields, and wooded riparian areas.

Observations of gambusia were documented during field surveys. However, no fish or macroinvertebrate sampling was conducted. No other observations of aquatic species were documented. Representative photographs of each stream flow regime type are provided in Appendix C.

With the exception of the white heelsplitter (S3), all aquatic species identified in the TVA Natural Heritage Database as potentially occurring within 10-miles of the Survey Area require aquatic habitats primarily comprised of medium to large rivers with moderate to fast moving currents. Perennial streams within the Survey Area do not meet habitat requirements for these species. White heelsplitters can be found in a variety of freshwater settings and perennial stream within the Survey Area may provide marginal habitat for this species. Additionally, proposed solar array layouts would be designed to avoid perennial stream and their buffers. Therefore, impacts to aquatic species within the Survey Area are not expected.

#### 4.0 CONCLUSIONS

The wood stork is federally protected under the ESA. Suitable roosting habitat for the wood stork does not exist within the Survey Area for the Project. However, suitable foraging habitat may be present near open water and large inundated wetlands. There are also large aquaculture/fish farms both north and south of the Project that may attract foraging wood storks. The Project will not affect fish farms or large open waters outside the immediate Project limits. For these reasons, the Project will have *no effect* on wood storks, and no further consultation is anticipated.

Only a few populations of Price's potato-bean are known to exist today because its distinct habitat requirements. Marginally suitable habitat for this plant was observed in the Survey Area. However, during surveys, no occurrences were identified within the Survey Area. Additionally, avoidance buffers will likely be places around intermittent and perennial streams and their associated riparian areas further protecting rare plants potentially inhabiting those areas. For these reasons, the Project will have *no effect* on Price's potato-bean, and no further consultation is anticipated.

Suitable roosting and foraging habitat for the NLEB was identified within the Survey Area. Based on preliminary designs, the Project is expected to result in up to 368 acres of forest clearing within an area identified as potentially suitable roosting habitat. Appendix B contains a figure that depicts potentially suitable NLEB roosting and foraging habitat as well as the specific area proposed for tree clearing. However, acoustic surveys were conducted at 11 sites (a total of 26 detector-nights) within the Survey Area in accordance with USFWS Guidelines (Appendix D). NLEB was determined to be likely absent based on automated identification results. Therefore, the Project is expected to have *no effect* to the northern long-eared bat.

According to the Mississippi Natural Heritage Program results, one state-listed species, the federally threatened wood stork, and 15 state ranked species of special concern have potential to occur within the Survey Area. Furthermore, MDWFP concluded that if best management practices are properly implemented, monitored, and maintained, the proposed project will have *no effect* to state protected species or their habitats.

Based on the natural heritage data provided by TVA, an osprey, and occurrences of 28 rare plants, and twelve aquatic species, have been known to occur within three and five miles of the Survey Area, respectively. Of those species, bur oak was the only rare species to be observed during field surveys and was limited to two individuals in the southeast corner of the Survey Area. Due to their location in a perennial stream riparian zone and proximity to the parcel boundary, these trees may easily be avoided.

MS Solar 6, LLC 4-1 Burns & McDonnell

#### 5.0 REFERENCES

- Chafin, Linda G. 2007. Field Guide to Rare Plants of Georgia. State Botanical Garden of Georgia, Athens, Georgia.
- Cornell University 2019a. The Cornell Lab. All About Birds. Wood Stork Life History. <a href="https://www.allaboutbirds.org/guide/Wood\_Stork/lifehistory">https://www.allaboutbirds.org/guide/Wood\_Stork/lifehistory</a>. Accessed March 1, 2021
- Cornell University 2019b. The Cornell Lab. All About Birds. Painted Bunting Life History. <a href="https://www.allaboutbirds.org/guide/Painted Bunting/lifehistory">https://www.allaboutbirds.org/guide/Painted Bunting/lifehistory</a>. Accessed April 27, 2021.
- Cornell University 2019c. The Cornell Lab. All About Birds. Scissor-tailed flycatcher Life History. <a href="https://www.allaboutbirds.org/guide/Scissor-tailed\_flycatcher/lifehistory">https://www.allaboutbirds.org/guide/Scissor-tailed\_flycatcher/lifehistory</a>. Accessed April 27, 2021.
- Cornell University 2019d. The Cornell Lab. All About Birds. American Kestrel Life History. <a href="https://www.allaboutbirds.org/guide/American\_Kestrel/lifehistory">https://www.allaboutbirds.org/guide/American\_Kestrel/lifehistory</a>. Accessed October 21, 2020.
- Cornell University 2019e. The Cornell Lab. All About Birds. Osprey Life History. <a href="https://www.allaboutbirds.org/guide/Osprey/lifehistory">https://www.allaboutbirds.org/guide/Osprey/lifehistory</a>. Accessed April 27, 2021.
- Cornell University. 2019b. The Cornell Lab. All About Birds. Lesser Yellowlegs Life History. <a href="https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory">https://www.allaboutbirds.org/guide/Lesser\_Yellowlegs/lifehistory</a>. Accessed October 20, 2020.
- Cornell University 2019c. The Cornell Lab. All About Birds. Marbled Godwit Life History. https://www.allaboutbirds.org/guide/Marbled\_Godwit/lifehistory. Accessed October 20, 2020.
- Cornell University 2019d. The Cornell Lab. All About Birds. Red-headed Woodpecker Life History. <a href="https://www.allaboutbirds.org/guide/Red-headed Woodpecker/lifehistory">https://www.allaboutbirds.org/guide/Red-headed Woodpecker/lifehistory</a>. Accessed October 20, 2020.
- Cornell University 2019e. The Cornell Lab. All About Birds. Rusty Blackbird Life History. https://www.allaboutbirds.org/guide/Rusty\_Blackbird/lifehistory. Accessed October 20, 2020.
- Cornell University 2019f. The Cornell Lab. All About Birds. Short-billed Dowitcher Life History. <a href="https://www.allaboutbirds.org/guide/Short-billed\_Dowitcher/lifehistory">https://www.allaboutbirds.org/guide/Short-billed\_Dowitcher/lifehistory</a>. Accessed October 20, 2020.
- Cornell University 2020g. The Cornell Lab. All About Birds. Willet Life History. <a href="https://www.allaboutbirds.org/guide/Willet/lifehistory">https://www.allaboutbirds.org/guide/Willet/lifehistory</a>. Accessed October 20, 2020.
- Flora of North America (FNA). 2020. *Carex microdonta*. www.efloras.org/florataxon.aspx?flora\_id=1&taxon\_id=242357334. Accessed August 31, 2020.
- Hilty, John. 2018. Illinois Wildflowers: Prairie Parsely (*Polytaenia nuttallii*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/pr\_parsley.html">https://www.illinoiswildflowers.info/prairie/plantx/pr\_parsley.html</a>. Accessed April 26, 2021.
- Hilty, John. 2019a. Illinois Wildflowers: Mead's Sedge (*Carex meadii*). https://www.illinoiswildflowers.info/grasses/plants/mead\_sedge.htm. Accessed April 26, 2021.

MS Solar 6, LLC 5-1 Burns & McDonnell

- Hilty, John. 2020a. Illinois Wildflowers: James' Sedge (*Carex jamesii*).

  <a href="https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm">https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020b. Illinois Wildflowers: Ohio Buckeye (*Aesculus glabra*). <a href="https://www.illinoiswildflowers.info/trees/tree">https://www.illinoiswildflowers.info/trees/tree</a> index.htm#oh buckeye. Accessed April 26, 2021.
- Hilty, John. 2020c. Illinois Wildflowers: Wild Hyacinth (*Camassia scilloides*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm">https://www.illinoiswildflowers.info/prairie/plantx/wild\_hyacinth.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020d. Illinois Wildflowers: Kingnut Hickory (*Carya laciniosa*). https://www.illinoiswildflowers.info/trees/plants/kn hickory.htm. Accessed August 31, 2020.
- Hilty, John. 2020e. Illinois Wildflowers: Canada Moonseed (*Menispermum canadense*). <a href="https://www.illinoiswildflowers.info/prairie/plants/canada\_moonseed.htm">https://www.illinoiswildflowers.info/prairie/plants/canada\_moonseed.htm</a>. Accessed August 31, 2020.
- Hilty, John. 2020f. Illinois Wildflowers: Heath Aster (*Symphyotrichum ericoides*). <a href="https://www.illinoiswildflowers.info/prairie/plantx/hth\_asterx.htm">https://www.illinoiswildflowers.info/prairie/plantx/hth\_asterx.htm</a>. Accessed August 31, 2020.
- Minnesota Wildflowers. 2021. A Field Guide to Flora of Minnesota: Great Plains Ladies' tresses (*Spiranthes magnicamporum*). <a href="https://www.minnesotawildflowers.info/flower/great-plains-ladies-tresses">https://www.minnesotawildflowers.info/flower/great-plains-ladies-tresses</a>. Accessed April 27, 2021.
- Minnesota Department of Natural Resources. 2021. Species Profile: *Cycleptus elongatus*. <a href="https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AFCJC040">https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AFCJC040</a> 10. Accessed December 7, 2021.
- Minnesota Department of Natural Resources. 2021. Species Profile: *Crystallaria asprella*. https://www.dnr.state.mn.us/rsg/profile.html?action=elementDetail&selectedElement=AFCQC01 010. Accessed December 7, 2021. Mississippi Museum of Natural Science. 2014. Endangered Species of Mississippi. Mississippi Department of Wildlife, Fisheries, and Parks, Mississippi Museum of Natural Science, Jackson, MS. <a href="https://www.mdwfp.com/media/256526/endangered\_species\_of\_mississippi.pdf">https://www.mdwfp.com/media/256526/endangered\_species\_of\_mississippi.pdf</a>.
- Mississippi Natural Heritage Program. 2018. Listed Species of Mississippi. Museum of Natural Science, Mississippi Dept. of Wildlife, Fisheries, and Parks, Jackson, MS. 6 pp. 24 September 2018.
- Missouri Botanical Garden. 2020a. Plant Finder: *Fraxinus profunda*.

  <u>www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=c987</u>.

  Accessed August 31, 2020.
- Missouri Botanical Garden. 2020b. Plant Finder. *Quercus macrocarpa*. <a href="https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=a902">https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?kempercode=a902</a> . Accessed August 31, 2020.
- Missouri Department of Conservation (MDC). 2020. Discover Nature: Field Guide. Celestial Lily (*Nemastylis geminiflora*). <a href="https://nature.mdc.mo.gov/discover-nature/field-guide/celestial-lily">https://nature.mdc.mo.gov/discover-nature/field-guide/celestial-lily</a>. Accessed August 31, 2020

- Missouri Department of Conservation (MDC). 2021. Discover Nature: Field Guide. Eastern Spotted Skunk (*Spilogale putorius*). <a href="https://nature.mdc.mo.gov/discover-nature/field-guide/eastern-spotted-skunk">https://nature.mdc.mo.gov/discover-nature/field-guide/eastern-spotted-skunk</a>. Accessed April 25, 2021.
- Montana Natural Heritage Program. 2021. Big-head Evax *Evax Prolifera*. Montanta Field Guide. <a href="http://FieldGuide.mt.gov/speciesDetail.aspx?elcode=PDAST3S050">http://FieldGuide.mt.gov/speciesDetail.aspx?elcode=PDAST3S050</a>. Accessed. April 27, 2021.
- NatureServeExplorer. 2021a. Cylinder Elimia (*Elimia cylindrica*).

  <a href="https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.113362/Elimia\_cylindracea">https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.113362/Elimia\_cylindracea</a>.

  Accessed December 7, 2021.
- NatureServeExplorer. 2021b. Flat Pigtoe (*Pleurobema marshalli*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.114612/Pleurobema\_marshalli. Accessed December 7, 2021.
- NatureServeExplorer. 2021c. Freckled Darter (*Percina lenticula*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.116207/Pleurobema\_taitianum. Accessed December 7, 2021.
- NatureServeExplorer. 2021d. Heavy Pigtoe (*Pleurobema taitianum*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.101698/Percina\_lenticula. Accessed December 7, 2021.
- NatureServeExplorer. 2021e. Southern Clubshell (*Pleurobema decisum*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.1066298/Pleurobema\_decisum. Accessed December 7, 2021.
- NatureServeExplorer. 2021f. Southern Combshell (*Epioblasma penita*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.109895/Epioblasma\_penita. Accessed December 7, 2021.
- NatureServeExplorer. 2021g. Stirrupshell (*Quadrula stapes*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.107836/Theliderma\_stapes. Accessed December 7, 2021.
- NatureServeExplorer. 2021h. Striped Bass (*Morone saxatilis*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.104731/Morone\_saxatilis. Accessed December 7, 2021.
- NatureServeExplorer. 2021h. White Heelsplitter (*Lasmigona compressa*). https://explorer.natureserve.org/Taxon/ELEMENT\_GLOBAL.2.111988/Lasmigona\_compressa. Accessed December 7, 2021.
- New York Natural Heritage Program. 2021. Online Conservation Guide for *Carya laciniosa*. Available from: https://guides.nynhp.org/big-shellbark-hickory/. Accessed April 26, 2021.
- Plants for a Future Database (PFAF). 2021. Nutmeg Hickory (*Carya myristiciformes*). https://pfaf.org/user/Plant.aspx?LatinName=Carya+myristiciformis. Accessed April 26, 2021.
- U.S. Forest Service (USFS) 2021. Fire Effects Information System (FEIS): *Quercus macrocarpa*. https://www.fs.fed.us/database/feis/plants/tree/quemac/all.html. Accessed April 26, 2021.

MS Solar 6, LLC 5-3 Burns & McDonnell

- University of Texas Austin. 2018. Thelesperma filifolium (Stiff Greenhead). Native Plants of North America. Lady Bird Johnson Wildflower Center. https://www.wildflower.org/plants/result.php?id\_plant=thfi. Accessed April 26, 2021.
- U.S. Fish and Wildlife Service (USFWS). 2019. <u>Range-wide Indiana Bat Survey Guidelines</u>. April 2019. Available via the internet at: <a href="https://www.fws.gov/arkansas-es/docs/FINAL%202019%20Range-wide%20IBat%20Survey%20Guidelines%204.10.19.pdf">https://www.fws.gov/arkansas-es/docs/FINAL%202019%20Range-wide%20IBat%20Survey%20Guidelines%204.10.19.pdf</a>
- U.S. Fish and Wildlife Service (USFWS). 2020b. Midwest Endangered Species: C
- U.S. Fish and Wildlife Service (USFWS) 2020. Frecklebelly Madtom (*Noturus munitus*). https://www.fws.gov/southeast/wildlife/fishes/frecklebelly-madtom/. Accessed December 7, 2021
- U.S. Fish and Wildlife Service (USFWS). 2021a. IPaC Report for Golden Triangle Solar Project. Mississippi Ecological Services Field Office. Jackson, MS. April 23, 2020.
- U.S. Fish and Wildlife Service (USFWS). 2021b. Mitchell's Satyr Butterfly Fact Sheet.

  <a href="https://www.fws.gov/midwest/endangered/insects/misa/misa\_fctsht.html">https://www.fws.gov/midwest/endangered/insects/misa/misa\_fctsht.html</a>. Accessed December 6, 2021.
- U.S. Forest Service. 2020. Plant of the Week: Indian Paintbrush (*Castilleja coccinea*). <a href="https://www.fs.fed.us/wildflowers/plant-of-the-week/Castilleja-coccinea.shtml">https://www.fs.fed.us/wildflowers/plant-of-the-week/Castilleja-coccinea.shtml</a>. Accessed August 31, 2020.
- U.S. Geological Survey (USGS). 2020. Plants of Louisiana *Agalinis oligophylla*. <a href="https://warcapps.usgs.gov/PlantID/Species/Details/1985">https://warcapps.usgs.gov/PlantID/Species/Details/1985</a>. Accessed October 21, 2020.

MS Solar 6, LLC 5-4 Burns & McDonnell





# United States Department of the Interior



#### FISH AND WILDLIFE SERVICE

Mississippi Ecological Services Field Office 6578 Dogwood View Parkway, Suite A Jackson, MS 39213-7856

Phone: (601) 965-4900 Fax: (601) 965-4340 http://www.fws.gov/mississippiES/endsp.html

In Reply Refer To: April 20, 2021

Consultation Code: 04EM1000-2021-SLI-0745

Event Code: 04EM1000-2021-E-01680 Project Name: Golden Triangle II

Subject: List of threatened and endangered species that may occur in your proposed project

location or may be affected by your proposed project

#### To Whom It May Concern:

The enclosed species list identifies threatened, endangered, proposed and candidate species, as well as proposed and final designated critical habitat, that may occur within the boundary of your proposed project and/or may be affected by your proposed project. The species list fulfills the requirements of the U.S. Fish and Wildlife Service (Service) under section 7(c) of the Endangered Species Act (Act) of 1973, as amended (16 U.S.C. 1531 *et seq.*).

New information based on updated surveys, changes in the abundance and distribution of species, changed habitat conditions, or other factors could change this list. Please feel free to contact us if you need more current information or assistance regarding the potential impacts to federally proposed, listed, and candidate species and federally designated and proposed critical habitat. Please note that under 50 CFR 402.12(e) of the regulations implementing section 7 of the Act, the accuracy of this species list should be verified after 90 days. This verification can be completed formally or informally as desired. The Service recommends that verification be completed by visiting the ECOS-IPaC website at regular intervals during project planning and implementation for updates to species lists and information. An updated list may be requested through the ECOS-IPaC system by completing the same process used to receive the enclosed list.

The purpose of the Act is to provide a means whereby threatened and endangered species and the ecosystems upon which they depend may be conserved. Under sections 7(a)(1) and 7(a)(2) of the Act and its implementing regulations (50 CFR 402 *et seq.*), Federal agencies are required to utilize their authorities to carry out programs for the conservation of threatened and endangered species and to determine whether projects may affect threatened and endangered species and/or designated critical habitat.

A Biological Assessment is required for construction projects (or other undertakings having similar physical impacts) that are major Federal actions significantly affecting the quality of the human environment as defined in the National Environmental Policy Act (42 U.S.C. 4332(2) (c)). For projects other than major construction activities, the Service suggests that a biological evaluation similar to a Biological Assessment be prepared to determine whether the project may affect listed or proposed species and/or designated or proposed critical habitat. Recommended contents of a Biological Assessment are described at 50 CFR 402.12.

If a Federal agency determines, based on the Biological Assessment or biological evaluation, that listed species and/or designated critical habitat may be affected by the proposed project, the agency is required to consult with the Service pursuant to 50 CFR 402. In addition, the Service recommends that candidate species, proposed species and proposed critical habitat be addressed within the consultation. More information on the regulations and procedures for section 7 consultation, including the role of permit or license applicants, can be found in the "Endangered Species Consultation Handbook" at:

http://www.fws.gov/endangered/esa-library/pdf/TOC-GLOS.PDF

Please be aware that bald and golden eagles are protected under the Bald and Golden Eagle Protection Act (16 U.S.C. 668 *et seq.*), and projects affecting these species may require development of an eagle conservation plan

(http://www.fws.gov/windenergy/eagle\_guidance.html). Additionally, wind energy projects should follow the wind energy guidelines (http://www.fws.gov/windenergy/) for minimizing impacts to migratory birds and bats.

Guidance for minimizing impacts to migratory birds for projects including communications towers (e.g., cellular, digital television, radio, and emergency broadcast) can be found at: http://www.fws.gov/migratorybirds/CurrentBirdIssues/Hazards/towers/towers.htm; http://www.towerkill.com; and http://www.fws.gov/migratorybirds/CurrentBirdIssues/Hazards/towers/comtow.html.

We appreciate your concern for threatened and endangered species. The Service encourages Federal agencies to include conservation of threatened and endangered species into their project planning to further the purposes of the Act. Please include the Consultation Tracking Number in the header of this letter with any request for consultation or correspondence about your project that you submit to our office. Submit consultation requests electronically to the following email: msfosection7consultation@fws.gov

#### Attachment(s):

- Official Species List
- USFWS National Wildlife Refuges and Fish Hatcheries
- Migratory Birds
- Wetlands

# **Official Species List**

This list is provided pursuant to Section 7 of the Endangered Species Act, and fulfills the requirement for Federal agencies to "request of the Secretary of the Interior information whether any species which is listed or proposed to be listed may be present in the area of a proposed action".

This species list is provided by:

**Mississippi Ecological Services Field Office** 6578 Dogwood View Parkway, Suite A Jackson, MS 39213-7856 (601) 965-4900

### **Project Summary**

Consultation Code: 04EM1000-2021-SLI-0745 Event Code: 04EM1000-2021-E-01680

Project Name: Golden Triangle II

Project Type: POWER GENERATION
Project Description: Site Evaluation for Solar Farm

Project Location:

Approximate location of the project can be viewed in Google Maps: <a href="https://www.google.com/maps/@33.3930148,-88.59959158688966,14z">https://www.google.com/maps/@33.3930148,-88.59959158688966,14z</a>



Counties: Lowndes County, Mississippi

### **Endangered Species Act Species**

There is a total of 3 threatened, endangered, or candidate species on this species list.

Species on this list should be considered in an effects analysis for your project and could include species that exist in another geographic area. For example, certain fish may appear on the species list because a project could affect downstream species.

IPaC does not display listed species or critical habitats under the sole jurisdiction of NOAA Fisheries<sup>1</sup>, as USFWS does not have the authority to speak on behalf of NOAA and the Department of Commerce.

See the "Critical habitats" section below for those critical habitats that lie wholly or partially within your project area under this office's jurisdiction. Please contact the designated FWS office if you have questions.

NOAA Fisheries, also known as the National Marine Fisheries Service (NMFS), is an
office of the National Oceanic and Atmospheric Administration within the Department of
Commerce.

#### **Mammals**

NAME STATUS

#### Northern Long-eared Bat Myotis septentrionalis

Threatened

No critical habitat has been designated for this species. Species profile: <a href="https://ecos.fws.gov/ecp/species/9045">https://ecos.fws.gov/ecp/species/9045</a>

#### **Birds**

NAME STATUS

#### Wood Stork Mycteria americana

Threatened

Population: AL, FL, GA, MS, NC, SC

No critical habitat has been designated for this species. Species profile: <a href="https://ecos.fws.gov/ecp/species/8477">https://ecos.fws.gov/ecp/species/8477</a>

#### **Flowering Plants**

NAME STATUS

#### Prices Potato-bean *Apios priceana*

Threatened

Population:

No critical habitat has been designated for this species. Species profile: <a href="https://ecos.fws.gov/ecp/species/7422">https://ecos.fws.gov/ecp/species/7422</a>

#### **Critical habitats**

THERE ARE NO CRITICAL HABITATS WITHIN YOUR PROJECT AREA UNDER THIS OFFICE'S JURISDICTION.

# **USFWS National Wildlife Refuge Lands And Fish Hatcheries**

Any activity proposed on lands managed by the <u>National Wildlife Refuge</u> system must undergo a 'Compatibility Determination' conducted by the Refuge. Please contact the individual Refuges to discuss any questions or concerns.

THERE ARE NO REFUGE LANDS OR FISH HATCHERIES WITHIN YOUR PROJECT AREA.

# **Migratory Birds**

Certain birds are protected under the Migratory Bird Treaty Act<sup>1</sup> and the Bald and Golden Eagle Protection Act<sup>2</sup>.

Any person or organization who plans or conducts activities that may result in impacts to migratory birds, eagles, and their habitats should follow appropriate regulations and consider implementing appropriate conservation measures, as described <u>below</u>.

- 1. The Migratory Birds Treaty Act of 1918.
- 2. The <u>Bald and Golden Eagle Protection Act</u> of 1940.
- 3. 50 C.F.R. Sec. 10.12 and 16 U.S.C. Sec. 668(a)

THERE ARE NO FWS MIGRATORY BIRDS OF CONCERN WITHIN THE VICINITY OF YOUR PROJECT AREA.

#### **Migratory Birds FAQ**

Tell me more about conservation measures I can implement to avoid or minimize impacts to migratory birds.

<u>Nationwide Conservation Measures</u> describes measures that can help avoid and minimize impacts to all birds at any location year round. Implementation of these measures is particularly important when birds are most likely to occur in the project area. When birds may be breeding in the area, identifying the locations of any active nests and avoiding their destruction is a very helpful impact minimization measure. To see when birds are most likely to occur and be breeding in your project area, view the Probability of Presence Summary. <u>Additional measures</u> or <u>permits</u> may be advisable depending on the type of activity you are conducting and the type of infrastructure or bird species present on your project site.

# What does IPaC use to generate the migratory birds potentially occurring in my specified location?

The Migratory Bird Resource List is comprised of USFWS <u>Birds of Conservation Concern</u> (<u>BCC</u>) and other species that may warrant special attention in your project location.

The migratory bird list generated for your project is derived from data provided by the <u>Avian Knowledge Network (AKN)</u>. The AKN data is based on a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u> and is queried and filtered to return a list of those birds reported as occurring in the 10km grid cell(s) which your project intersects, and that have been identified as warranting special attention because they are a BCC species in that area, an eagle (<u>Eagle Act</u> requirements may apply), or a species that has a particular vulnerability to offshore activities or development.

Again, the Migratory Bird Resource list includes only a subset of birds that may occur in your project area. It is not representative of all birds that may occur in your project area. To get a list of all birds potentially present in your project area, please visit the <u>AKN Phenology Tool</u>.

# What does IPaC use to generate the probability of presence graphs for the migratory birds potentially occurring in my specified location?

The probability of presence graphs associated with your migratory bird list are based on data provided by the <u>Avian Knowledge Network (AKN)</u>. This data is derived from a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u>.

Probability of presence data is continuously being updated as new and better information becomes available. To learn more about how the probability of presence graphs are produced and how to interpret them, go the Probability of Presence Summary and then click on the "Tell me about these graphs" link.

# How do I know if a bird is breeding, wintering, migrating or present year-round in my project area?

To see what part of a particular bird's range your project area falls within (i.e. breeding, wintering, migrating or year-round), you may refer to the following resources: The Cornell Lab of Ornithology All About Birds Bird Guide, or (if you are unsuccessful in locating the bird of interest there), the Cornell Lab of Ornithology Neotropical Birds guide. If a bird on your migratory bird species list has a breeding season associated with it, if that bird does occur in your project area, there may be nests present at some point within the timeframe specified. If "Breeds elsewhere" is indicated, then the bird likely does not breed in your project area.

#### What are the levels of concern for migratory birds?

Migratory birds delivered through IPaC fall into the following distinct categories of concern:

- 1. "BCC Rangewide" birds are <u>Birds of Conservation Concern</u> (BCC) that are of concern throughout their range anywhere within the USA (including Hawaii, the Pacific Islands, Puerto Rico, and the Virgin Islands);
- 2. "BCC BCR" birds are BCCs that are of concern only in particular Bird Conservation Regions (BCRs) in the continental USA; and
- 3. "Non-BCC Vulnerable" birds are not BCC species in your project area, but appear on your list either because of the <u>Eagle Act</u> requirements (for eagles) or (for non-eagles) potential susceptibilities in offshore areas from certain types of development or activities (e.g. offshore energy development or longline fishing).

Although it is important to try to avoid and minimize impacts to all birds, efforts should be made, in particular, to avoid and minimize impacts to the birds on this list, especially eagles and BCC species of rangewide concern. For more information on conservation measures you can implement to help avoid and minimize migratory bird impacts and requirements for eagles, please see the FAQs for these topics.

#### Details about birds that are potentially affected by offshore projects

For additional details about the relative occurrence and abundance of both individual bird species and groups of bird species within your project area off the Atlantic Coast, please visit the <a href="Northeast Ocean Data Portal">Northeast Ocean Data Portal</a>. The Portal also offers data and information about other taxa besides birds that may be helpful to you in your project review. Alternately, you may download the bird model results files underlying the portal maps through the <a href="NOAA NCCOS Integrative Statistical">NOAA NCCOS Integrative Statistical</a>

Modeling and Predictive Mapping of Marine Bird Distributions and Abundance on the Atlantic Outer Continental Shelf project webpage.

Bird tracking data can also provide additional details about occurrence and habitat use throughout the year, including migration. Models relying on survey data may not include this information. For additional information on marine bird tracking data, see the <u>Diving Bird Study</u> and the <u>nanotag studies</u> or contact <u>Caleb Spiegel</u> or <u>Pam Loring</u>.

#### What if I have eagles on my list?

If your project has the potential to disturb or kill eagles, you may need to <u>obtain a permit</u> to avoid violating the Eagle Act should such impacts occur.

#### **Proper Interpretation and Use of Your Migratory Bird Report**

The migratory bird list generated is not a list of all birds in your project area, only a subset of birds of priority concern. To learn more about how your list is generated, and see options for identifying what other birds may be in your project area, please see the FAO "What does IPaC use to generate the migratory birds potentially occurring in my specified location". Please be aware this report provides the "probability of presence" of birds within the 10 km grid cell(s) that overlap your project; not your exact project footprint. On the graphs provided, please also look carefully at the survey effort (indicated by the black vertical bar) and for the existence of the "no data" indicator (a red horizontal bar). A high survey effort is the key component. If the survey effort is high, then the probability of presence score can be viewed as more dependable. In contrast, a low survey effort bar or no data bar means a lack of data and, therefore, a lack of certainty about presence of the species. This list is not perfect; it is simply a starting point for identifying what birds of concern have the potential to be in your project area, when they might be there, and if they might be breeding (which means nests might be present). The list helps you know what to look for to confirm presence, and helps guide you in knowing when to implement conservation measures to avoid or minimize potential impacts from your project activities, should presence be confirmed. To learn more about conservation measures, visit the FAQ "Tell me about conservation measures I can implement to avoid or minimize impacts to migratory birds" at the bottom of your migratory bird trust resources page.

### Wetlands

Impacts to <u>NWI wetlands</u> and other aquatic habitats may be subject to regulation under Section 404 of the Clean Water Act, or other State/Federal statutes.

For more information please contact the Regulatory Program of the local <u>U.S. Army Corps of Engineers District</u>.

Please note that the NWI data being shown may be out of date. We are currently working to update our NWI data set. We recommend you verify these results with a site visit to determine the actual extent of wetlands on site.

WETLAND INFORMATION WAS NOT AVAILABLE WHEN THIS SPECIES LIST WAS GENERATED. PLEASE VISIT <u>HTTPS://www.fws.gov/wetlands/data/mapper.html</u> or contact the field office for further information.



# Mississippi Department of Wildlife, Fisheries, and Parks

Sam Polles, Ph.D. Executive Director

March 18, 2021

Burns and McDonnell 3650 Mansell Road Suite 300 Atlanta , GA 30022

Re: Golden Triangle

Solar Project - Phase

Ш

Lowndes County, MS

Project # Internal Id 1708

#### To Jesse Brown:

In response to your request for information dated **February 25, 2021**, we have searched our database for occurrences of state or federally listed species and species of special concern that occur within 2 miles of the site of the proposed project. Please find our concerns and recommendations below.

#### The following species of concern may occur within 2 miles of the proposed project area:

Scientific Name	Common Name	Federal Status	State Status	State Rank
Evax prolifera	Big-head Evax			S1
Spilogale putorius	Eastern Spotted Skunk			S1
Carex gracilescens	Slender Sedge			S1
Thelesperma filifolium	Stiff Greenthread			S1
Tyrannus forficatus	Scissor-tailed Flycatcher			S1B,S1N
Carex jamesii	Nebraska Sedge			S1S2
Carya laciniosa	Big Shellbark Hickory			S2
Quercus macrocarpa	Bur Oak			S2

Scientific Name	Common Name	Federal Status	State Status	State Rank
Aesculus glabra	Ohio Buckeye			S2
Polytaenia nuttallii	Prairie Parsley			S2
Nemastylis geminiflora	Prairie Pleatleaf			S2
Mycteria americana	Wood Stork	LT	LE	S2N
Carex microdonta	Small-toothed Sedge			S3
Passerina ciris	Painted Bunting			S3B
Carex meadii	Mead's Sedge			S3S4
Carya myristiciformis	Nutmeg Hickory			S3S4

#### State Rank

- **S1** Critically imperiled in Mississippi because of extreme rarity (5 or fewer occurrences or vey few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.
- S2 Imperiled in Mississippi because of rarity (6 to 20 occurrences or few remaining individuals or acres) or because of some factor(s) making it vulnerable to extirpation.
- S3 Rare or uncommon in Mississippi (on the order of 21 to 100 occurrences).

#### **State and Federal Status**

- LE Endangered A species which is in danger of extinction throughout all or a significant portion of its range.
- LT Threatened A species likely to become endangered in foreseeable future throughout all or a significant portion of its range.

Based on the information provided, we conclude that if best management practices are properly implemented, monitored, and maintained (particularly measures to prevent, or at least, minimize negative impacts to water quality), the proposed project likely poses no threat to listed species or their habitats.

#### Recommendations:

As listed above, there are 16 species of concern in our database within a 2-mile radius of the proposed Golden Triangle Solar Project in Lowndes County, MS (33.385400, -88.620229). Of the 16 species listed, the Wood Stork is listed as federally threaten and state endangered. Many of the listed species are on the decline because of degradation or destruction of essential habitat needed to support them. Concerns for this project is the potential for sediment and pollutants to enter nearby into nearby bodies of water, such as North Branch Magowah Creek. Activities that modify the landscape can be detrimental because they can adversely affect water quality by increasing herbicide and pesticide load, silt load, exhaust runoff from roads, and other unintentional pollutants. These factors may negatively impact habitat conditions by detrimentally affecting respiration, feeding, and reproduction of amphibians, bats, birds, crayfishes, fishes, insects, turtles, and vegetation. Wood Storks inhabit freshwater wetlands and require receding water levels and concentrations of fish. Wood Stork populations have declined throughout its range because of habitat alteration from the construction of canals, levees, and other such structures that have interfered with normal water cycles. Portions within or adjacent to the boundaries of this project site are underlain by hydric soils and may be designated wetlands. If this project is approved, we ask that serious consideration be given to the cumulative impacts of wetland disturbance and elimination, and that appropriate, in-kind mitigation be provided. We recommend that best management practices such as streamside management zones (SMZs) be properly implemented, monitored, and maintained for compliance, specifically measures that will prevent suspended silt and contaminants from leaving the site in storm water run-off as this may negatively affect water quality and habitat conditions within nearby streams and waterbodies (https://www.mfc.ms.gov/forest-health/water-quality-and-forestry-best-management-practices/). If possible, we ask to


leave the trees on the boundaries of streams and the outlined property, as they can act as corridors for wildlife and erosion control.


Please feel free to contact us if we can provide any additional infromation, resources, or assistance that will help minimize negative impacts to the species and/or ecological communities identified in this review. We are happy to work with you to ensure that our state's precious natural heritage is conserved and preserved for future Mississippians.


Completed by Quentin Fairchild

The Mississippi Natural Heritage Program (MNHP) has compiled a database that is the most complete source of information about Mississippi's rare, threatened, and endangered plants, animals, and ecological communities. The quantity and quality of data collected by MNHP are dependent on the research and observations of many individuals and organizations. In many cases, this information is not the result of comprehensive or site-specific field surveys; most natural areas in Mississippi have not been thoroughly surveyed and new occurrences of plant and animal species are often discovered. Heritage reports summarize the existing information known to the MNHP at the time of the request and cannot always be considered a definitive statement on the presence, absence or condition of biological elements on a particular site.













Source: Esri, NRCS SSURGO Soils, Origis, Burns & McDonnell

Issued: 4/20/20


















Source: Esri, NRCS SSURGO Soils, NAIP 2020, Origis, Burns & McDonnell

Issued: 4/28/202









**Photograph 1**: Example of suitable bat roosting habitat within the Golden Triangle II Survey Area. Pictured: Shagbark hickories with exfoliating bark and open understory.






Photograph 2: Example of suitable bat roosting habitat within Golden Triangle II Survey Area. Pictured: Large Snag.



**Photograph 3**: Example of suitable roosting habitat within Golden Triangle Survey Area. Pictured: Scattered snags throughout forested areas.






**Photograph 4**: Example of suitable bat roosting habitat within the Golden Triangle II Survey Area. Pictured: Shagbark hickories with exfoliating bark and open understory.





**Photograph 5:** Example of suitable bat roosting habitat within Golden Triangle II Survey Area. Pictured: Hollow snag.





Photograph 6: Bur Oak (Quercus macrocarpa) observation within Golden Triangle II Survey Area.





Photograph 7: Example of bur oak acorn within Golden Triangle II Survey Area..



Photograph 8: Example of chalk outcrop within Golden Triangle II Survey Area.





**Photograph 9**: Representative photograph of pasture community type within Survey Area.



**Photograph 10**: Representative photograph of PFO wetland habitat within the Survey Area.





Photograph 11: Representative photograph of forested habitat community type within Survey Area.



Photograph 12: Representative photograph of active agriculture field habitat community type within Survey Area – between crop rotations.





**Photograph 9**: General representation of perennial streams within the Survey Area.



Photograph 10: General representation of intermittent streams within the Survey Area





Photograph 11: General representation of ephemeral stream features within Survey Area.



**Photograph 12**: General representation of PUB features within the Survey Area.





Photograph 13: General representation of PEM wetland features within the Survey Area



**Photograph 14**: General representation of forested area in southwest portion of Survey Area. Primarily composed of eastern red cedar; determined not suitable NLEB habitat.







October 5, 2021

Logan Barber Terrestrial Zoologist Tennessee Valley Authority

Re: Acoustic Bat Survey Report for the Origis Golden Triangle 2 Solar Project in Lowndes County, Mississippi Burns & McDonnell Project Number 131497

Dear Mr. Barber:

Burns & McDonnell has been retained by Origis Energy (Origis) to conduct environmental permitting for the Golden Triangle 2 Solar Project (Project). The initial review of protected species information for the Project indicated that potential habitat for the federally threatened northern long-eared bat (*Myotis septentrionalis*) may be impacted by the tree clearing activities (Figure A-1 in Appendix A). Based on recent discussions with the Tennessee Valley Authority (TVA) it was determined that presence/absence surveys should be conducted to assess potential occurrence of the northern long-eared bat. Accordingly, Burns & McDonnell conducted presence/absence surveys for the northern long-eared bat using acoustic methods.

## **METHODS**

The following methods were included in a survey plan submitted to the TVA on July 29, 2021. The survey protocol followed the most current (2020) U.S. Fish and Wildlife (USFWS) Rangewide Indiana Bat Survey Guidelines (Guidelines). Based on the Protected Species Habitat Assessment Report finalized on July 6, 2021, approximately 358 acres of potential northern longeared bat roosting habitat occurs within the Project area. The level of effort specified in the Guidelines for nonlinear projects is 8 detector-nights per 123 acres (or less) of habitat. Therefore, the required level of survey effort on this Project is 24 detector-nights.

The locations of the deployed detectors were determined in the field and based on site conditions by qualified bat biologists (Appendix D). Areas of suitable roosting habitat and proposed acoustic sites are shown in Figure A-1. A total of 11 sites were surveyed. Habitat types sampled included corridors, field/forest edges, ponds, and streams. Forested habitat in the vicinity of the sites were generally dominated by white oak (*Quercus alba*), southern red oak (*Quercus falcata*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), mockernut hickory (*Carya tomentosa*), water hickory (*Carya aquatica*), willow oak (*Quercus phellos*), cherrybark oak (*Quercus pagoda*), swamp chestnut oak (*Quercus michauxii*), green ash (*Fraxinus pennsylvanica*), box elder (*Acer negundo*), and red maple (*Acer rubrum*). Fields near the sample sites were in an early successional state due to crop growth/harvesting, and regular management/mowing as well as repeated cattle grazing. Corn is planted in late spring and covers the row crop fields on approximately 125 acres.



Logan Barber Tennessee Valley Authority October 5, 2021 Page 2

Wildlife Acoustics SM2BAT, SM3BAT, and SM4FS acoustic detectors (detectors) with omnidirectional microphones were used to survey the Project area. One detector was placed at each site and a total of 11 sites were sampled, (Table 1; Appendix B - Photographs). Microphones were positioned at least 3 meters from obstructions and with minimal clutter within 10 meters. The detectors were set to record from 30 minutes before sunset to 30 minutes after sunrise. Recordings were collected in full spectrum, with settings optimized for detection of northern long-eared bat (Table 1). Cable connections and settings were carefully assessed in the field to determine the detector would be operating properly.

Table 1: Key settings for the acoustic detectors used for acoustic bat surveys.

Setting Type	Setting*
Detector model	Wildlife Acoustics SM2BAT, SM3BAT, or SM4FS
Nightly recording time	0.5 hours before sunset to 0.5 hours after sunrise
Recording format	Full-spectrum .wav
Sample frequency	384 kHz
Minimum trigger frequency	16 kHz
Trigger level	12 dB
Trigger window	3 seconds

<sup>\*</sup>All other settings used default values for recording bats.

Bat calls were classified to species using Kaleidoscope Pro with the Bats of North America classifier version 5.4.0 with a 0 (neutral) setting. If Kaleidoscope indicated likely presence (p < 0.05) of northern long-eared bat, all bat calls from that sight and night would be reviewed manually by Josiah Maine. Call characteristics would be compared to known representative calls and known call parameters to determine species.

## RESULTS

Detectors were placed at sites 1, 2, 3, 4, and 5 on August 3, 2021, and were retrieved on August 5, 2021 (Table 2). Detectors were placed at sites 6, 7, 8, and 9 on August 4, 2021, and retrieved on August 6, 2021. At sites 10 and 11, detectors were placed on August 5, 2021, and retrieved on August 10, 2021. Weather during the survey period included temperatures ranging from 66 degrees Fahrenheit to 86 degrees Fahrenheit, wind speeds of 3 to 10 miles per hour, and no precipitation. Winds did not exceed 9 miles per hour for more than 30 minutes (Table C-12 in Appendix C); therefore, weather conditions were within the bounds of the USFWS Guidelines. Detectors generally functioned properly, except at site 11, where the acoustic data exceeded the



Logan Barber Tennessee Valley Authority October 5, 2021 Page 3

card capacity the evening of August 8. At 26 complete detector-nights, the level of survey effort was sufficient to meet the requirements of the Guidelines.

Table 2: Acoustic survey site coordinates and habitat.

Site	<b>Detector-nights</b>	Lat	Long	Habitat
1	2	33.4016513	-88.6010734	Corridor/forest edge
2	2	33.4019296	-88.5983261	Field/forest edge by pond
3	2	33.3979317	-88.5953295	Forest opening
4	2	33.3920261	-88.5993483	Field/forest edge
5	2	33.3890347	-88.6055889	Field/forest edge
6	2	33.3934448	-88.6147555	Corridor/forest opening
7	2	33.3891569	-88.6144079	Field/forest edge
8	2	33.3871545	-88.6250538	Field/forest edge
9	2	33.3825378	-88.6225526	Field/forest edge
10	5	33.3843164	-88.6017695	Field/forest edge by stream
11	3	33.3835537	-88.607802	Field/forest edge

Across the 11 sites, 9 species were determined to be present by the Kaleidoscope Pro classifier: big brown bat (*Eptesicus fuscus*), eastern red bat (*Lasiurus borealis*), hoary bat (*Lasiurus cinereus*), Seminole bat (*Lasiurus seminolus*), little brown bat (*Myotis lucifugus*), southeastern bat (*Myotis austroriparius*), tricolored bat (*Perimyotis subflavus*), evening bat (*Nycticeius humeralis*), and Mexican free-tailed bat (*Tadarida brasiliensis*). No threatened or endangered species were determined present at any of the sites; therefore, qualitative analysis (manual review) was not necessary. Data tables with detailed analysis results for each site are included in Appendix C.

## **CONCLUSION**

Acoustic surveys were conducted at 11 sites (a total of 26 detector-nights) for the Project, in accordance with USFWS Guidelines. Northern long-eared bat was determined to be likely absent based on automated identification results. Therefore, impacts to northern long-eared bat from implementation of this Project are not anticipated.



Logan Barber Tennessee Valley Authority October 5, 2021 Page 4

If you have any questions or comments regarding this bat study plan, please contact me by phone at 816-448-7519 or by email at jjmaine@burnsmcd.com.

Sincerely,

Josiah Maine

**Environmental Scientist** 


South May


Attachments:

Appendix A - Figures

Appendix B - Acoustic Survey Photographs

Appendix C - Data Tables









Photograph B-1: View of acoustic site 1, facing north.



Photograph B-2: View of acoustic site 2, facing east.





Photograph B-3: View of acoustic site 3, facing west.



Photograph B-4: View of acoustic site 4, facing east.





Photograph B-5: View of acoustic site 5, facing south.



Photograph B-6: View of acoustic site 6, facing north.





Photograph B-7: View of acoustic site 7, facing east.



Photograph B-8: View of acoustic site 8, facing southwest.





Photograph B-9: View of acoustic site 9, facing west.



Photograph B-10: View of acoustic site 10, facing south.





Photograph B-11: View of acoustic site 11, facing south.



Table C-1: Automated identification results for site 1. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	211	0
Eastern red bat	Lasiurus borealis	372	0
Hoary bat	Lasiurus cinereus	1	1
Seminole bat	Lasiurus seminolus	24	1
Southeastern bat	Myotis austroriparius	12	0
Little brown bat	Myotis lucifugus	135	0
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	97	1
Tricolored bat	Perimyotis subflavus	318	0
Mexican free-tailed bat	Tadarida brasiliensis	32	< 0.01
Unidentified bat		927	
Noise		3181	
Total bat files		2129	

Table C-2: Automated identification results for site 2. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	257	0
Eastern red bat	Lasiurus borealis	665	0
Hoary bat	Lasiurus cinereus	5	1
Seminole bat	Lasiurus seminolus	35	1
Southeastern bat	Myotis austroriparius	5	< 0.01
Little brown bat	Myotis lucifugus	79	0.41
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	59	1
Tricolored bat	Perimyotis subflavus	120	< 0.01
Mexican free-tailed bat	Tadarida brasiliensis	133	0
Unidentified bat		791	
Noise		2943	
Total bat files		2149	

Table C-3: Automated identification results for site 3. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	73	0
Eastern red bat	Lasiurus borealis	135	0
Hoary bat	Lasiurus cinereus		1
Seminole bat	Lasiurus seminolus	24	1
Southeastern bat	Myotis austroriparius	1	0.01
Little brown bat	Myotis lucifugus	10	1
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	29	1
Tricolored bat	Perimyotis subflavus	23	0.31
Mexican free-tailed bat	Tadarida brasiliensis	10	0.07
Unidentified bat		277	
Noise		51	
Total bat files		582	

Table C-4: Automated identification results for site 4. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	117	0
Eastern red bat	Lasiurus borealis	322	0
Hoary bat	Lasiurus cinereus	59	0
Seminole bat	Lasiurus seminolus	26	1
Southeastern bat	Myotis austroriparius	2	0.55
Little brown bat	Myotis lucifugus	66	< 0.01
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	76	1
Tricolored bat	Perimyotis subflavus	87	0
Mexican free-tailed bat	Tadarida brasiliensis	41	0
Unidentified bat		595	
Noise		2495	
Total bat files		1391	

Table C-5: Automated identification results for site 5. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	73	0
Eastern red bat	Lasiurus borealis	133	0
Hoary bat	Lasiurus cinereus	1	1
Seminole bat	Lasiurus seminolus	67	0.01
Southeastern bat	Myotis austroriparius		1
Little brown bat	Myotis lucifugus	10	1
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	15	1
Tricolored bat	Perimyotis subflavus	77	0
Mexican free-tailed bat	Tadarida brasiliensis	79	0
Unidentified bat		296	
Noise		1238	
Total bat files		751	

Table C-6: Automated identification results for site 6. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	282	0
Eastern red bat	Lasiurus borealis	212	0
Hoary bat	Lasiurus cinereus	1	1
Seminole bat	Lasiurus seminolus	3	1
Southeastern bat	Myotis austroriparius	4	< 0.01
Little brown bat	Myotis lucifugus	49	< 0.01
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	40	1
Tricolored bat	Perimyotis subflavus	85	0
Mexican free-tailed bat	Tadarida brasiliensis	3	1
Unidentified bat		260	
Noise		275	
Total bat files		939	

Table C-7: Automated identification results for site 7. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	3	0.38
Eastern red bat	Lasiurus borealis	2	0.13
Hoary bat	Lasiurus cinereus		1
Seminole bat	Lasiurus seminolus	2	0.60
Southeastern bat	Myotis austroriparius	14	0
Little brown bat	Myotis lucifugus	3	0.18
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis		1
Tricolored bat	Perimyotis subflavus	13	0
Mexican free-tailed bat	Tadarida brasiliensis	12	0
Unidentified bat		68	
Noise		1097	
Total bat files		117	

Table C-8: Automated identification results for site 8. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	121	0
Eastern red bat	Lasiurus borealis	38	0
Hoary bat	Lasiurus cinereus	4	1
Seminole bat	Lasiurus seminolus	10	0.86
Southeastern bat	Myotis austroriparius	2	< 0.01
Little brown bat	Myotis lucifugus	12	0.08
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	16	1
Tricolored bat	Perimyotis subflavus	11	0.15
Mexican free-tailed bat	Tadarida brasiliensis	3	1
Unidentified bat		147	
Noise		7302	
Total bat files		364	

Table C-9: Automated identification results for site 9. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	48	0
Eastern red bat	Lasiurus borealis	16	0
Hoary bat	Lasiurus cinereus		1
Seminole bat	Lasiurus seminolus	1	1
Southeastern bat	Myotis austroriparius	3	< 0.01
Little brown bat	Myotis lucifugus	28	0
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis		1
Tricolored bat	Perimyotis subflavus	4	0.34
Mexican free-tailed bat	Tadarida brasiliensis	1	1
Unidentified bat		75	
Noise		926	
Total bat files		176	

Table C-10: Automated identification results for site 10. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	178	0
Eastern red bat	Lasiurus borealis	173	0
Hoary bat	Lasiurus cinereus	14	0.97
Seminole bat	Lasiurus seminolus	78	< 0.01
Southeastern bat	Myotis austroriparius	14	0
Little brown bat	Myotis lucifugus	59	< 0.01
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	82	1
Tricolored bat	Perimyotis subflavus	163	0
Mexican free-tailed bat	Tadarida brasiliensis	83	0
Unidentified bat		573	
Noise		8386	
Total bat files		1417	

Table C-11: Automated identification results for site 11. MLE scores less than 0.05 indicate presence as determined by Kaleidoscope.

Common Name	Scientific Name	Files	MLE
Big brown bat	Eptesicus fuscus	52	0
Eastern red bat	Lasiurus borealis	25	0
Hoary bat	Lasiurus cinereus		1
Seminole bat	Lasiurus seminolus	14	0.58
Southeastern bat	Myotis austroriparius	7	0
Little brown bat	Myotis lucifugus	36	0
Northern long-eared bat	Myotis septenrionalis		1
Evening bat	Nycticeius humeralis	47	0.02
Tricolored bat	Perimyotis subflavus	43	0
Mexican free-tailed bat	Tadarida brasiliensis	22	0
Unidentified bat		170	
Noise		5482	
Total bat files		416	

Table C-12: Weather conditions from the Golden Triangle Regional Airport during the acoustic surveys.

	Temperature	Wind Speed	
Date & Time	(° <b>F</b> )	(mph)	Weather Summary
08/03/2021 19:25 CDT	78.8	5.75	clear
08/03/2021 19:30 CDT	77	5.75	clear
08/03/2021 19:35 CDT	77	5.75	clear
08/03/2021 19:40 CDT	77	5.75	clear
08/03/2021 19:45 CDT	75.2	4.6	clear
08/03/2021 19:50 CDT	75.2	5.75	clear
08/03/2021 19:55 CDT	75.2	5.75	clear
08/03/2021 19:56 CDT	75.92	5.75	clear
08/03/2021 20:00 CDT	75.2	5.75	clear
08/03/2021 20:05 CDT	75.2	4.6	clear
08/03/2021 20:10 CDT	75.2	4.6	clear
08/03/2021 20:15 CDT	75.2	4.6	clear
08/03/2021 20:20 CDT	75.2	4.6	clear
08/03/2021 20:25 CDT	75.2	4.6	clear
08/03/2021 20:30 CDT	75.2	3.45	clear
08/03/2021 20:35 CDT	75.2	3.45	clear
08/03/2021 20:40 CDT	73.4	3.45	clear
08/03/2021 20:45 CDT	73.4	3.45	clear
08/03/2021 20:50 CDT	73.4	3.45	clear
08/03/2021 20:55 CDT	73.4	4.6	clear
08/03/2021 20:56 CDT	73.94	4.6	clear
08/03/2021 21:00 CDT	73.4	3.45	clear
08/03/2021 21:05 CDT	73.4	4.6	clear
08/03/2021 21:10 CDT	73.4	4.6	clear
08/03/2021 21:15 CDT	73.4	3.45	clear
08/03/2021 21:20 CDT	73.4	3.45	clear
08/03/2021 21:25 CDT	71.6	3.45	clear
08/03/2021 21:30 CDT	73.4	4.6	clear
08/03/2021 21:35 CDT	71.6	4.6	clear
08/03/2021 21:40 CDT	71.6	5.75	clear
08/03/2021 21:45 CDT	71.6	5.75	clear
08/03/2021 21:50 CDT	71.6	5.75	clear
08/03/2021 21:55 CDT	69.8	5.75	clear
08/03/2021 21:56 CDT	69.98	5.75	clear
08/03/2021 22:00 CDT	71.6	5.75	clear
08/03/2021 22:05 CDT	71.6	5.75	clear
08/03/2021 22:10 CDT	71.6	5.75	clear
08/03/2021 22:15 CDT	69.8	5.75	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/03/2021 22:20 CDT	69.8	4.6	clear
08/03/2021 22:25 CDT	71.6	4.6	clear
08/03/2021 22:30 CDT	71.6	3.45	clear
08/03/2021 22:35 CDT	69.8	4.6	clear
08/03/2021 22:40 CDT	71.6	5.75	clear
08/03/2021 22:45 CDT	69.8	4.6	clear
08/03/2021 22:50 CDT	69.8	4.6	clear
08/03/2021 22:55 CDT	69.8	5.75	clear
08/03/2021 22:56 CDT	69.98	4.6	clear
08/03/2021 23:00 CDT	69.8	4.6	clear
08/03/2021 23:05 CDT	69.8	4.6	clear
08/03/2021 23:10 CDT	69.8	4.6	clear
08/03/2021 23:15 CDT	69.8	0	clear
08/03/2021 23:20 CDT	69.8	0	clear
08/03/2021 23:25 CDT	69.8	0	clear
08/03/2021 23:30 CDT	69.8	0	clear
08/03/2021 23:35 CDT	69.8	0	clear
08/03/2021 23:40 CDT	69.8	0	clear
08/03/2021 23:45 CDT	68	0	clear
08/03/2021 23:50 CDT	69.8	0	clear
08/03/2021 23:55 CDT	69.8	4.6	clear
08/03/2021 23:56 CDT	69.08	4.6	clear
08/04/2021 00:00 CDT	69.8	4.6	clear
08/04/2021 00:05 CDT	68	3.45	clear
08/04/2021 00:10 CDT	68	3.45	clear
08/04/2021 00:15 CDT	68	0	clear
08/04/2021 00:20 CDT	68	0	clear
08/04/2021 00:25 CDT	66.2	0	clear
08/04/2021 00:30 CDT	66.2	3.45	clear
08/04/2021 00:35 CDT	66.2	3.45	clear
08/04/2021 00:40 CDT	66.2	5.75	clear
08/04/2021 00:45 CDT	68	5.75	clear
08/04/2021 00:50 CDT	68	5.75	clear
08/04/2021 00:55 CDT	66.2	4.6	clear
08/04/2021 00:56 CDT	66.92	4.6	clear
08/04/2021 01:00 CDT	66.2	3.45	clear
08/04/2021 01:05 CDT	66.2	3.45	clear
08/04/2021 01:10 CDT	66.2	0	clear
08/04/2021 01:15 CDT	66.2	3.45	clear
08/04/2021 01:20 CDT	66.2	4.6	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/04/2021 01:25 CDT	66.2	3.45	clear
08/04/2021 01:30 CDT	66.2	3.45	clear
08/04/2021 01:35 CDT	66.2	4.6	clear
08/04/2021 01:40 CDT	66.2	4.6	clear
08/04/2021 01:45 CDT	66.2	3.45	clear
08/04/2021 01:50 CDT	66.2	3.45	clear
08/04/2021 01:55 CDT	66.2	3.45	clear
08/04/2021 01:56 CDT	66.02	3.45	clear
08/04/2021 02:00 CDT	66.2	3.45	clear
08/04/2021 02:05 CDT	66.2	0	clear
08/04/2021 02:10 CDT	66.2	0	clear
08/04/2021 02:15 CDT	66.2	0	clear
08/04/2021 02:20 CDT	66.2	0	clear
08/04/2021 02:25 CDT	66.2	3.45	clear
08/04/2021 02:30 CDT	64.4	4.6	clear
08/04/2021 02:35 CDT	64.4	3.45	clear
08/04/2021 02:40 CDT	64.4	3.45	clear
08/04/2021 02:45 CDT	64.4	4.6	clear
08/04/2021 02:50 CDT	64.4	4.6	clear
08/04/2021 02:55 CDT	64.4	4.6	clear
08/04/2021 02:56 CDT	64.94	4.6	clear
08/04/2021 03:00 CDT	64.4	4.6	clear
08/04/2021 03:05 CDT	64.4	4.6	clear
08/04/2021 03:10 CDT	64.4	4.6	clear
08/04/2021 03:15 CDT	64.4	4.6	clear
08/04/2021 03:20 CDT	64.4	4.6	clear
08/04/2021 03:25 CDT	64.4	4.6	clear
08/04/2021 03:30 CDT	64.4	3.45	clear
08/04/2021 03:35 CDT	64.4	3.45	clear
08/04/2021 03:40 CDT	64.4	3.45	clear
08/04/2021 03:45 CDT	64.4	3.45	clear
08/04/2021 03:50 CDT	64.4	3.45	clear
08/04/2021 03:55 CDT	64.4	3.45	clear
08/04/2021 03:56 CDT	64.94	3.45	clear
08/04/2021 04:00 CDT	64.4	0	clear
08/04/2021 04:05 CDT	64.4	0	clear
08/04/2021 04:10 CDT	64.4	3.45	clear
08/04/2021 04:15 CDT	66.2	0	clear
08/04/2021 04:20 CDT	66.2	3.45	clear
08/04/2021 04:25 CDT	64.4	4.6	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/04/2021 04:30 CDT	64.4	3.45	clear
08/04/2021 04:35 CDT	64.4	0	clear
08/04/2021 04:40 CDT	64.4	0	clear
08/04/2021 04:45 CDT	64.4	0	clear
08/04/2021 04:50 CDT	64.4	0	clear
08/04/2021 04:55 CDT	64.4	0	clear
08/04/2021 04:56 CDT	64.04	0	clear
08/04/2021 05:00 CDT	64.4	0	clear
08/04/2021 05:05 CDT	64.4	0	clear
08/04/2021 05:10 CDT	64.4	3.45	clear
08/04/2021 05:15 CDT	64.4	4.6	clear
08/04/2021 05:20 CDT	64.4	4.6	clear
08/04/2021 05:25 CDT	64.4	4.6	clear
08/04/2021 05:30 CDT	64.4	4.6	clear
08/04/2021 05:35 CDT	64.4	4.6	clear
08/04/2021 05:40 CDT	62.6	3.45	clear
08/04/2021 05:45 CDT	62.6	3.45	clear
08/04/2021 05:50 CDT	62.6	3.45	clear
08/04/2021 05:55 CDT	62.6	3.45	clear
08/04/2021 05:56 CDT	62.96	3.45	clear
08/04/2021 06:00 CDT	62.6	3.45	clear
08/04/2021 06:05 CDT	62.6	0	clear
08/04/2021 06:10 CDT	62.6	3.45	clear
08/04/2021 06:15 CDT	62.6	3.45	clear
08/04/2021 06:20 CDT	62.6	3.45	clear
08/04/2021 06:25 CDT	62.6	3.45	clear
08/04/2021 06:30 CDT	62.6	3.45	clear
08/04/2021 06:35 CDT	62.6	3.45	clear
08/04/2021 06:40 CDT	62.6	3.45	clear
08/04/2021 19:25 CDT	78.8	3.45	thin scattered
08/04/2021 19:30 CDT	78.8	4.6	thin scattered
08/04/2021 19:35 CDT	78.8	4.6	clear
08/04/2021 19:40 CDT	78.8	3.45	clear
08/04/2021 19:45 CDT	77	3.45	clear
08/04/2021 19:50 CDT	78.8	3.45	thin scattered
08/04/2021 19:55 CDT	77	5.75	scattered
08/04/2021 19:56 CDT	77	4.6	broken
08/04/2021 20:00 CDT	77	4.6	broken
08/04/2021 20:05 CDT	77	4.6	broken
08/04/2021 20:10 CDT	75.2	4.6	scattered

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/04/2021 20:15 CDT	75.2	4.6	broken
08/04/2021 20:20 CDT	75.2	4.6	broken
08/04/2021 20:25 CDT	75.2	3.45	broken
08/04/2021 20:30 CDT	75.2	3.45	broken
08/04/2021 20:35 CDT	75.2	4.6	broken
08/04/2021 20:40 CDT	75.2	4.6	broken
08/04/2021 20:45 CDT	75.2	4.6	scattered
08/04/2021 20:50 CDT	75.2	4.6	scattered
08/04/2021 20:55 CDT	73.4	4.6	thin scattered
08/04/2021 20:56 CDT	73.94	4.6	thin scattered
08/04/2021 21:00 CDT	73.4	4.6	thin scattered
08/04/2021 21:05 CDT	73.4	4.6	clear
08/04/2021 21:10 CDT	73.4	4.6	clear
08/04/2021 21:15 CDT	73.4	5.75	clear
08/04/2021 21:20 CDT	73.4	6.91	clear
08/04/2021 21:25 CDT	71.6	6.91	clear
08/04/2021 21:30 CDT	71.6	6.91	clear
08/04/2021 21:35 CDT	71.6	5.75	clear
08/04/2021 21:40 CDT	71.6	6.91	clear
08/04/2021 21:45 CDT	71.6	5.75	clear
08/04/2021 21:50 CDT	71.6	5.75	clear
08/04/2021 21:55 CDT	71.6	5.75	clear
08/04/2021 21:56 CDT	71.96	5.75	clear
08/04/2021 22:00 CDT	71.6	5.75	clear
08/04/2021 22:05 CDT	71.6	5.75	clear
08/04/2021 22:10 CDT	71.6	4.6	clear
08/04/2021 22:15 CDT	71.6	4.6	clear
08/04/2021 22:20 CDT	71.6	4.6	clear
08/04/2021 22:25 CDT	71.6	5.75	clear
08/04/2021 22:30 CDT	71.6	4.6	clear
08/04/2021 22:35 CDT	71.6	3.45	clear
08/04/2021 22:40 CDT	71.6	3.45	clear
08/04/2021 22:45 CDT	71.6	3.45	clear
08/04/2021 22:50 CDT	71.6	3.45	clear
08/04/2021 22:55 CDT	71.6	3.45	clear
08/04/2021 22:56 CDT	71.06	3.45	clear
08/04/2021 23:00 CDT	69.8	3.45	clear
08/04/2021 23:05 CDT	71.6	3.45	clear
08/04/2021 23:10 CDT	69.8	3.45	clear
08/04/2021 23:15 CDT	69.8	3.45	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/04/2021 23:20 CDT	69.8	3.45	clear
08/04/2021 23:25 CDT	69.8	4.6	clear
08/04/2021 23:30 CDT	69.8	3.45	clear
08/04/2021 23:35 CDT	69.8	3.45	clear
08/04/2021 23:40 CDT	69.8	3.45	clear
08/04/2021 23:45 CDT	69.8	4.6	clear
08/04/2021 23:50 CDT	68	4.6	clear
08/04/2021 23:55 CDT	68	4.6	clear
08/04/2021 23:56 CDT	69.08	3.45	clear
08/05/2021 00:00 CDT	69.8	3.45	clear
08/05/2021 00:05 CDT	69.8	4.6	clear
08/05/2021 00:10 CDT	69.8	4.6	clear
08/05/2021 00:15 CDT	69.8	4.6	clear
08/05/2021 00:20 CDT	69.8	4.6	clear
08/05/2021 00:25 CDT	68	4.6	clear
08/05/2021 00:30 CDT	68	3.45	clear
08/05/2021 00:35 CDT	68	4.6	clear
08/05/2021 00:40 CDT	68	4.6	clear
08/05/2021 00:45 CDT	68	4.6	clear
08/05/2021 00:50 CDT	68	4.6	clear
08/05/2021 00:55 CDT	68	3.45	clear
08/05/2021 00:56 CDT	68	3.45	clear
08/05/2021 01:00 CDT	68	4.6	clear
08/05/2021 01:05 CDT	66.2	4.6	clear
08/05/2021 01:10 CDT	68	4.6	clear
08/05/2021 01:15 CDT	66.2	4.6	clear
08/05/2021 01:20 CDT	66.2	4.6	clear
08/05/2021 01:25 CDT	66.2	4.6	clear
08/05/2021 01:30 CDT	66.2	4.6	clear
08/05/2021 01:35 CDT	66.2	0	clear
08/05/2021 01:40 CDT	66.2	0	clear
08/05/2021 01:45 CDT	66.2	3.45	clear
08/05/2021 01:50 CDT	68	3.45	clear
08/05/2021 01:55 CDT	66.2	3.45	clear
08/05/2021 01:56 CDT	66.92	3.45	clear
08/05/2021 02:00 CDT	66.2	3.45	clear
08/05/2021 02:05 CDT	66.2	3.45	clear
08/05/2021 02:10 CDT	66.2	3.45	clear
08/05/2021 02:15 CDT	66.2	3.45	clear
08/05/2021 02:20 CDT	66.2	3.45	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/05/2021 02:25 CDT	66.2	4.6	clear
08/05/2021 02:30 CDT	66.2	3.45	clear
08/05/2021 02:35 CDT	66.2	3.45	clear
08/05/2021 02:40 CDT	66.2	3.45	clear
08/05/2021 02:45 CDT	66.2	4.6	clear
08/05/2021 02:50 CDT	66.2	4.6	clear
08/05/2021 02:55 CDT	66.2	4.6	clear
08/05/2021 02:56 CDT	66.02	4.6	clear
08/05/2021 03:00 CDT	66.2	3.45	clear
08/05/2021 03:05 CDT	66.2	3.45	clear
08/05/2021 03:10 CDT	66.2	4.6	clear
08/05/2021 03:15 CDT	66.2	3.45	clear
08/05/2021 03:20 CDT	64.4	0	clear
08/05/2021 03:25 CDT	66.2	3.45	clear
08/05/2021 03:30 CDT	66.2	3.45	clear
08/05/2021 03:35 CDT	64.4	3.45	clear
08/05/2021 03:40 CDT	64.4	3.45	clear
08/05/2021 03:45 CDT	66.2	0	clear
08/05/2021 03:50 CDT	66.2	0	clear
08/05/2021 03:55 CDT	64.4	0	clear
08/05/2021 03:56 CDT	64.94	0	clear
08/05/2021 04:00 CDT	64.4	0	clear
08/05/2021 04:05 CDT	66.2	0	clear
08/05/2021 04:10 CDT	66.2	0	clear
08/05/2021 04:15 CDT	66.2	3.45	clear
08/05/2021 04:20 CDT	64.4	0	clear
08/05/2021 04:25 CDT	64.4	0	clear
08/05/2021 04:30 CDT	66.2	0	clear
08/05/2021 04:35 CDT	64.4	0	clear
08/05/2021 04:40 CDT	64.4	0	clear
08/05/2021 04:45 CDT	64.4	3.45	clear
08/05/2021 04:50 CDT	64.4	3.45	clear
08/05/2021 04:55 CDT	64.4	0	clear
08/05/2021 04:56 CDT	64.94	0	clear
08/05/2021 05:00 CDT	64.4	0	clear
08/05/2021 05:05 CDT	64.4	3.45	clear
08/05/2021 05:10 CDT	64.4	0	clear
08/05/2021 05:15 CDT	64.4	0	clear
08/05/2021 05:20 CDT	64.4	0	clear
08/05/2021 05:25 CDT	64.4	3.45	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/05/2021 05:30 CDT	64.4	0	clear
08/05/2021 05:40 CDT	64.4	4.6	clear
08/05/2021 05:45 CDT	64.4	3.45	clear
08/05/2021 05:50 CDT	64.4	4.6	clear
08/05/2021 05:55 CDT	64.4	4.6	clear
08/05/2021 05:56 CDT	64.04	4.6	clear
08/05/2021 06:05 CDT	64.4	0	clear
08/05/2021 06:10 CDT	64.4	0	clear
08/05/2021 06:15 CDT	64.4	0	clear
08/05/2021 06:20 CDT	64.4	3.45	clear
08/05/2021 06:25 CDT	64.4	3.45	clear
08/05/2021 06:30 CDT	64.4	3.45	clear
08/05/2021 06:35 CDT	64.4	3.45	clear
08/05/2021 06:40 CDT	64.4	4.6	clear
08/05/2021 19:25 CDT	82.4	3.45	broken
08/05/2021 19:35 CDT	80.6	3.45	scattered
08/05/2021 19:40 CDT	80.6	3.45	scattered
08/05/2021 19:45 CDT	80.6	3.45	thin scattered
08/05/2021 19:50 CDT	80.6	3.45	scattered
08/05/2021 19:55 CDT	80.6	3.45	broken
08/05/2021 19:56 CDT	80.06	3.45	broken
08/05/2021 20:00 CDT	80.6	3.45	broken
08/05/2021 20:05 CDT	80.6	0	broken
08/05/2021 20:10 CDT	78.8	0	overcast
08/05/2021 20:15 CDT	78.8	3.45	overcast
08/05/2021 20:20 CDT	78.8	3.45	overcast
08/05/2021 20:25 CDT	78.8	3.45	overcast
08/05/2021 20:30 CDT	78.8	3.45	overcast
08/05/2021 20:35 CDT	78.8	0	overcast
08/05/2021 20:40 CDT	80.6	3.45	overcast
08/05/2021 20:45 CDT	80.6	5.75	overcast
08/05/2021 20:50 CDT	80.6	4.6	overcast
08/05/2021 20:55 CDT	80.6	4.6	overcast
08/05/2021 20:56 CDT	80.96	4.6	overcast
08/05/2021 21:00 CDT	80.6	5.75	overcast
08/05/2021 21:05 CDT	80.6	6.91	overcast
08/05/2021 21:10 CDT	80.6	5.75	overcast
08/05/2021 21:15 CDT	78.8	4.6	overcast
08/05/2021 21:20 CDT	78.8	3.45	overcast
08/05/2021 21:25 CDT	78.8	3.45	overcast

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/05/2021 21:30 CDT	78.8	3.45	overcast
08/05/2021 21:35 CDT	77	4.6	overcast
08/05/2021 21:40 CDT	78.8	3.45	overcast
08/05/2021 21:45 CDT	78.8	3.45	overcast
08/05/2021 21:50 CDT	78.8	0	overcast
08/05/2021 21:55 CDT	78.8	0	overcast
08/05/2021 21:56 CDT	78.08	0	overcast
08/05/2021 22:00 CDT	78.8	0	overcast
08/05/2021 22:05 CDT	78.8	0	overcast
08/05/2021 22:10 CDT	78.8	0	overcast
08/05/2021 22:15 CDT	78.8	0	overcast
08/05/2021 22:20 CDT	78.8	0	broken
08/05/2021 22:25 CDT	77	0	broken
08/05/2021 22:30 CDT	77	0	broken
08/05/2021 22:35 CDT	77	0	broken
08/05/2021 22:40 CDT	78.8	0	broken
08/05/2021 22:45 CDT	78.8	0	broken
08/05/2021 22:50 CDT	77	0	broken
08/05/2021 22:55 CDT	77	0	broken
08/05/2021 22:56 CDT	77	0	broken
08/05/2021 23:00 CDT	75.2	0	overcast
08/05/2021 23:05 CDT	75.2	0	overcast
08/05/2021 23:10 CDT	75.2	3.45	overcast
08/05/2021 23:15 CDT	75.2	0	overcast
08/05/2021 23:20 CDT	75.2	0	overcast
08/05/2021 23:25 CDT	77	0	overcast
08/05/2021 23:30 CDT	77	0	overcast
08/05/2021 23:35 CDT	77	0	overcast
08/05/2021 23:40 CDT	77	0	overcast
08/05/2021 23:45 CDT	77	0	overcast
08/05/2021 23:50 CDT	77	3.45	overcast
08/05/2021 23:55 CDT	77	3.45	overcast
08/05/2021 23:56 CDT	77	3.45	overcast
08/06/2021 00:00 CDT	77	3.45	overcast
08/06/2021 00:05 CDT	75.2	0	overcast
08/06/2021 00:10 CDT	77	0	overcast
08/06/2021 00:15 CDT	78.8	0	overcast
08/06/2021 00:20 CDT	78.8	0	overcast
08/06/2021 00:25 CDT	78.8	3.45	overcast
08/06/2021 00:30 CDT	77	5.75	overcast

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/06/2021 00:35 CDT	75.2	8.06	overcast
08/06/2021 00:40 CDT	75.2	6.91	overcast
08/06/2021 00:45 CDT	75.2	4.6	overcast
08/06/2021 00:50 CDT	75.2	4.6	overcast
08/06/2021 00:55 CDT	75.2	0	overcast
08/06/2021 00:56 CDT	75.02	0	overcast
08/06/2021 01:00 CDT	75.2	3.45	overcast
08/06/2021 01:05 CDT	75.2	0	overcast
08/06/2021 01:10 CDT	75.2	0	overcast
08/06/2021 01:15 CDT	75.2	3.45	overcast
08/06/2021 01:20 CDT	75.2	3.45	overcast
08/06/2021 01:25 CDT	75.2	0	overcast
08/06/2021 01:30 CDT	75.2	0	overcast
08/06/2021 01:35 CDT	75.2	0	overcast
08/06/2021 01:40 CDT	75.2	0	overcast
08/06/2021 01:45 CDT	75.2	0	overcast
08/06/2021 01:50 CDT	75.2	0	overcast
08/06/2021 01:55 CDT	75.2	0	overcast
08/06/2021 01:56 CDT	75.92	0	overcast
08/06/2021 02:00 CDT	75.2	0	overcast
08/06/2021 02:05 CDT	75.2	0	overcast
08/06/2021 02:10 CDT	75.2	0	overcast
08/06/2021 02:15 CDT	75.2	0	broken
08/06/2021 02:20 CDT	75.2	0	broken
08/06/2021 02:25 CDT	75.2	3.45	broken
08/06/2021 02:30 CDT	75.2	3.45	broken
08/06/2021 02:35 CDT	75.2	6.91	thin scattered
08/06/2021 02:40 CDT	75.2	5.75	thin scattered
08/06/2021 02:45 CDT	75.2	5.75	thin scattered
08/06/2021 02:50 CDT	75.2	4.6	thin scattered
08/06/2021 02:55 CDT	75.2	4.6	clear
08/06/2021 02:56 CDT	75.92	4.6	clear
08/06/2021 03:00 CDT	75.2	4.6	clear
08/06/2021 03:05 CDT	75.2	0	clear
08/06/2021 03:10 CDT	77	0	thin scattered
08/06/2021 03:15 CDT	77	3.45	thin scattered
08/06/2021 03:20 CDT	77	0	thin scattered
08/06/2021 03:25 CDT	77	3.45	thin scattered
08/06/2021 03:30 CDT	77	0	thin scattered
08/06/2021 03:35 CDT	75.2	0	thin scattered

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/06/2021 03:40 CDT	75.2	0	thin scattered
08/06/2021 03:45 CDT	75.2	0	thin scattered
08/06/2021 03:50 CDT	75.2	0	thin scattered
08/06/2021 03:55 CDT	75.2	3.45	thin scattered
08/06/2021 03:56 CDT	75.02	3.45	thin scattered
08/06/2021 04:00 CDT	75.2	3.45	thin scattered
08/06/2021 04:05 CDT	75.2	3.45	thin scattered
08/06/2021 04:10 CDT	75.2	5.75	thin scattered
08/06/2021 04:15 CDT	75.2	4.6	clear
08/06/2021 04:20 CDT	75.2	3.45	clear
08/06/2021 04:25 CDT	75.2	3.45	clear
08/06/2021 04:30 CDT	75.2	3.45	clear
08/06/2021 04:35 CDT	75.2	3.45	clear
08/06/2021 04:40 CDT	75.2	3.45	clear
08/06/2021 04:45 CDT	75.2	0	clear
08/06/2021 04:50 CDT	75.2	3.45	clear
08/06/2021 04:55 CDT	75.2	3.45	clear
08/06/2021 04:56 CDT	75.02	3.45	clear
08/06/2021 05:00 CDT	75.2	3.45	clear
08/06/2021 05:05 CDT	75.2	3.45	thin scattered
08/06/2021 05:10 CDT	73.4	3.45	thin scattered
08/06/2021 05:15 CDT	73.4	0	thin scattered
08/06/2021 05:20 CDT	75.2	3.45	scattered
08/06/2021 05:25 CDT	75.2	3.45	broken
08/06/2021 05:30 CDT	73.4	0	broken
08/06/2021 05:35 CDT	75.2	0	broken
08/06/2021 05:40 CDT	75.2	3.45	overcast
08/06/2021 05:45 CDT	75.2	4.6	overcast
08/06/2021 05:50 CDT	75.2	3.45	overcast
08/06/2021 05:55 CDT	75.2	5.75	overcast
08/06/2021 05:56 CDT	75.02	5.75	overcast
08/06/2021 06:00 CDT	75.2	6.91	overcast
08/06/2021 06:05 CDT	75.2	6.91	overcast
08/06/2021 06:10 CDT	75.2	5.75	overcast
08/06/2021 06:15 CDT	75.2	6.91	overcast
08/06/2021 06:20 CDT	75.2	6.91	overcast
08/06/2021 06:25 CDT	75.2	6.91	overcast
08/06/2021 06:30 CDT	75.2	8.06	overcast
08/06/2021 06:35 CDT	75.2	6.91	overcast
08/06/2021 06:40 CDT	75.2	6.91	overcast

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/06/2021 19:25 CDT	80.6	4.6	scattered
08/06/2021 19:30 CDT	80.6	4.6	thin scattered
08/06/2021 19:35 CDT	80.6	5.75	thin scattered
08/06/2021 19:40 CDT	80.6	5.75	scattered
08/06/2021 19:45 CDT	80.6	4.6	broken
08/06/2021 19:50 CDT	80.6	4.6	scattered
08/06/2021 19:55 CDT	80.6	0	scattered
08/06/2021 19:56 CDT	80.06	0	scattered
08/06/2021 20:00 CDT	78.8	0	scattered
08/06/2021 20:05 CDT	77	0	broken
08/06/2021 20:10 CDT	78.8	0	broken
08/06/2021 20:15 CDT	78.8	0	broken
08/06/2021 20:20 CDT	75.2	0	scattered
08/06/2021 20:25 CDT	77	0	scattered
08/06/2021 20:30 CDT	77	0	scattered
08/06/2021 20:35 CDT	77	0	scattered
08/06/2021 20:40 CDT	78.8	4.6	scattered
08/06/2021 20:45 CDT	78.8	4.6	thin scattered
08/06/2021 20:50 CDT	78.8	3.45	thin scattered
08/06/2021 20:55 CDT	78.8	0	thin scattered
08/06/2021 20:56 CDT	78.98	0	thin scattered
08/06/2021 21:00 CDT	78.8	0	thin scattered
08/06/2021 21:05 CDT	75.2	4.6	clear
08/06/2021 21:10 CDT	78.8	4.6	clear
08/06/2021 21:15 CDT	78.8	3.45	clear
08/06/2021 21:20 CDT	78.8	5.75	clear
08/06/2021 21:25 CDT	77	5.75	clear
08/06/2021 21:30 CDT	77	4.6	clear
08/06/2021 21:35 CDT	78.8	4.6	clear
08/06/2021 21:40 CDT	78.8	4.6	clear
08/06/2021 21:45 CDT	78.8	4.6	clear
08/06/2021 21:50 CDT	78.8	4.6	clear
08/06/2021 21:55 CDT	78.8	5.75	clear
08/06/2021 21:56 CDT	78.08	4.6	clear
08/06/2021 22:00 CDT	78.8	6.91	clear
08/06/2021 22:05 CDT	78.8	4.6	clear
08/06/2021 22:10 CDT	77	3.45	clear
08/06/2021 22:20 CDT	73.4	4.6	clear
08/06/2021 22:25 CDT	73.4	4.6	clear
08/06/2021 22:30 CDT	73.4	4.6	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/06/2021 22:35 CDT	73.4	3.45	clear
08/06/2021 22:40 CDT	73.4	0	clear
08/06/2021 22:45 CDT	75.2	0	clear
08/07/2021 05:50 CDT		0	
08/07/2021 05:51 CDT	69.8	0	
08/07/2021 05:55 CDT	69.8	0	
08/07/2021 05:56 CDT	69.98	0	mist
08/07/2021 06:00 CDT	69.8	0	mist
08/07/2021 06:03 CDT	69.8	0	mist
08/07/2021 06:05 CDT	69.8	0	mist
08/07/2021 06:10 CDT	69.8	0	mist
08/07/2021 06:15 CDT	69.8	0	mist
08/07/2021 06:20 CDT	69.8	0	mist
08/07/2021 06:25 CDT	69.8	0	mist
08/07/2021 06:30 CDT	69.8	0	mist
08/07/2021 06:31 CDT	69.8	0	mist
08/07/2021 06:35 CDT	69.8	0	mist
08/07/2021 06:40 CDT	71.6	3.45	mist
08/07/2021 19:20 CDT	84.2	0	clear
08/07/2021 19:25 CDT	84.2	0	clear
08/07/2021 19:30 CDT	84.2	0	clear
08/07/2021 19:35 CDT	82.4	0	clear
08/07/2021 19:40 CDT	82.4	0	clear
08/07/2021 19:45 CDT	82.4	0	clear
08/07/2021 19:50 CDT	82.4	0	clear
08/07/2021 19:55 CDT	80.6	0	clear
08/07/2021 19:56 CDT	80.96	0	clear
08/07/2021 20:00 CDT	80.6	0	clear
08/07/2021 20:05 CDT	80.6	0	clear
08/07/2021 20:10 CDT	80.6	0	clear
08/07/2021 20:15 CDT	80.6	0	clear
08/07/2021 20:20 CDT	80.6	0	clear
08/07/2021 20:25 CDT	80.6	0	clear
08/07/2021 20:30 CDT	80.6	0	clear
08/07/2021 20:35 CDT	80.6	0	clear
08/07/2021 20:40 CDT	78.8	0	clear
08/07/2021 20:45 CDT	78.8	0	clear
08/07/2021 20:50 CDT	78.8	0	clear
08/07/2021 20:55 CDT	77	0	clear
08/07/2021 20:56 CDT	77	0	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/07/2021 21:00 CDT	77	0	clear
08/07/2021 21:05 CDT	78.8	0	clear
08/07/2021 21:10 CDT	78.8	0	clear
08/07/2021 21:15 CDT	78.8	0	clear
08/07/2021 21:25 CDT	77	0	clear
08/07/2021 21:30 CDT	77	0	clear
08/07/2021 21:35 CDT	77	0	clear
08/07/2021 21:40 CDT	77	0	clear
08/07/2021 21:45 CDT	77	0	clear
08/07/2021 21:50 CDT	75.2	0	clear
08/07/2021 21:55 CDT	75.2	0	clear
08/07/2021 21:56 CDT	75.92	0	clear
08/07/2021 22:00 CDT	75.2	0	clear
08/07/2021 22:05 CDT	75.2	0	clear
08/07/2021 22:10 CDT	75.2	3.45	clear
08/07/2021 22:15 CDT	75.2	3.45	clear
08/07/2021 22:20 CDT	73.4	3.45	clear
08/07/2021 22:25 CDT	73.4	3.45	clear
08/07/2021 22:30 CDT	73.4	3.45	clear
08/07/2021 22:35 CDT	73.4	3.45	clear
08/07/2021 22:40 CDT	73.4	3.45	clear
08/07/2021 22:45 CDT	73.4	3.45	clear
08/07/2021 22:50 CDT	73.4	0	clear
08/07/2021 22:55 CDT	73.4	0	clear
08/07/2021 22:56 CDT	73.94	0	clear
08/07/2021 23:00 CDT	73.4	3.45	clear
08/07/2021 23:05 CDT	73.4	0	clear
08/07/2021 23:10 CDT	73.4	3.45	clear
08/07/2021 23:15 CDT	73.4	0	clear
08/07/2021 23:20 CDT	73.4	0	clear
08/07/2021 23:25 CDT	73.4	0	clear
08/07/2021 23:30 CDT	73.4	0	clear
08/07/2021 23:35 CDT	73.4	0	clear
08/07/2021 23:40 CDT	73.4	0	clear
08/07/2021 23:50 CDT	73.4	3.45	clear
08/07/2021 23:55 CDT	73.4	3.45	clear
08/07/2021 23:56 CDT	73.04	3.45	clear
08/08/2021 00:00 CDT	73.4	0	clear
08/08/2021 00:05 CDT	73.4	0	clear
08/08/2021 00:10 CDT	73.4	0	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/08/2021 00:15 CDT	71.6	0	clear
08/08/2021 00:20 CDT	71.6	0	clear
08/08/2021 00:25 CDT	73.4	0	clear
08/08/2021 00:30 CDT	71.6	0	clear
08/08/2021 00:35 CDT	73.4	0	clear
08/08/2021 00:40 CDT	73.4	0	clear
08/08/2021 00:45 CDT	73.4	0	clear
08/08/2021 00:50 CDT	73.4	0	clear
08/08/2021 00:55 CDT	73.4	0	clear
08/08/2021 00:56 CDT	73.04	0	clear
08/08/2021 01:00 CDT	73.4	0	clear
08/08/2021 01:05 CDT	73.4	0	clear
08/08/2021 01:10 CDT	73.4	0	clear
08/08/2021 01:15 CDT	73.4	0	clear
08/08/2021 01:20 CDT	71.6	0	clear
08/08/2021 01:25 CDT	71.6	0	clear
08/08/2021 01:30 CDT	71.6	0	clear
08/08/2021 01:35 CDT	71.6	0	clear
08/08/2021 01:40 CDT	71.6	0	clear
08/08/2021 01:45 CDT	71.6	0	clear
08/08/2021 01:50 CDT	71.6	0	clear
08/08/2021 01:55 CDT	71.6	0	clear
08/08/2021 01:56 CDT	71.06	0	clear
08/08/2021 02:00 CDT	71.6	0	clear
08/08/2021 02:05 CDT	71.6	0	clear
08/08/2021 02:10 CDT	71.6	0	clear
08/08/2021 02:15 CDT	71.6	0	clear
08/08/2021 02:20 CDT	71.6	0	clear
08/08/2021 02:25 CDT	71.6	0	clear
08/08/2021 02:30 CDT	71.6	0	clear
08/08/2021 02:35 CDT	71.6	0	clear
08/08/2021 02:40 CDT	71.6	0	clear
08/08/2021 02:45 CDT	71.6	0	clear
08/08/2021 02:50 CDT	71.6	0	clear
08/08/2021 02:55 CDT	71.6	0	clear
08/08/2021 02:56 CDT	71.96	0	clear
08/08/2021 03:00 CDT	71.6	0	clear
08/08/2021 03:05 CDT	71.6	0	clear
08/08/2021 03:10 CDT	71.6	0	mist
08/08/2021 03:15 CDT	69.8	0	mist

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/08/2021 03:20 CDT	69.8	0	clear
08/08/2021 03:25 CDT	69.8	0	clear
08/08/2021 03:30 CDT	69.8	0	clear
08/08/2021 03:35 CDT	69.8	0	clear
08/08/2021 03:40 CDT	71.6	0	clear
08/08/2021 03:45 CDT	71.6	0	clear
08/08/2021 03:50 CDT	71.6	3.45	clear
08/08/2021 03:55 CDT	71.6	0	clear
08/08/2021 03:56 CDT	71.06	0	clear
08/08/2021 04:00 CDT	69.8	0	clear
08/08/2021 04:05 CDT	69.8	0	clear
08/08/2021 04:10 CDT	71.6	0	clear
08/08/2021 04:15 CDT	71.6	0	clear
08/08/2021 04:20 CDT	69.8	0	clear
08/08/2021 04:25 CDT	69.8	0	clear
08/08/2021 04:30 CDT	69.8	0	clear
08/08/2021 04:35 CDT	69.8	0	clear
08/08/2021 04:40 CDT	69.8	0	clear
08/08/2021 04:45 CDT	71.6	0	clear
08/08/2021 04:50 CDT	71.6	0	clear
08/08/2021 04:55 CDT	71.6	3.45	clear
08/08/2021 04:56 CDT	71.06	3.45	clear
08/08/2021 05:00 CDT	71.6	3.45	clear
08/08/2021 05:05 CDT	69.8	3.45	clear
08/08/2021 05:10 CDT	69.8	3.45	clear
08/08/2021 05:15 CDT	69.8	3.45	clear
08/08/2021 05:20 CDT	69.8	3.45	clear
08/08/2021 05:25 CDT	71.6	4.6	clear
08/08/2021 05:30 CDT	71.6	0	clear
08/08/2021 05:35 CDT	71.6	0	clear
08/08/2021 05:40 CDT	71.6	0	clear
08/08/2021 05:45 CDT	71.6	0	clear
08/08/2021 05:50 CDT	71.6	0	clear
08/08/2021 05:55 CDT	71.6	0	clear
08/08/2021 05:56 CDT	71.06	0	clear
08/08/2021 06:00 CDT	69.8	0	mist
08/08/2021 06:03 CDT	69.8	0	mist
08/08/2021 06:05 CDT	69.8	0	mist
08/08/2021 06:10 CDT	69.8	0	mist
08/08/2021 06:15 CDT	69.8	0	mist

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/08/2021 06:20 CDT	69.8	0	mist
08/08/2021 06:25 CDT	69.8	0	mist
08/08/2021 06:30 CDT	69.8	0	mist
08/08/2021 06:35 CDT	71.6	0	mist
08/08/2021 06:37 CDT	71.6	0	mist
08/08/2021 06:40 CDT	71.6	0	mist
08/08/2021 19:20 CDT	84.2	5.75	clear
08/08/2021 19:25 CDT	84.2	5.75	clear
08/08/2021 19:35 CDT	82.4	5.75	clear
08/08/2021 19:40 CDT	82.4	5.75	clear
08/08/2021 19:45 CDT	82.4	5.75	clear
08/08/2021 19:50 CDT	82.4	6.91	clear
08/08/2021 19:55 CDT	82.4	6.91	clear
08/08/2021 19:56 CDT	82.94	6.91	clear
08/08/2021 20:00 CDT	82.4	6.91	clear
08/08/2021 20:05 CDT	82.4	5.75	clear
08/08/2021 20:10 CDT	82.4	6.91	clear
08/08/2021 20:15 CDT	82.4	5.75	clear
08/08/2021 20:20 CDT	82.4	5.75	thin scattered
08/08/2021 20:25 CDT	82.4	5.75	scattered
08/08/2021 20:30 CDT	82.4	5.75	broken
08/08/2021 20:35 CDT	82.4	5.75	broken
08/08/2021 20:40 CDT	82.4	5.75	broken
08/08/2021 20:45 CDT	82.4	5.75	scattered
08/08/2021 20:50 CDT	80.6	5.75	scattered
08/08/2021 20:55 CDT	80.6	5.75	thin scattered
08/08/2021 20:56 CDT	80.96	5.75	thin scattered
08/08/2021 21:00 CDT	80.6	6.91	thin scattered
08/08/2021 21:05 CDT	80.6	5.75	clear
08/08/2021 21:10 CDT	80.6	5.75	clear
08/08/2021 21:15 CDT	80.6	5.75	clear
08/08/2021 21:20 CDT	80.6	5.75	clear
08/08/2021 21:25 CDT	80.6	6.91	clear
08/08/2021 21:30 CDT	80.6	5.75	clear
08/08/2021 21:35 CDT	80.6	5.75	clear
08/08/2021 21:40 CDT	80.6	5.75	clear
08/08/2021 21:45 CDT	80.6	5.75	clear
08/08/2021 21:50 CDT	80.6	4.6	clear
08/08/2021 21:55 CDT	80.6	4.6	clear
08/08/2021 21:56 CDT	80.96	4.6	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/08/2021 22:00 CDT	80.6	4.6	clear
08/08/2021 22:05 CDT	80.6	4.6	clear
08/08/2021 22:10 CDT	80.6	5.75	clear
08/08/2021 22:15 CDT	80.6	5.75	clear
08/08/2021 22:20 CDT	80.6	5.75	clear
08/08/2021 22:30 CDT	80.6	5.75	clear
08/08/2021 22:35 CDT	80.6	5.75	clear
08/08/2021 22:40 CDT	78.8	6.91	clear
08/08/2021 22:45 CDT	78.8	6.91	clear
08/08/2021 22:50 CDT	78.8	6.91	clear
08/08/2021 22:55 CDT	78.8	5.75	clear
08/08/2021 22:56 CDT	78.98	5.75	clear
08/08/2021 23:00 CDT	78.8	5.75	clear
08/08/2021 23:05 CDT	78.8	5.75	clear
08/08/2021 23:10 CDT	78.8	5.75	clear
08/08/2021 23:15 CDT	78.8	5.75	clear
08/08/2021 23:20 CDT	78.8	5.75	clear
08/08/2021 23:25 CDT	78.8	6.91	clear
08/08/2021 23:30 CDT	78.8	5.75	clear
08/08/2021 23:35 CDT	78.8	5.75	clear
08/08/2021 23:40 CDT	78.8	4.6	clear
08/08/2021 23:45 CDT	77	4.6	clear
08/08/2021 23:50 CDT	77	5.75	clear
08/08/2021 23:55 CDT	77	5.75	clear
08/08/2021 23:56 CDT	77	5.75	clear
08/09/2021 00:00 CDT	77	5.75	clear
08/09/2021 00:05 CDT	78.8	5.75	clear
08/09/2021 00:10 CDT	78.8	6.91	clear
08/09/2021 00:15 CDT	78.8	6.91	clear
08/09/2021 00:20 CDT	77	5.75	clear
08/09/2021 00:25 CDT	75.2	6.91	clear
08/09/2021 00:30 CDT	75.2	5.75	clear
08/09/2021 00:35 CDT	75.2	5.75	clear
08/09/2021 00:40 CDT	75.2	5.75	clear
08/09/2021 00:45 CDT	75.2	5.75	clear
08/09/2021 00:50 CDT	75.2	6.91	clear
08/09/2021 00:55 CDT	75.2	8.06	clear
08/09/2021 00:56 CDT	75.92	8.06	clear
08/09/2021 01:00 CDT	75.2	6.91	clear
08/09/2021 01:05 CDT	75.2	5.75	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/09/2021 01:10 CDT	75.2	5.75	clear
08/09/2021 01:15 CDT	77	5.75	clear
08/09/2021 01:20 CDT	77	6.91	clear
08/09/2021 01:25 CDT	77	5.75	clear
08/09/2021 01:30 CDT	75.2	5.75	clear
08/09/2021 01:35 CDT	75.2	5.75	clear
08/09/2021 01:40 CDT	75.2	5.75	clear
08/09/2021 01:45 CDT	75.2	5.75	clear
08/09/2021 01:50 CDT	75.2	5.75	clear
08/09/2021 01:55 CDT	75.2	6.91	clear
08/09/2021 01:56 CDT	75.92	6.91	clear
08/09/2021 02:00 CDT	75.2	6.91	clear
08/09/2021 02:05 CDT	75.2	5.75	clear
08/09/2021 02:10 CDT	75.2	5.75	clear
08/09/2021 02:15 CDT	75.2	3.45	clear
08/09/2021 02:20 CDT	75.2	3.45	clear
08/09/2021 02:25 CDT	75.2	3.45	clear
08/09/2021 02:30 CDT	75.2	4.6	clear
08/09/2021 02:35 CDT	75.2	4.6	clear
08/09/2021 02:40 CDT	75.2	4.6	clear
08/09/2021 02:45 CDT	75.2	4.6	clear
08/09/2021 02:50 CDT	75.2	3.45	clear
08/09/2021 02:55 CDT	75.2	3.45	clear
08/09/2021 02:56 CDT	75.92	3.45	clear
08/09/2021 03:00 CDT	75.2	3.45	clear
08/09/2021 03:05 CDT	75.2	3.45	clear
08/09/2021 03:10 CDT	73.4	4.6	clear
08/09/2021 03:15 CDT	73.4	5.75	clear
08/09/2021 03:20 CDT	73.4	5.75	clear
08/09/2021 03:25 CDT	73.4	4.6	clear
08/09/2021 03:30 CDT	73.4	4.6	clear
08/09/2021 03:35 CDT	73.4	4.6	clear
08/09/2021 03:40 CDT	73.4	4.6	clear
08/09/2021 03:45 CDT	73.4	4.6	clear
08/09/2021 03:50 CDT	73.4	5.75	clear
08/09/2021 03:55 CDT	73.4	5.75	clear
08/09/2021 03:56 CDT	73.94	5.75	clear
08/09/2021 04:00 CDT	73.4	5.75	clear
08/09/2021 04:05 CDT	73.4	5.75	clear
08/09/2021 04:10 CDT	73.4	5.75	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/09/2021 04:15 CDT	75.2	4.6	clear
08/09/2021 04:20 CDT	73.4	4.6	clear
08/09/2021 04:25 CDT	73.4	5.75	clear
08/09/2021 04:30 CDT	73.4	5.75	clear
08/09/2021 04:35 CDT	73.4	5.75	clear
08/09/2021 04:40 CDT	73.4	5.75	clear
08/09/2021 04:45 CDT	73.4	4.6	clear
08/09/2021 04:50 CDT	73.4	4.6	clear
08/09/2021 04:55 CDT	73.4	4.6	clear
08/09/2021 04:56 CDT	73.94	4.6	clear
08/09/2021 05:00 CDT	73.4	3.45	clear
08/09/2021 05:05 CDT	73.4	3.45	clear
08/09/2021 05:10 CDT	73.4	0	clear
08/09/2021 05:15 CDT	73.4	0	clear
08/09/2021 05:20 CDT	75.2	0	clear
08/09/2021 05:25 CDT	73.4	0	clear
08/09/2021 05:30 CDT	73.4	0	clear
08/09/2021 05:35 CDT	73.4	0	clear
08/09/2021 05:45 CDT	75.2	0	clear
08/09/2021 05:50 CDT	73.4	0	clear
08/09/2021 05:55 CDT	73.4	0	clear
08/09/2021 05:56 CDT	73.94	0	clear
08/09/2021 06:00 CDT	73.4	0	clear
08/09/2021 06:10 CDT	75.2	0	clear
08/09/2021 06:15 CDT	73.4	0	clear
08/09/2021 06:20 CDT	73.4	0	clear
08/09/2021 06:25 CDT	75.2	0	clear
08/09/2021 06:30 CDT	75.2	0	clear
08/09/2021 06:35 CDT	75.2	0	clear
08/09/2021 06:40 CDT	75.2	0	clear
08/09/2021 19:20 CDT	86	4.6	clear
08/09/2021 19:25 CDT	86	5.75	clear
08/09/2021 19:30 CDT	86	5.75	clear
08/09/2021 19:35 CDT	86	4.6	clear
08/09/2021 19:40 CDT	86	5.75	clear
08/09/2021 19:45 CDT	84.2	5.75	clear
08/09/2021 19:50 CDT	84.2	5.75	clear
08/09/2021 19:55 CDT	84.2	5.75	clear
08/09/2021 19:56 CDT	84.92	4.6	clear
08/09/2021 20:00 CDT	84.2	5.75	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/09/2021 20:05 CDT	84.2	5.75	clear
08/09/2021 20:10 CDT	84.2	4.6	clear
08/09/2021 20:15 CDT	84.2	5.75	clear
08/09/2021 20:20 CDT	84.2	5.75	clear
08/09/2021 20:25 CDT	84.2	5.75	clear
08/09/2021 20:30 CDT	84.2	5.75	clear
08/09/2021 20:35 CDT	84.2	4.6	clear
08/09/2021 20:40 CDT	84.2	4.6	clear
08/09/2021 20:45 CDT	84.2	4.6	clear
08/09/2021 20:50 CDT	82.4	4.6	clear
08/09/2021 20:55 CDT	82.4	3.45	clear
08/09/2021 20:56 CDT	82.94	3.45	clear
08/09/2021 21:00 CDT	82.4	4.6	clear
08/09/2021 21:05 CDT	82.4	3.45	clear
08/09/2021 21:10 CDT	82.4	3.45	clear
08/09/2021 21:15 CDT	80.6	4.6	clear
08/09/2021 21:20 CDT	80.6	3.45	clear
08/09/2021 21:25 CDT	80.6	4.6	clear
08/09/2021 21:30 CDT	80.6	5.75	clear
08/09/2021 21:35 CDT	80.6	5.75	clear
08/09/2021 21:40 CDT	80.6	5.75	clear
08/09/2021 21:45 CDT	80.6	5.75	clear
08/09/2021 21:50 CDT	80.6	4.6	clear
08/09/2021 21:55 CDT	80.6	4.6	clear
08/09/2021 21:56 CDT	80.96	4.6	clear
08/09/2021 22:00 CDT	80.6	4.6	clear
08/09/2021 22:05 CDT	80.6	4.6	clear
08/09/2021 22:10 CDT	80.6	4.6	clear
08/09/2021 22:15 CDT	80.6	4.6	clear
08/09/2021 22:20 CDT	80.6	9.21	clear
08/09/2021 22:25 CDT	82.4	8.06	clear
08/09/2021 22:30 CDT	82.4	5.75	clear
08/09/2021 22:35 CDT	82.4	6.91	clear
08/09/2021 22:40 CDT	82.4	5.75	clear
08/09/2021 22:45 CDT	82.4	6.91	clear
08/09/2021 22:50 CDT	82.4	5.75	clear
08/09/2021 22:55 CDT	82.4	6.91	clear
08/09/2021 22:56 CDT	82.04	6.91	clear
08/09/2021 23:00 CDT	82.4	8.06	clear
08/09/2021 23:05 CDT	80.6	6.91	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/09/2021 23:10 CDT	80.6	6.91	clear
08/09/2021 23:15 CDT	80.6	5.75	clear
08/09/2021 23:20 CDT	80.6	8.06	clear
08/09/2021 23:25 CDT	80.6	6.91	clear
08/09/2021 23:30 CDT	80.6	9.21	clear
08/09/2021 23:35 CDT	80.6	8.06	clear
08/09/2021 23:40 CDT	80.6	6.91	clear
08/09/2021 23:45 CDT	80.6	6.91	clear
08/09/2021 23:50 CDT	80.6	8.06	clear
08/09/2021 23:55 CDT	80.6	8.06	clear
08/09/2021 23:56 CDT	80.06	8.06	clear
08/10/2021 00:00 CDT	80.6	8.06	clear
08/10/2021 00:05 CDT	80.6	9.21	clear
08/10/2021 00:10 CDT	80.6	9.21	clear
08/10/2021 00:15 CDT	80.6	8.06	clear
08/10/2021 00:20 CDT	80.6	8.06	clear
08/10/2021 00:25 CDT	80.6	6.91	clear
08/10/2021 00:30 CDT	78.8	6.91	clear
08/10/2021 00:35 CDT	78.8	5.75	clear
08/10/2021 00:40 CDT	78.8	8.06	clear
08/10/2021 00:45 CDT	78.8	6.91	clear
08/10/2021 00:50 CDT	78.8	5.75	clear
08/10/2021 00:55 CDT	78.8	5.75	clear
08/10/2021 00:56 CDT	78.98	5.75	clear
08/10/2021 01:00 CDT	78.8	6.91	clear
08/10/2021 01:05 CDT	78.8	5.75	clear
08/10/2021 01:10 CDT	78.8	4.6	clear
08/10/2021 01:15 CDT	77	4.6	clear
08/10/2021 01:20 CDT	78.8	5.75	clear
08/10/2021 01:25 CDT	78.8	6.91	clear
08/10/2021 01:30 CDT	78.8	8.06	clear
08/10/2021 01:35 CDT	78.8	5.75	clear
08/10/2021 01:40 CDT	78.8	6.91	clear
08/10/2021 01:45 CDT	78.8	5.75	clear
08/10/2021 01:50 CDT	77	6.91	clear
08/10/2021 01:55 CDT	77	8.06	clear
08/10/2021 01:56 CDT	77	6.91	clear
08/10/2021 02:00 CDT	77	5.75	clear
08/10/2021 02:05 CDT	77	4.6	clear
08/10/2021 02:10 CDT	77	4.6	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/10/2021 02:15 CDT	77	4.6	clear
08/10/2021 02:20 CDT	75.2	4.6	clear
08/10/2021 02:25 CDT	75.2	4.6	clear
08/10/2021 02:30 CDT	77	4.6	clear
08/10/2021 02:35 CDT	77	4.6	clear
08/10/2021 02:40 CDT	75.2	5.75	clear
08/10/2021 02:45 CDT	75.2	4.6	clear
08/10/2021 02:50 CDT	77	4.6	clear
08/10/2021 02:55 CDT	77	4.6	clear
08/10/2021 02:56 CDT	77	4.6	clear
08/10/2021 03:00 CDT	77	4.6	clear
08/10/2021 03:05 CDT	77	3.45	clear
08/10/2021 03:10 CDT	77	0	clear
08/10/2021 03:15 CDT	77	0	clear
08/10/2021 03:20 CDT	77	0	clear
08/10/2021 03:25 CDT	77	3.45	clear
08/10/2021 03:30 CDT	75.2	0	clear
08/10/2021 03:35 CDT	75.2	0	clear
08/10/2021 03:40 CDT	75.2	0	clear
08/10/2021 03:45 CDT	75.2	0	clear
08/10/2021 03:50 CDT	75.2	0	clear
08/10/2021 03:55 CDT	75.2	0	clear
08/10/2021 03:56 CDT	75.92	0	clear
08/10/2021 04:00 CDT	75.2	0	clear
08/10/2021 04:05 CDT	75.2	0	clear
08/10/2021 04:10 CDT	75.2	3.45	clear
08/10/2021 04:15 CDT	75.2	0	clear
08/10/2021 04:20 CDT	75.2	4.6	clear
08/10/2021 04:25 CDT	75.2	4.6	clear
08/10/2021 04:30 CDT	77	4.6	clear
08/10/2021 04:35 CDT	77	3.45	clear
08/10/2021 04:40 CDT	77	0	clear
08/10/2021 04:45 CDT	75.2	0	clear
08/10/2021 04:50 CDT	75.2	3.45	clear
08/10/2021 04:55 CDT	75.2	0	clear
08/10/2021 04:56 CDT	75.92	0	clear
08/10/2021 05:00 CDT	75.2	3.45	clear
08/10/2021 05:05 CDT	75.2	3.45	clear
08/10/2021 05:10 CDT	75.2	3.45	clear
08/10/2021 05:15 CDT	75.2	0	clear

Date & Time	Temperature (°F)	Wind Speed (mph)	Weather Summary
08/10/2021 05:20 CDT	75.2	0	clear
08/10/2021 05:25 CDT	75.2	3.45	clear
08/10/2021 05:30 CDT	73.4	5.75	clear
08/10/2021 05:35 CDT	75.2	6.91	clear
08/10/2021 05:45 CDT	75.2	5.75	clear
08/10/2021 05:50 CDT	75.2	6.91	clear
08/10/2021 05:55 CDT	75.2	5.75	clear
08/10/2021 05:56 CDT	75.92	5.75	clear
08/10/2021 06:00 CDT	75.2	4.6	clear
08/10/2021 06:05 CDT	75.2	3.45	clear
08/10/2021 06:10 CDT	75.2	3.45	clear
08/10/2021 06:15 CDT	75.2	5.75	clear
08/10/2021 06:20 CDT	75.2	4.6	clear
08/10/2021 06:25 CDT	75.2	4.6	clear
08/10/2021 06:30 CDT	75.2	4.6	clear
08/10/2021 06:35 CDT	75.2	3.45	clear
08/10/2021 06:40 CDT	75.2	3.45	clear

Source: Weather data for KGTR from www.mesowest.utah.edu



Golden Triangle II Acoustic Bat Survey				
Name/Education	Experience	Project Role		
Josiah Maine B.S. Wildlife Biology M.S. Zoology	10 years conducting bat surveys, acoustic bat identification, and environmental permitting	Lead Bat Biologist; Survey Development, Acoustic Site Selection, Data Analytics, Primary Report Author		
Grant Dickerson B.S Wildlife Biology M.S. Biology	5 years conducting bat surveys	Bat Biologist; Site Selection and Acoustic Monitor Deployment		
Brooke Hines B.S. Wildlife Biology M.S. Biology, Applied Ecology	21 years in protected bat species consultation to include, habitat and biological assessments, writing state and federal guidance for bat surveys, and coordination with state and federal agencies on policy implementation	Bat Biologist; Senior Technical Lead/Review of Bat Survey Plan/Report		
Jesse Brown B.A. Biology M.S. Biology	12 years in environmental permitting and protected species evaluations	Field Survey Coordination, Habitat Assessment, Acoustic Monitor Deployment, Deputy PM		

# Josiah J. Maine

Curriculum Vitae 9400 Ward Pkwy Kansas City, MO 64114 jjmaine@burnsmcd.com 816-448-7519

#### **EDUCATION**

2014 Southern Illinois University, Carbondale, IL

M.S. Zoology, conferred December 2014

Thesis: Trophic Ecology of Insectivorous Bats in Agroecosystems

**2011** Kansas State University, Manhattan, KS

B.S. Fisheries, Wildlife, and Conservation Biology, conferred December 2011

BAT NETTING, TRACKING, AND ACOUSTIC EXPERIENCE

**Bat Biologist, ITC Cardinal-Hickory Creek Transmission Line Project**, Clayton County, IA, August 2020

Lead bat biologist responsible for conducting bat surveys for a potential transmission line that crossed a national wildlife refuge. Surveys included acoustic surveys at two sites and mist-net surveys at two sites. Species captured included little brown bat, tricolored bat, and eastern red bat.

Approximate time: 4 detector-nights, 4 nights of netting

**Bat Biologist, Duke UL06 Pipeline Replacement Project**, Campbell County, KY, June 2020 **Lead bat biologist** responsible for conducting acoustic presence/absence surveys for a pipeline replacement project. One acoustic site was surveyed for two detector-nights. Automated identification software indicated that northern long-eared bats may occur on the site. Follow-up manual vetting conducted by Josiah Maine identified calls that had characteristics consistent with northern long-eared bats.

Approximate time: 2 detector-nights

**Bat Biologist and Land Manager, Chariton Hills Conservation Bank,** Northeast Missouri, Jan. 2017 - Present

Lead bat biologist and land manager for the Chariton Hills Conservation Bank. Played a key role in the development of the bat conservation bank, which was be the first conservation bank for Region 3 of the U.S. Fish & Wildlife Service. Identified suitable properties and led the acoustic, mist-netting, and tracking studies on the bank properties, focused on Indiana bat and northern long-eared bat. Made significant contributions to the management plan for the bank. As the land manager, coordinates permitting, biological studies, forest management, property security, and recreational activities for the bank.

Approximate time: 14 netting survey nights, 14 nights of acoustics, 2 Indiana bats and 1 northern long-eared bat captured

Bat Biologist, Alta Call Center Project, Fort Myers, FL, October 2019

**Lead bat biologist** responsible for conducting Florida bonneted bat presence/absence surveys for a development site using acoustic equipment. This study allowed the client to evaluate potential risk of the project to bats, in coordination with the U.S. Fish & Wildlife Service. *Approximate time:* 18 detector-nights

# **Bat Biologist, Boone Area Improvements Transmission Line**, Boone and Kanawha Counties, WV, 2018

**Lead bat biologist** responsible for conducting presence/absence surveys for a potential transmission line. Surveys included acoustic surveys at 53 sites, with follow-up netting at 3 sites. Species captured included eastern small-footed bat, hoary bat, eastern red bat, and big brown bat. *Approximate time: 106 detector-nights, 9 nights of netting* 

#### Bat Biologist, Kossuth Wind Farm, Iowa, 2018

**Lead bat biologist** responsible for conducting presence/absence surveys for a wind farm site using acoustic equipment. Target species included northern long-eared bat, little brown bat, and tricolored bat. This study allowed the client to evaluate potential risk of the project to bats, in coordination with the U.S. Fish & Wildlife Service.

Approximate time: 32 detector-nights

# Bat Biologist, Bat Mitigation Property Assessment, Iowa, 2018

Lead bat biologist responsible for conducting presence/absence surveys for several potential bat mitigation properties using acoustic equipment. Target species included Indiana bat, northern long-eared bat, little brown bat, and tricolored bat. Acoustic bat identification was conducted with Kaleidoscope software followed by manual vetting. The purpose of the study was to determine species presence and suitability for multiple properties in Iowa that could serve as bat mitigation sites in the future.

Approximate time: 8 nights of acoustics, Indiana bats and northern long-eared bats identified

# Bat Biologist, Deuel Wind Project, Deuel County, SD, 2017

**Lead bat biologist** responsible for conducting a season-long bat study using acoustic equipment. The purpose of the bat study was to identify overall levels and seasonal patterns of bat activity within and below the rotor-swept area of a proposed wind farm. This study allowed the client to evaluate potential risk of the project to bats, in coordination with the U.S. Fish & Wildlife Service.

Approximate time: 90 nights of acoustics

## Bat Biologist, Whispering Willow Wind Project, Franklin County, IA, 2017

Lead bat biologist responsible for conducting bat study for a pre-construction wind project. The study included presence/absence surveys for the northern long-eared bat and a season-long bat study using acoustic equipment on meteorological towers. Northern long-eared bat acoustic data was processed through automated identification software, and manual vetting (qualitative identification by Josiah Maine) was conducted. Acoustic recordings were provided to USFWS to bolster data on the distribution of the northern long-eared bat in Iowa.

Approximate time: 200 nights of acoustics, northern long-eared bats identified

# Bat Biologist, Dakota Access Pipeline Project, Iowa, 2015

Assistant bat biologist responsible for performing sensitive species evaluations on a proposed crude oil pipeline crossing the states of Iowa and Illinois. The project included habitat assessments for the northern long-eared bat, Indiana bat, eastern massasauga and eastern prairie-fringed orchid. Assisted another permitted biologist with acoustic bat surveys followed by netting, telemetry, and emergence surveys for threatened and endangered bats.

Approximate time: 10 netting survey nights, 15 nights of acoustics, 5 Indiana bats and 3 northern long-eared bats captured

# Bat Telemetry Volunteer, Champaign, IL, Aug. 2014

**Assistant bat biologist** for netting and telemetry study. Duties: Capture, handle, and identify bats of Illinois, including Indiana bat, little brown bat, northern long-eared bat, red bat, evening bat, and big brown bat. Attach radio transmitters to multiple Indiana bats and northern long-eared bats. Track bats using radio telemetry. Use acoustic detectors to monitor bat activity. *Approximate time: 4 net survey nights, 3 northern long-eared bats captured* 

# **Graduate Research Assistant, Southern Illinois University**, Carbondale, IL, Aug. 2012 to Dec. 2014

Assistant bat biologist for netting surveys and lead bat biologist for acoustic surveys. Duties: Capture, handle, identify, and band bats of southern Illinois, including Indiana bat, northern long-eared bat, gray bat, little brown bat, southeastern bat, evening bat, big brown bat, tricolored bat, and red bat. Collect bat feces for diet analysis and perform acoustic bat surveys. Monitor bat activity using acoustic bat detectors, bat identification software, and qualitative ID. Design a large-scale experiment to test the strength of crop pest suppression by bats. Approximate time: 10 net survey nights, 100+ nights of acoustics, 2 Indiana bats and 2 northern long-eared bats independently identified

**Bat Survey Technician, Sanders Environmental**, Belefonte, PA, May 2010 to Aug. 2010 **Assistant bat biologist** for netting surveys. Duties: Capture, handle, and identify bats of Pennsylvania and West Virginia, including northern long-eared bat, little brown bat, big brown bat, red bat, hoary bat, Rafinesque's big-eared bat, and Virginia big-eared bat. *Approximate time:* 60 net survey nights, 10 northern long-eared bats captured

#### **BAT HIBERNACULA SURVEYS**

**Fall Swarm Survey Volunteer**, Great Scott Cave and Pilot Knob Mine, MO, October 2019 **Assistant bat biologist** for fall swarm surveys at two major Indiana bat and gray bat hibernacula. Duties: set up harp traps, capture, handle, and identify Indiana bat, little brown bat, gray bat, and eastern small-footed bat. *Approximate time:* 8 hrs, 200+ Indiana bats and 16 gray bats independently identified

## Hibernacula Survey Volunteer, Wolf River Cave, TN, March 2014

**Assistant bat biologist** for hibernaculum survey. Duties: Capture, handle, identify, and WNS-swab Indiana bat, little brown bat, northern long-eared bat, and tricolored bat in the second largest Indiana bat hibernaculum in Tennessee. Observe many species of hibernating bats, including Rafinesque's big-eared bat and gray bat. *Approximate time: 6 hrs, 12 Indiana bats and 10 northern long-eared bats independently identified* 

#### Hibernacula Survey Volunteer, Blackball Mine, IL, Dec. 2013

**Assistant bat biologist** for hibernaculum survey. Capture and transport hibernating big brown bats for a lab-based experiment. Visually identify and count hibernating bats, including big brown bat, tricolored bat, northern long-eared bat, little brown bat, and Indiana bat. *Approximate time:* 6 hrs

**Hibernacula Survey Volunteer,** Shawnee National Forest, IL, Feb. 2013 **Assistant bat biologist** for hibernaculum survey. Conduct a winter mine survey. Capture and transport hibernating big brown bats for a lab-based experiment. Visually identify and count

hibernating bats, including big brown bat, Rafinesque's big-eared bat, northern long-eared bat, southeastern bat, gray bat, and Indiana bat.

Approximate time: 6 hrs

#### **EMPLOYMENT**

#### 2015-Present Staff Environmental Scientist, Burns & McDonnell, Kansas City, MO

As a bat biologist, supervise and conduct threatened and endangered bat surveys, including habitat assessments, acoustic surveys, mist-netting, radio-tracking, and roost emergence counts.

Conduct habitat assessments and consultations assessing the impacts of proposed infrastructure projects on a variety of protected species across the U.S. Write Bird and Bat Conservation Strategies for wind energy facilities, analyzing ecological data and providing recommendations regarding best management practices to avoid impacts to bird and bat populations.

#### 2012-2014 Graduate Research Assistant, Southern Illinois University, Carbondale, IL

Duties: Capture, handle, identify, and band bats of southern Illinois, including Indiana bat, northern long-eared bat, little brown bat, gray bat, southeastern bat, Rafinesque's big-eared bat, evening bat, big brown bat, tricolored bat, and eastern red bat. Collect bat feces for diet analysis and perform acoustic bat surveys. Monitor bat activity using acoustic bat detectors, bat identification software, and qualitative ID. Design a large-scale experiment to test the strength of crop pest suppression by bats. Assist surveys for bat populations and whitenose syndrome in caves and reclaimed mines of Illinois and Tennessee.

#### 2012 Research Technician, Virginia Tech, Vermillion, SD

Duties: Capture and band federally threatened Piping Plovers and federally endangered Least Terns.

## 2010-2012 Research Technician, Kansas State University, Manhattan, KS

Duties: Conduct surveys for fish and aquatic macroinvertebrates across KS and NM, identify invertebrates in the lab, and lead a study on the diet of predacious macroinvertebrates from the Gila River, NM.

#### **2010** Research Technician, Sanders Environmental, Belefonte, PA

Duties: Capture, handle, and identify bats of Pennsylvania and West Virginia, including northern long-eared bat, little brown bat, big brown bat, red bat, hoary bat, Rafinesque's big-eared bat, and Virginia big-eared bat. Apply forearm bands to several species, including Virginia big-eared bat. Affix radio transmitters and conduct radiotelemetry on multiple species, including Rafinesque's big-eared bat.

#### **PUBLICATIONS**

**Maine, J.** and J. Boyles. 2015. Bats initiate vital agroecological interactions in corn. *Proceedings of the National Academy of Sciences*. 112 (40): 12438-12443

**Maine, J.** and J. Boyles. 2015. Land cover influences dietary specialization of insectivorous bats globally. *Mammal Research*. 60 (4): 343-351

- Maas, B., D. S. Karp, S. Bumrungsri, K. Darras, D. Gonthier, J. C.-C. Huang, C. A. Lindell, J. Maine, L. Mestre, N. L. Michel, E. B. Morrison, I. Perfecto, S. M. Philpott, C. H. Sekercioglu, R. M. Silva, P. Taylor, T. Tscharntke, S. A. Van Bael, C. J. Whelan, and K. Williams-Guillén. 2015. Bird and Bat Predation Services in Tropical Forests and Agroforestry Landscapes. *Biological Reviews*.
- Neiles, B., C. S. Carey, A. Araujo, D. Burkhart, L. J. Kirschman, B. LaBumbard, S. LaGrange, J. Maine, A. M. Rombenso, M. N. Wood, and J. G. Boyles. 2015. Writing Your Way into High Impact Factor Journals. *Bulletin of the Ecological Society of America*. 96 (2): 312-316.
- **Maine, J.**, J. Whitney, and K. Gido. 2014. Dietary Overlap of Invertivorous Fishes and Macroinvertebrates in the Gila River, NM. *The Southwestern Naturalist*. 59 (2): 292-295.

#### REFERENCES

James (Gene) Gardner Senior Environmental Scientist Burns & McDonnell 573-353-2726 jegardner@burnsmcd.com

Dr. Justin Boyles Assistant Professor Southern Illinois University 618-453-6945 jgboyles@siu.edu

# **BROOKE A. HINES, MS CWB**

# **ENVIRONMENTAL PROJECT MANAGER**



Brooke is an environmental project manager with extensive experience in ESA Section 7 consultation, NEPA documentation, CWA coordination, and federal, state, and municipality environmental permitting. Lead senior biologist for effects analysis on listed bat species, habitat assessments, presence/absence surveys, and federal and state species-specific studies.

Brooke is co-author of acoustic publications focusing on presence/absence methodology.

Brooke has over 21 years of experience with the ESA Section 7 consultation for threatened and endangered bat species, in public and private sectors. She has extensive experience with habitat and biological assessments, writing state and federal guidance

# **EDUCATION**

- MS, Biology, Applied Ecology
- BS, Natural Resources Wildlife Management

# REGISTRATIONS

- ESA Section 10 T&E Recovery Permit/USFWS
- Certified Wildlife Biologist/The Wildlife Society
- ► Remote Pilot sUAS Part 107 Certification/FAA

YEARS WITH BURNS & MCDONNELL

21 YEARS OF EXPERIENCE

for bat surveys, and coordination with state and federal agencies on policy implementation. She also has extensive experience with bat population estimates which include state-wide bat hibernacula and summer emergence counts and acoustic surveys. Brooke has also overseen numerous research studies on White-nose Syndrome and co-authored acoustic publications. Brooke has worked on numerous ecological and energy development projects throughout the U.S. focusing on federally endangered bats.

Ms. Hines' published research focuses on acoustic methodology and probability of detection. She has extensive experience conducting bat habitat surveys, presence/absence surveys including mist-net, acoustic, and harp traps, and telemetry surveys. Surveys focus on federally-listed species that include the Indiana bat (*Myotis sodalis*), gray bat (*M. grisescens*), northern long-eared bat (M. septentrionalis), and Virginia big-eared bat (*Corynorhinus townsendii virginianus*). Ms. Hines holds a U.S. Fish and Wildlife Service (USFWS) Section 10(A)(1)(a) recovery permit (#TE31355B-3) to handle Indiana bats, gray bats, northern long-eared bats, Virginia big-eared bats, and Ozark big- eared bats (*C. t. ingens*).


Ms. Hines is familiar with multi-agency coordination, data management, technical report writing, and field crew management. Her experience includes many ecological field techniques, including habitat assessments, bridge surveys, mistnetting, cave, and portal surveys, as well as acoustic monitoring and bat call ID and analysis. Ms. Hines was a member of the writing team for the first USFWS Range-wide Indiana bat Summer Guidelines (2014) and has extensive experience with qualitative identification of bat calls to manually verify acoustic software outputs.





CREATE AMAZING.

Burns & McDonnell Atlanta Regional Office 3650 Mansell Road, Suite 300 Alpharetta, GA 30022 www.burnsmcd.com





400 West Summit Hill Drive, Knoxville, Tennessee 37902

April 13, 2021

Mr. Barry White Director Mississippi Department of Archives and History Historic preservation Division Post Office Box 571 Jackson, Mississippi 39205-0521

Dear Mr. White:

TENNESSEE VALLEY AUTHORITY (TVA), POWER PURCHASE AGREEMENT (PPA), GOLDEN TRIANGLE SOLAR, SECOND PHASE LOWNDES COUNTY, MISSISSIPPI (TVA TRACKING NUMBER – CID 79857) (33.39170, -88.61037)

TVA proposes to enter into a PPA with Orgis Energy for an approximately 150 megawatt (MW) solar photovoltaic (PV) generating facility and a 50 MW Battery Energy Storage System (BESS) in Lowndes County, Mississippi. The project would be located within seven parcels totaling 1,520 acres. TVA recommends that the area of potential effects (APE) be considered to be the area of proposed ground-disturbance, where physical effects could occur including the PV facility, associated access routes, interconnect, and the BESS, as well as areas within a half-mile radius of the project within which the project would be visible, where visual effects on aboveground resources could occur. In order to obtain maximum flexibility in design to avoid both cultural and environmental resources, the archaeological survey area consists of the entire 1,520 acres.

Orgis Energy contracted with Burns & McDonnell to conduct a Phase I Cultural Resources survey. For your review, please find Burn & McDonnell's scope of work (SOW) for the Phase I Cultural Resources survey enclosed. Pursuant to 36 CFR § 800.4(b)(1), TVA finds that the SOW presented here represents a reasonable and good faith effort to carry out identification efforts.

By this letter, TVA is initiating consultation regarding the proposed undertaking. TVA is proposing to conduct a Phase I Cultural Resources survey of the APE as described in the enclosed SOW. In addition, TVA is proposing to have a kick-off meeting with all parties prior to starting fieldwork. Please let us know your availability.

Pursuant to 36 C.F.R. Part 800.3(f)(2), TVA is initiating consultation with federally recognized Indian tribes regarding historic properties within the proposed project's APE that may be of religious and cultural significance and are eligible for the National Register of Historic Places.

Mr. Barry White Page 2 April 13, 2021

Please contact Michaelyn Harle by email at <a href="mailto:mharle@tva.gov">mharle@tva.gov</a> with your comments.

Sincerely,

Clinton E. Jones

Manager

Cultural Compliance

MSH:ABM Enclosures

## INTERNAL COPIES NOT TO BE INCLUDED WITH OUTGOING LETTER:

S. Dawn Booker, BR 2C-C
J. Taylor Cates, BR 2C-C
Michael C. Easley, BR 2C-C
Michaelyn S. Harle, WT 11C-K
Brandon J. Hartline, BR 2C-C
Susan R. Jacks, WT 11C-K
Dana M. Nelson, WT 11C-K
Rebecca C. Tolene, WT 7B-K
William B. Wells, III BR 2C-C
W. Douglas White, WT 11C-K
ECM, ENVRecords



400 West Summit Hill Drive, Knoxville, Tennessee 37902

January 28, 2022

Mr. Barry White Director Mississippi Department of Archives and History Historic Preservation Division Post Office Box 571 Jackson, Mississippi 39205-0521

Dear Mr. White:

TENNESSEE VALLEY AUTHORITY (TVA), POWER PURCHASE AGREEMENT (PPA), GOLDEN TRIANGLE SOLAR, SECOND PHASE, LOWNDES COUNTY, MISSISSIPPI (TVA TRACKING NUMBER – CID 79857) (33.39170, -88.61037) (MDAH PROJECT LOG #04-063-21)

In a letter dated March 11, 2021, TVA initiated consultation with your office regarding TVA's proposal to enter into a PPA with Origis Energy for an approximately 150 megawatt (MW) solar photovoltaic (PV) generating facility and a 50 MW Battery Energy Storage System (BESS) in Lowndes County, Mississippi. The project would be located within seven parcels totaling 1,520 acres.

TVA, in consultation, determined that the area of potential effects (APE) to be the area of proposed ground disturbance, where physical effects could occur, including the PV facility, associated access routes, interconnect, and the BESS, as well as areas within a half-mile radius of the project within which the project would be visible, where visual effects on aboveground resources could occur. To obtain maximum flexibility in design to avoid both cultural and environmental resources the archaeological survey area consisted of the entire 1,520 acres.

Origis Energy contracted with Burns and McDonald to conduct the Phase I Cultural Resources survey in accordance with the scope of work that was provided to your office. The resulting report titled, *Phase 1 Cultural Resources Survey for the Golden Triangle II Solar Farm Project, Lowndes County, Mississippi* can be downloaded.

Please note that a portion of the survey area was included in the original Phase I survey, as described in the Golden Triangle I addendum report we provided previously (*Phase 1 Cultural Resources Survey for the Golden Triangle I Solar Farm Project, Lowndes County, Mississippi Addendum Report*).

Mr. Barry White Page 2 January 28, 2022

Burns and McDonald identified 12 archaeological sites and 15 isolated finds as a result of the current survey.

- Site 22LO1100, is a Late Archaic through Late Woodland period possible camp site. Burns and McDonald recommends site 22LO1100 remain unassessed for inclusion in the National Register of Historic Places (NRHP); the phase I survey indicated the potential presence of intact deposits, though no such deposits were identified.
- Sites 22LO1114 and 22LO1098 are historic scatters with intact features. Burns and McDonald recommend both 22LO1114 and 22LO1098 remain unassessed for inclusion in the NRHP.
- Site 22LO1117 is an extant portion of the Smith Oaks Harness Horse Racing Track. The 0.5-mile dirt track was constructed by Thomas Wilburn II in the late 1940s or early 1950s. No artifacts were identified in association with the racetrack. Burns and McDonald recommend 22LO1117 eligible for inclusion in the NRHP under Criteria A and B.
- The remaining archaeological sites represent late nineteenth to early twentieth century
  artifact scatters. While some of these sites may represent tenant or sharecropper
  occupations, few artifacts were recovered, no intact deposits were identified, and the sites
  have been affected by ground disturbance. TVA finds that the sites do not retain sufficient
  integrity to be considered eligible for listing on the NRHP.

The survey resulted in the documentation of 25 architectural resources on seven properties within the visual APE. An additional eight architectural resources (Resources 15, 41a, 41b, 42, 43, and 44a-44c) were previously evaluated as part of the Golden Triangle I Solar Project and were determined, in consultation with your office, not eligible for NRHP listing (Shaver et. al 2021). Two properties were previously identified during the additional parcel in the addendum report for Golden Triangle I.

- Resource 02 (46 in the addendum report) is a collection of structures associated with the Smith Oaks Plantation. Burns and McDonald recommend, and TVA agrees, that one of the structures, a circa 1880s former commissary/store building, be considered eligible for the NRHP. As sufficient analysis of the Smith Oaks Cemetery has not been done, the cemetery should be considered unassessed for NRHP eligibility. Per our discussion in our December consultation letter for Golden Triangle I, TVA finds that both resources' integrity of setting has been reduced due to the loss of the original dwelling and associated agricultural buildings. TVA finds that although there would be a visual effect to the Smith Oakes Commissary and Cemetery the effect would not be adverse.
- Resource 07 (Resource 47 in the addendum report), a former African American cemetery
  identified through archival information, was not identified during the field study. The
  cemetery warrants additional NRHP eligibility analysis and should be considered
  unassessed. The cemetery is located outside the project footprint. The viewshed to the
  project area is obscured by existing pond berms and the undertaking would not have an
  effect to this property should it be determined eligible.
- Resource 05 is a collection of structures associated with the former Oakland Plantation.
   Oakland Plantation includes a circa 1840s dwelling (05a), a circa 1975 barn (05b), two small

Mr. Barry White Page 3 January 28, 2022

early-twentieth-century buildings (05c and 05d) located south of the dwelling, and a non-historic-age building behind the dwelling that appears to be a secondary residence. Burns and McDonald recommends Resource 05a eligible for NRHP listing under Criteria A and C, and Resources 05c and 05d are recommended as contributing resources. Resource 5 is located approximately 0.31 miles outside of the project footprint and would not be directly affected by the undertaking. The viewshed of Resource 05a, c, and d is heavily obscured by dense vegetation and would not have an adverse visual effect to historic properties.

 Burns and McDonald recommends Resources 01, 03, 04, and 06 not eligible for the NRHP due to lack of architectural distinction, inability to associate the property with significant events or persons, and/or architectural integrity.

TVA has read the enclosed report and agrees with the recommendations of the authors. In order to avoid adverse effects to unassessed sites (22LO1100, 22LO1114, 22LO1098, 22LO1117), the boundaries of the sites, along with a 100-foot buffer, have been added to the "exclusion areas" within the project area. TVA will avoid these exclusion areas and will not allow, in these areas, any development, disturbance, or other construction activities associated with the development of the project or future activities associated with the operation and maintenance of the solar array. In order to ensure avoidance of these sites during the life of the project, Origis and TVA will sign the attached draft legal agreement.

Pursuant to 36 CFR Part 800.3(f)(2), TVA is consulting with federally recognized Indian tribes regarding properties within the survey area that may be of religious and cultural significance to them and eligible for the NRHP.

Pursuant to 36 CFR Part 800.5(c) we are notifying you of TVA's finding of no adverse effect, providing the documentation specified in § 800.11(e); and inviting you to review the finding. Also, we are seeking your agreement with TVA's eligibility determinations and finding that the undertaking as currently planned will have no adverse effects on historic properties.

If you have any questions, please contact Michaelyn Harle by email, mharle@tva.gov.

Sincerely,

James W. Osborne, Jr.

Manager

**Cultural Compliance** 

MSH:ABM Enclosures


## **References Cited:**

Shaver, Douglas, Brandy Harris, Kim House, and Jess Kepka

2021 Phase I Cultural Resources Survey for the Proposed Golden Triangle Solar Farm Project, Lowndes County, Mississippi. Prepared by Burns and McDonnell Engineering Company, Inc., Kansas City, Missouri, for Origis Energy.

# INTERNAL COPIES NOT TO BE INCLUDED WITH OUTGOING LETTER:

S. Dawn Booker, BR 2C-C
J. Taylor Cates, WT 11C-K
Michael C. Easley, BR 2C-C
Michaelyn S. Harle, WT 11C-K
Brandon J. Hartline, BR 2C-C
Susan R. Jacks, WT 11C-K
Dana M. Nelson, WT 11B-K
Rebecca C. Tolene, WT 11C-K
William B. Wells, III BR 2C-C
W. Douglas White, WT 11C-K
ECM, ENVRecords





February 17, 2022

Mr. James W. Osborne, Jr, Tennessee Valley Authority 400 West Summit Hill Drive Knoxville, Tennessee 37902

RE: Tennessee Valley Authority (TVA), Origis Energy Solar (Golden Triangle I) (33.45111, -88.62775) (TVA Tracking Number – CIDV 79483). MDAH Project Log 02-133-22 (12-139-21, 05-135-20), Lowndes County, Mississippi

Dear Mr. Osborne:

We have received your February 15, 2022 communication regarding the Golden Triangle II Solar Farm Project, Lowndes County, Mississippi in accordance with our responsibilities under Section 106 of the National Historic Preservation Act and 36 CFR Part 800.

MDAH concurs with the archaeological and architectural resource eligibility determinations and the finding that the undertaking as currently planned will have no adverse effects on historic properties. Additionally, MDAH concurs with the proposed temporary barriers to avoid effects to significant archaeological sites. Please consider this comment letter as conclusion of the Mississippi SHPO's consultation requirements for this undertaking.

If you have any questions or require additional information, please do not hesitate to contact me.

Sincerely,

Barry White

Deputy State Historic Preservation Officer

FOR: Katie Blount

Bany white

State Historic Preservation Officer

Cc: Michaelyn Harle – TVA

Amy McCampbell - TVA



P.O. Box 571 Jackson, MS 39205-0571 601-576-6850 mdah.ms.gov

May 27, 2022

Mr. James W. Osborne, Jr. Tennessee Valley Authority 400 West Summit Hill Drive Knoxville, Tennessee 37902

RE: Phase I Cultural Resources Survey for the Proposed Golden Triangle II Solar

Farm Project, Revised, (TVA) MDAH Project Log #04-162-22, Report #22-0162,

Lowndes County

Dear Mr. Osborne:

We have reviewed the March, 2022, cultural resources survey, by Douglas Shaver and Brandy Harris, Principal Investigators, with Burns & McDowell Engineering Company, Inc., received on April 27, 2022, for the above referenced project in accordance with our responsibilities under Section 106 of the National Historic Preservation Act and 36 CFR Part 800. After review, MDAH archaeology section concurs with avoidance plans for sites 22Lo1114, 1115, 1116 and 1119, and that the project will have No Adverse Effect on these sites.

MDAH architectural section concludes that the multiple resources in the APE which are eligible for listing under Criterion A: Agriculture and Criterion C: Architecture, will not be adversely affected by this project.

As such, we have no objections with the proposed undertaking.

There remains the possibility that unrecorded cultural resources may be encountered during the project. Should this occur, we would appreciate your contacting this office immediately in order that we may offer appropriate comments under 36 CFR 800.13.

If you have any questions, please do not hesitate to contact us at (601) 576-6940.

Sincerely.

Hal Bell

Review and Compliance Officer

Hal Bell

FOR: Katie Blount

State Historic Preservation Officer



# United States Department of the Interior



### FISH AND WILDLIFE SERVICE

Mississippi Ecological Services Field Office 6578 Dogwood View Parkway, Suite A Jackson, MS 39213-7856

Phone: (601) 965-4900 Fax: (601) 965-4340 http://www.fws.gov/mississippiES/endsp.html

In Reply Refer To: March 03, 2022

Project code: 2022-0015575 Project Name: Golden Triangle

Subject: Verification letter for the 'Golden Triangle' project under the January 5, 2016,

Programmatic Biological Opinion on Final 4(d) Rule for the Northern Long-eared Bat

and Activities Excepted from Take Prohibitions.

#### Dear Elizabeth Hamrick:

The U.S. Fish and Wildlife Service (Service) received on March 03, 2022 your effects determination for the 'Golden Triangle' (the Action) using the northern long-eared bat (*Myotis septentrionalis*) key within the Information for Planning and Consultation (IPaC) system. This IPaC key assists users in determining whether a Federal action is consistent with the activities analyzed in the Service's January 5, 2016, Programmatic Biological Opinion (PBO). The PBO addresses activities excepted from "take" prohibitions applicable to the northern long-eared bat under the Endangered Species Act of 1973 (ESA) (87 Stat.884, as amended; 16 U.S.C. 1531 et seq.).

Based upon your IPaC submission, the Action is consistent with activities analyzed in the PBO. The Action may affect the northern long-eared bat; however, any take that may occur as a result of the Action is not prohibited under the ESA Section 4(d) rule adopted for this species at 50 CFR §17.40(o). Unless the Service advises you within 30 days of the date of this letter that your IPaC-assisted determination was incorrect, this letter verifies that the PBO satisfies and concludes your responsibilities for this Action under ESA Section 7(a)(2) with respect to the northern long-eared bat.

Please report to our office any changes to the information about the Action that you submitted in IPaC, the results of any bat surveys conducted in the Action area, and any dead, injured, or sick northern long-eared bats that are found during Action implementation. If the Action is not completed within one year of the date of this letter, you must update and resubmit the information required in the IPaC key.

This IPaC-assisted determination allows you to rely on the PBO for compliance with ESA Section 7(a)(2) <u>only</u> for the northern long-eared bat. It **does not** apply to the following ESA-protected species that also may occur in the Action area:

- Monarch Butterfly Danaus plexippus Candidate
- Prices Potato-bean *Apios priceana* Threatened
- Wood Stork Mycteria americana Threatened

If the Action may affect other federally listed species besides the northern long-eared bat, a proposed species, and/or designated critical habitat, additional consultation between you and this Service office is required. If the Action may disturb bald or golden eagles, additional coordination with the Service under the Bald and Golden Eagle Protection Act is recommended.

[1] Take means to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct [ESA Section 3(19)].

## **Action Description**

You provided to IPaC the following name and description for the subject Action.

#### 1. Name

Golden Triangle

## 2. Description

The following description was provided for the project 'Golden Triangle':

The Tennessee Valley Authority (TVA) has entered into a Power Purchase Agreement with MS Solar 6, LLC to purchase power and environmental attributes generated by the proposed Golden Triangle 2 Solar Project in Lowndes County, Mississippi. The project would be constructed by MS Solar 6 and is expected to generate up to 150 megawatts (MW) of alternating current (AC) capacity with a 50 MW AC – 200-Megawatt hour Battery Energy Storage System. Total land impacts for implementation of the project, or proposed action, would be less than the overall project site (approx. 1,525 acres). Up to 270 of the 480 acres of forest on site may be removed.

Approximate location of the project can be viewed in Google Maps: <a href="https://www.google.com/maps/@33.39304685">https://www.google.com/maps/@33.39304685</a>,-88.61355841205881,14z



## **Determination Key Result**

This Federal Action may affect the northern long-eared bat in a manner consistent with the description of activities addressed by the Service's PBO dated January 5, 2016. Any taking that may occur incidental to this Action is not prohibited under the final 4(d) rule at 50 CFR §17.40(o). Therefore, the PBO satisfies your responsibilities for this Action under ESA Section 7(a)(2) relative to the northern long-eared bat.

### Determination Key Description: Northern Long-eared Bat 4(d) Rule

This key was last updated in IPaC on May 15, 2017. Keys are subject to periodic revision.

This key is intended for actions that may affect the threatened northern long-eared bat.

The purpose of the key for Federal actions is to assist determinations as to whether proposed actions are consistent with those analyzed in the Service's PBO dated January 5, 2016.

Federal actions that may cause prohibited take of northern long-eared bats, affect ESA-listed species other than the northern long-eared bat, or affect any designated critical habitat, require ESA Section 7(a)(2) consultation in addition to the use of this key. Federal actions that may affect species proposed for listing or critical habitat proposed for designation may require a conference under ESA Section 7(a)(4).

# **Determination Key Result**

This project may affect the threatened Northern long-eared bat; therefore, consultation with the Service pursuant to Section 7(a)(2) of the Endangered Species Act of 1973 (87 Stat.884, as amended; 16 U.S.C. 1531 et seq.) is required. However, based on the information you provided, this project may rely on the Service's January 5, 2016, *Programmatic Biological Opinion on Final 4(d) Rule for the Northern Long-Eared Bat and Activities Excepted from Take Prohibitions* to fulfill its Section 7(a)(2) consultation obligation.

# **Qualification Interview**

1. Is the action authorized, funded, or being carried out by a Federal agency? *Yes* 

2. Have you determined that the proposed action will have "no effect" on the northern long-eared bat? (If you are unsure select "No")

No

3. Will your activity purposefully **Take** northern long-eared bats?

No

4. [Semantic] Is the project action area located wholly outside the White-nose Syndrome Zone?

#### Automatically answered

No

5. [Semantic] Is the project action area located within 0.25 miles of a known northern longeared bat hibernaculum?

Note: The map queried for this question contains proprietary information and cannot be displayed. If you need additional information, please contact your State wildlife agency

#### Automatically answered

No

6. [Semantic] Is the project action area located within 150 feet of a known occupied northern long-eared bat maternity roost tree?

Note: The map queried for this question contains proprietary information and cannot be displayed. If you need additional information, please contact your State wildlife agency

## Automatically answered

No

# **Project Questionnaire**

If the project includes forest conversion, report the appropriate acreages below. Otherwise, type '0' in questions 1-3.

1. Estimated total acres of forest conversion:

270

2. If known, estimated acres of forest conversion from April 1 to October 31

135

3. If known, estimated acres of forest conversion from June 1 to July 31

135

# If the project includes timber harvest, report the appropriate acreages below. Otherwise, type '0' in questions 4-6.

4. Estimated total acres of timber harvest

0

5. If known, estimated acres of timber harvest from April 1 to October 31

0

6. If known, estimated acres of timber harvest from June 1 to July 31

0

# If the project includes prescribed fire, report the appropriate acreages below. Otherwise, type '0' in questions 7-9.

7. Estimated total acres of prescribed fire

0

8. If known, estimated acres of prescribed fire from April 1 to October 31

n

9. If known, estimated acres of prescribed fire from June 1 to July 31

0

# If the project includes new wind turbines, report the megawatts of wind capacity below. Otherwise, type '0' in question 10.

10. What is the estimated wind capacity (in megawatts) of the new turbine(s)?

0

# **IPaC User Contact Information**

Agency: Tennessee Valley Authority

Name: Elizabeth Hamrick Address: 400 W Summit Hill Dr

City: Knoxville

State: TN Zip: 37902

Email ecburton@tva.gov

Phone: 5034492373



400 West Summit Hill Drive, Knoxville, Tennessee 37902

March 7, 2022

Mr. Stephen Ricks U.S. Fish and Wildlife Service Ecological Services 6578 Dogwood View Parkway, Suite A Jackson, Mississippi 39213

Dear Mr. Ricks:

TENNESSEE VALLEY AUTHORITY (TVA) - GOLDEN TRIANGLE II SOLAR and BATTERY ENERGY STORAGE SYSTEM (BESS) PROJECT

TVA has entered into a Power Purchase Agreement with MS Solar 6, LLC to purchase power generated by the proposed Golden Triangle II Solar and BESS Project in Lowndes County, Mississippi. The project would be constructed by MS Solar 6 and is expected to generate up to 150 megawatts (MW) of alternating current (AC) capacity with a 50 MW AC – 200-Megawatt hour BESS. Total land impacts for implementation of the project, or proposed action, would be less than the overall project site (Approximately 1,525 acres). Up to 270 of the 480 acres of forest on site may be removed. Wetlands and open water ponds would be avoided. The installation of temporary and permanent bridges and/or culverts could affect up to 300 linear feet of intermittent and perennial streams. Best Management Practices for stream crossings would be implemented, such as using a bridge span that avoids direct impacts to the stream beds and top of bank, placing geotextile fabric along bridges to minimize dirt and debris from entering the stream channel, and minimizing the width of each crossing to the smallest width allowable (within safety requirements). Please refer to the attached Protected Species Habitat Assessment Report (PSHAR) and the draft Environmental Assessment (https://www.tva.com/environment/environmental-stewardship/environmental-reviews/nepadetail/golden-triangle-ii-solar-project) for a more detailed project description, figures, and photos.

Review of the TVA Regional Natural Heritage database and the U.S. Fish and Wildlife Service Information for Planning and Consultation (IPaC) website identified four species listed as federally endangered, threatened, candidate, or delisted and monitored under the Endangered Species Act (ESA) that have the potential to occur within the project area in Lowndes County, Mississippi. These species include one plant (Price's potato-bean), one bird (wood stork), one mammal (northern long-eared bat [NLEB]) and one insect (monarch butterfly) that have the potential to occur within Lowndes County based on historic range, proximity to known occurrence records, biological characteristics, and/or physiographic characteristics. No federally designated critical habitats for these species are present within or adjacent to the project action area, therefore no adverse modification of critical habitats would occur. Section 7 consultation is not required for candidate species.

Mr. Stephen Ricks Page 2 March 7, 2022

Field surveys were conducted by biologists from Burns and McDonnell Engineering Company, Inc from March 8 - 13, 2021, to determine whether suitable habitat for federally listed species occurs within the project action area. Suitable roosting habitat for the wood stork does not exist within the survey area for the project. However, suitable foraging habitat may be present near open water. There are no large, inundated wetlands within the Project Area. The project footprint includes 20 small open waters (PUB) and several small emergent wetland areas (collectively totaling approximately 26 acres). Proposed panel layouts avoid any impacts on wetlands. There are also large aquaculture/fish farms directly north of the project that may attract foraging wood storks. The project would not affect fish farms or large open waters outside the immediate project limits. Therefore, TVA has determined there would be no effect to wood stork or wood stork habitat resulting from implementation of the proposed action.

No individual Price's potato-bean were observed during field surveys; however, potential habitat for the species was observed within the survey area. Potential habitat for the species occurs in ravine areas that would not be developed during project construction or operation. Thus, TVA has determined there would be no effect on Price's potato-bean.

There are no known records documenting the presence of NLEB from within Lowndes County, Mississippi. No known hibernacula or maternity roosts occur within five miles of the project site. Phase 1 Bat Habitat Assessments were conducted from March 8 - 13, 2021, using the 2019 and 2020 Range-Wide Indiana Bat Summer Survey Guidelines as a surrogate protocol for determining presence/absence of the federally threatened NLEB. No caves, mines, buildings, bridges or potential winter roosting structures were identified during field surveys of the project footprint. Suitable summer roosting habitat for the NLEB was observed within forested areas at the project site, and suitable foraging habitat was observed within stream corridors, fence rows, wetlands, and forests throughout the project site. As mentioned above, the project footprint includes approximately 26 acres of small open waters and small wetlands. Proposed panel layouts avoid any impacts to wetlands. The installation of temporary and permanent bridges and/or culverts could affect up to 300 linear feet of intermittent and perennial streams. Best Management Practices would be used around these bodies of water, minimizing sedimentation and avoiding changes to hydrology.

Up to 270 acres of forest that could provide summer roosting habitat for NLEBs may be removed. Phase 2 Presence/Absence surveys were conducted by Burns & McDonnell on August 3 through August 10, 2021 to assess potential occurrence of the NLEB using acoustic detectors. Proposed plans were submitted to USFWS on July 29, 2021. Acoustic surveys were conducted at 11 sites (a total of 26 detector-nights). Using the Kaleidoscope Pro classifier, no threatened or endangered species were determined to be present. No manual vetting was required. See attached Acoustic Survey Report. Despite these results, efforts would still be made to minimize clearing of forested areas to the greatest extent practicable as the Project designs progress.

TVA has determined that while removal of suitable roosting habitat could have indirect adverse effects on NLEB and result in 'incidental take' as defined in the ESA, this 'incidental take' is excepted from ESA Section 9 Take Prohibitions. In addition, presence/absence surveys

Mr. Stephen Ricks Page 3 March 7, 2022

determined that the likelihood of presence of this species in the project area is low. Based on the USFWS' online Northern Long-Eared Bat 4(d) rule determination key accessed on March 3, 2022 (see verification letter attached, Project Code: 2022-0015575), this project may affect NLEB and may rely on the Service's January 5, 2016, Programmatic Biological Opinion (PBO) on Final 4(d) Rule for the Northern Long-Eared Bat and Activities Excepted from Take Prohibitions to fulfill its Section 7(a)(2) consultation obligation.

We respectfully request concurrence with our determination. However, it is our understanding that unless the Service advises us within 30 days of March 3, 2022, when the NLEB verification letter was generated, that our IPaC-assisted determination was incorrect, that the verification letter adequately verifies that the PBO satisfies and concludes our responsibilities for this Action under ESA Section 7(a)(2) with respect to the NLEB.

Should you have any questions or wish to discuss the project in more detail, please contact Elizabeth Hamrick at ecburton@tva.gov.

Sincerely,

W. Douglas White

Manager

Biological Compliance

Will-Dholes

JCT:ABM Enclosures

# INTERNAL COPIES, NOT INCLUDED WITH OUTBOUND LETTER:

Adam J. Dattilo, WT 11C-K Elizabeth B. Hamrick, WT 11C-K Susan R. Jacks, WT 11 D-K J. Taylor Johnson, BR 2C-C W. Douglas White, WT11C-K ECM, ENVRecords



# **United States Department of the Interior**

## FISH AND WILDLIFE SERVICE

Mississippi Ecological Services Field Office 6578 Dogwood View Parkway, Suite A Jackson, Mississippi 39213 Phone: (601)965-4900 Fax: (601)965-4340



11 March 2022

IN REPLY REFER TO: 2022-0017318

W. Douglas White Tennessee Valley Authority 400 West Summit Hill Drive Knoxville, Tennessee 37902


Dear Mr. White:

The Fish and Wildlife Service (Service) has reviewed your correspondence dated March 7, 2022 regarding the proposed Golden Triangle II Solar and Battery Energy Storage System (BESS) Project, located in Lowndes County, Mississippi. Our comments are submitted in accordance with the Endangered Species Act (ESA) (87 Stat. 884, as amended; 16 U.S.C. 1531 et seq.).

The Service has received the Northern Long-eared Bat (NLEB) Verification Letter and concurs with the determination that the proposed project may affect the NLEB, but that any resulting incidental take of the NLEB is not prohibited by the final 4(d) rule. We have also reviewed your "no effect" determination for the wood stork and Price's potato bean and offer no additional comments regarding your determination of effects for these species. No further coordination is required with this office unless there are changes in scope or location of the proposed project.

If you have any questions, please contact Ashley Seagroves in our office, telephone: (601) 321-1126, email: lauren\_seagroves@fws.gov or visit our website at https://www.fws.gov/office/mississippi-ecological-services.

Sincerely,
James A. Austin
Acting Field Supervisor
Mississippi Field Office



### **Enclosure**

EPA comments on the Draft Environmental Assessment for the Tennessee Valley Authority Golden Triangle II
Battery Energy Storage System in Lowndes County, Mississippi

<u>Air Quality</u>: The DEA identified minor direct impacts to air quality over the 17 months of construction and the operation of the project site. Also, temporary effects on Green House Gas emissions are expected during construction.

Recommendation: The EPA recommends considering the following additional measures: Clean Diesel: implementation of diesel controls, cleaner fuel, and cleaner construction practices for on-road and off-road equipment used for transportation, soil movement, or other construction activities, including:

- Strategies and technologies that reduce unnecessary idling, including auxiliary power units, the use of electric equipment, and strict enforcement of idling limits; and
- Use of Clean Diesel through add-on control technologies like diesel particulate filters and diesel oxidation catalysts, repowers, or newer, cleaner equipment. For more information on diesel emission controls in construction projects, please see: <a href="https://northeastdiesel.org/construction.html">https://northeastdiesel.org/construction.html</a>

<u>Water Resources:</u> Section 3.3.2.2 of the DEA identified that the Proposed Action would result in 160 linear feet of minor, direct, permanent impacts to jurisdictional streams over eight stream crossings at locations where permanent culverts will be installed for construction of access roads.

Recommendation: The EPA notes that a portion of the 160 linear feet of jurisdictional intermittent and perennial stream impacts is subject to regulatory oversight of the U.S. Army Corps of Engineers, Mississippi Department for Environmental Quality. The EPA recommends that any contractor working on-site use best management practices. The EPA also recommends that site grading, excavation, and construction plans include implementable measures to prevent erosion and sediment runoff from the project site during and after construction.